

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | | |
|---|--|---|--|
| <p>(51) International Patent Classification ⁶ : A61K 38/17, 31/70, 39/395, C12N 15/11, C12Q 1/68, G01N 33/50, 33/574, A01K 67/027, C12N 15/00</p> | | <p>A2</p> | <p>(11) International Publication Number: WO 98/35693</p> <p>(43) International Publication Date: 20 August 1998 (20.08.98)</p> |
| <p>(21) International Application Number: PCT/IB98/00781</p> <p>(22) International Filing Date: 13 February 1998 (13.02.98)</p> <p>(30) Priority Data: 08/800,929 13 February 1997 (13.02.97) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 08/800,929 (CIP) Filed on 13 February 1997 (13.02.97)</p> <p>(71) Applicant (for all designated States except US): UNIVERSITY OF OTTAWA [CA/CA]; 650 Cumberland, Ottawa, Ontario K1N 6N5 (CA).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): KORNELUK, Robert [CA/CA]; 1901 Tweed Avenue, Ottawa, Ontario K1G 2L8 (CA). MACKENZIE, Alexander, E. [CA/CA]; 35 Rockcliffe Way, Ottawa, Ontario K1M 1A3 (CA). LISTON, Peter [CA/CA]; 1 Second Avenue, Ottawa, Ontario K1S 2H2 (CA). BAIRD, Stephen [CA/CA]; 20 Julian Avenue, Ottawa, Ontario K1Y 0S5 (CA). TSANG, Benjamin [CA/CA];</p> | | <p>1053 Carling Avenue, Ottawa, Ontario K1Y 4E9 (CA). PRATT, Christine [CA/CA]; 31 Long Gate Court, Nepean, Ontario K2J 4E7 (CA).</p> <p>(74) Agent: DEETH WILLIAMS WALL; National Bank Building, Suite 400, 150 York Street, Toronto, Ontario M5H 3S5 (CA).</p> <p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published Without international search report and to be republished upon receipt of that report.</p> | |
| <p>(54) Title: DETECTION AND MODULATION OF THE IAPS AND NAIP FOR THE DIAGNOSIS AND TREATMENT OF PROLIFERATIVE DISEASE</p> <p>(57) Abstract</p> <p>Disclosed are diagnostic and prognostic methods and kits for the detection and treatment of proliferative diseases such as cancer (e.g., ovarian cancer, breast cancer, and lymphoma). Also disclosed are therapeutics for treating proliferative diseases (and methods for identifying such therapeutics) that utilize IAP and NAIP antisense nucleic acid molecules, antibodies which specifically bind IAP and NAIP polypeptides, and compounds that reduce the biological activities of IAP and NAIP polypeptides.</p> | | | |

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | | | |
|----|--------------------------|----|--|----|--|----|--------------------------|
| AL | Albania | ES | Spain | LS | Lesotho | SI | Slovenia |
| AM | Armenia | FI | Finland | LT | Lithuania | SK | Slovakia |
| AT | Austria | FR | France | LU | Luxembourg | SN | Senegal |
| AU | Australia | GA | Gabon | LV | Latvia | SZ | Swaziland |
| AZ | Azerbaijan | GB | United Kingdom | MC | Monaco | TD | Chad |
| BA | Bosnia and Herzegovina | GE | Georgia | MD | Republic of Moldova | TG | Togo |
| BB | Barbados | GH | Ghana | MG | Madagascar | TJ | Tajikistan |
| BE | Belgium | GN | Guinea | MK | The former Yugoslav Republic of Macedonia | TM | Turkmenistan |
| BF | Burkina Faso | GR | Greece | ML | Mali | TR | Turkey |
| BG | Bulgaria | HU | Hungary | MN | Mongolia | TT | Trinidad and Tobago |
| BJ | Benin | IE | Ireland | MR | Mauritania | UA | Ukraine |
| BR | Brazil | IL | Israel | MW | Malawi | UG | Uganda |
| BY | Belarus | IS | Iceland | MX | Mexico | US | United States of America |
| CA | Canada | IT | Italy | NE | Niger | UZ | Uzbekistan |
| CF | Central African Republic | JP | Japan | NL | Netherlands | VN | Viet Nam |
| CG | Congo | KE | Kenya | NO | Norway | YU | Yugoslavia |
| CH | Switzerland | KG | Kyrgyzstan | NZ | New Zealand | ZW | Zimbabwe |
| CI | Côte d'Ivoire | KP | Democratic People's Republic of Korea | PL | Poland | | |
| CM | Cameroon | KR | Republic of Korea | PT | Portugal | | |
| CN | China | KZ | Kazakhstan | RO | Romania | | |
| CU | Cuba | LC | Saint Lucia | RU | Russian Federation | | |
| CZ | Czech Republic | LI | Liechtenstein | SD | Sudan | | |
| DE | Germany | LK | Sri Lanka | SE | Sweden | | |
| DK | Denmark | LR | Liberia | SG | Singapore | | |
| EE | Estonia | | | | | | |

DETECTION AND MODULATION OF THE IAPs AND NAIP FOR THE DIAGNOSIS AND TREATMENT OF PROLIFERATIVE DISEASE

5 Background of the Invention

The invention relates to the diagnosis and treatment of proliferative disease, in particular, cancer.

One mechanism by which cells die is referred to as apoptosis, or programmed cell death. Apoptosis often occurs as a normal part of the development and maintenance of healthy tissues, and is now known to play a critical role in embryonic development. The failure of a normal apoptotic response has been implicated in the development of cancer; autoimmune disorders, such as lupus erythematosus and multiple sclerosis; and in viral infections, including those associated with herpes virus, poxvirus, and adenovirus.

Compared to the numerous growth promoting genes identified to date (>100) relatively few genes have been isolated that regulate apoptosis. Baculoviruses encode proteins termed inhibitors of apoptosis proteins (IAPs) which inhibit the apoptosis that would otherwise occur when insect cells are infected by the baculovirus. The baculovirus IAP genes include sequences encoding a ring zinc finger-like motif (RZF), which is presumed to be directly involved in DNA binding, and two N-terminal domains that consist of a 70 amino acid repeat motif termed a BIR domain (Baculovirus IAP Repeat). Mammalian IAP family members, and a related anti-apoptotic polypeptide, NAIP, have recently been identified.

Both normal cell types and cancer cell types display a wide range of susceptibility to apoptotic triggers. Many normal cell types undergo temporary growth arrest in response to a sub-lethal dose of radiation or cytotoxic chemical, while cancer cells in the vicinity undergo apoptosis. This provides the crucial treatment "window" of appropriate toxicity that allows successful anti-cancer therapy. It is therefore not surprising that resistance of tumor cells to apoptosis is emerging as a major category of cancer treatment failure. Finding compounds which overcome or prevent this resistance would greatly improve cancer therapies.

Summary of the Invention

30 We have discovered that IAP and NAIP overexpression are specifically associated
with a wide range of cancer types including ovarian cancer, adenocarcinoma, lymphoma, and

- 2 -

pancreatic cancer. The presence of a fragmented IAP polypeptide in the nucleus, and an overexpression of an IAP polypeptide in the presence of a p53 mutation correlates with a cancer diagnosis, a poor prognosis, and a resistance to numerous chemotherapeutic cancer drugs. In addition, we have found that an therapeutic agent that reduces the biological activity of an IAP polypeptide will induce apoptosis in a cell expressing the polypeptide (*e.g.*, a cell that is proliferating in a proliferative disease). These discoveries provide diagnostic and prognostic methods for the detection and treatment of proliferative diseases, and provide therapeutic compounds useful for the treatment of proliferative diseases, particularly cancer.

In a first aspect, the invention features a method for enhancing apoptosis in a cell from a mammal with a proliferative disease, the method including administering to the cell a compound that inhibits the biological activity of an IAP polypeptide or a NAIP polypeptide, the compound being administered to the cell in an amount sufficient to enhance apoptosis in the cell. In one embodiment of this aspect of the invention, the cell is proliferating in the proliferative disease. In another embodiment, the biological activity is the level of expression of the polypeptide (measured, for example, by assaying the amount of polypeptide present in the cell); the level of expression of an mRNA molecule encoding the polypeptide; or an apoptosis-inhibiting activity.

In various embodiments of the first aspect of the invention, the polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP. In other embodiment, the polypeptide is NAIP, XIAP, HIAP-1, or HIAP-2. In other preferred embodiments, the mammal is a human or a mouse, and the proliferative disease is cancer, for example, a cancer in a tissue selected from the group consisting of ovary, breast, pancreas, lymph node, skin, blood, lung, brain, kidney, liver, nasopharyngeal cavity, thyroid, central nervous system, prostate, colon, rectum, cervix, endometrium, and lung.

In various preferred embodiments of the first aspect of the invention, the compound is a negative regulator of an IAP or an NAIP-dependent anti-apoptotic pathway; a fragment of the IAP polypeptide, the fragment including a ring zinc finger and having no more than two BIR domains; a nucleic acid molecule encoding a ring zinc finger domain of the IAP polypeptide; a compound that prevents cleavage of the IAP polypeptide or the NAIP polypeptide; a purified antibody or a fragment thereof that specifically binds to the IAP polypeptide or the NAIP polypeptide; a ribozyme; or an antisense nucleic acid molecule have

- 3 -

a nucleic acid sequence that is complementary to the coding strand of a nucleic acid sequence encoding the IAP polypeptide or the NAIP polypeptide. Preferably, the cleavage is decreased by at least 20% in the cell; the antibody binds to a BIR domain of the IAP polypeptide or the NAIP polypeptide; the nucleic acid sequence encoding the IAP polypeptide or the NAIP polypeptide has about 50% or greater identity with the nucleotide sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, or the nucleic acid sequence of NAIP; the antisense nucleic acid molecule decreases the level of the nucleic acid sequence encoding the IAP polypeptide or the NAIP polypeptide by at least 20%, the level being measured in the cytoplasm of the cell; the antisense nucleic acid molecule is encoded by a virus vector; or the antisense nucleic acid molecule is encoded by transgene.

In a second aspect, the invention features a method for detecting a proliferative disease or an increased likelihood of the proliferative disease in a mammal that includes: (a) contacting an IAP or a NAIP nucleic acid molecule that is greater than about 18 nucleotides in length with a preparation of nucleic acid from a cell of the mammal, the cell proliferating in the disease, the cell from a tissue; and (b) measuring the amount of nucleic acid from the cell of the mammal that hybridizes to the molecule, an increase in the amount from the cell of the mammal relative to a control indicating an increased likelihood of the mammal having or developing a proliferative disease. In one embodiment, the method further includes the steps of: (a) contacting the molecule with a preparation of nucleic acid from the control, wherein the control is a cell from the tissue of a second mammal, the second mammal lacking a proliferative disease; and (b) measuring the amount of nucleic acid from the control, an increase in the amount of the nucleic acid from the cell of the mammal that hybridizes to the molecule relative to the amount of the nucleic acid from the control indicating an increased likelihood of the mammal having or developing a proliferative disease.

In one embodiment of the methods of the second aspect of the invention, the method further includes the steps of: (a) providing a pair of oligonucleotides having sequence identity to or being complementary to a region of the IAP or the NAIP nucleic acid molecule; (b) combining the pair of oligonucleotides with the nucleic acid under conditions suitable for polymerase chain reaction-mediated nucleic acid amplification; and (c) isolating the

- 4 -

amplified nucleic acid or fragment thereof. Preferably, the amplification is carried out using a reverse-transcription polymerase chain reaction (*e.g.*, RACE).

In one embodiment of the second aspect of the invention, the method provides measuring the nucleic acid having a nucleotide sequence that has about 50% or greater
5 identity with the nucleotide sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, or the nucleic acid sequence of NAIP. In other embodiments, the method provides measuring the nucleic acid having a nucleotide sequence that has about 50% or greater identity with the nucleotide sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, or NAIP.

10 In a third aspect, the invention features a method for detecting a proliferative disease or an increased likelihood of developing the disease in a mammal, the method including measuring the level of biological activity of an IAP polypeptide or a NAIP polypeptide in a sample of the mammal, an increase in the level of the IAP polypeptide or the NAIP polypeptide relative to a sample from a control mammal being an indication that the mammal
15 has the disease or increased likelihood of developing the disease. In various embodiments, the sample includes a cell that is proliferating in the disease from the mammal, the cell from a tissue; and the sample from a control mammal is from the tissue, the sample consisting of healthy cells. In another embodiment, the mammal and the control mammal are the same.

In various embodiments of the third aspect of the invention, the biological activity is
20 the level of expression of the polypeptide (measured, for example, by assaying the amount of the polypeptide present in the cell); wherein the biological activity is the level of expression of an mRNA molecule encoding the polypeptide; or wherein the biological activity is an apoptosis-inhibiting activity. In another embodiment, the polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP. In other
25 embodiments, the polypeptide is NAIP, XIAP, HIAP-1, or HIAP-2.

In a fourth aspect, the invention features a method for identifying a compound enhances apoptosis in an affected cell that is proliferating in a proliferative disease that includes exposing a cell that overexpresses an IAP polypeptide or a NAIP polypeptide to a candidate compound, a decrease the level of biological activity of the polypeptide indicating
30 the presence of a compound that enhances apoptosis in the affected cell that is proliferating in the proliferative disease.

- 5 -

In a fifth aspect, the invention features a method for identifying a compound that enhances apoptosis in an affected cell that is proliferating in a proliferative disease that includes the steps of: (a) providing a cell including a nucleic acid molecule encoding a IAP polypeptide or a nucleic acid molecule encoding a NAIP polypeptide, the nucleic acid molecule being expressed in the cell; and (b) contacting the cell with a candidate compound and monitoring level of biological activity of the IAP polypeptide or the NAIP polypeptide in the cell, a decrease in the level of biological activity of the IAP polypeptide or the NAIP polypeptide in the cell in response to the candidate compound relative to a cell not contacted with the candidate compound indicating the presence of a compound that enhances apoptosis in the affected cell that is proliferating in the proliferative disease. Preferably, the cell further expresses a p53 polypeptide associated with the proliferative disease.

In various embodiments of the fourth and fifth aspects of the invention, the biological activity is the level of expression of the polypeptide (measured, for example, by assaying the amount of the polypeptide present in the cell); wherein the biological activity is the level of expression of an mRNA molecule encoding the polypeptide; or wherein the biological activity is an apoptosis-inhibiting activity. In another embodiment, the polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP. In other embodiments, the polypeptide is NAIP, XIAP, HIAP-1, or HIAP-2.

In a sixth aspect, the invention features a method for determining the prognosis of a mammal diagnosed with a proliferative disease that includes the steps of: (a) isolating a sample from a tissue from the mammal; and (b) determining whether the sample has an increased an level of biological activity of an IAP polypeptide or an NAIP polypeptide relative to a control sample, an increase in the level in the sample being an indication that the mammal has a poor prognosis. In various embodiments of this aspect of the invention, the sample includes a cells that is proliferating in the proliferative disease and the control sample is from the tissue, the control sample consisting of healthy cells; and the sample and the control sample are from the mammal. Preferably, the sample further includes a cell expressing a p53 polypeptide associated with the proliferative disease.

In various embodiments of the sixth aspect of the invention, the biological activity is the level of expression of the polypeptide (measured, for example, by assaying the amount of the polypeptide present in the cell); wherein the biological activity is the level of expression

- 6 -

of an mRNA molecule encoding the polypeptide; or wherein the biological activity is an apoptosis-inhibiting activity. In another embodiment, the polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP. In other embodiments, the polypeptide is NAIP, XIAP, HIAP-1, or HIAP-2. In a preferred
5 embodiment, the level is assayed by measuring the amount of IAP peptide of less than 64 kDa present in the sample.

In a seventh aspect, the invention features a method for determining the prognosis of a mammal diagnosed with a proliferative disease that includes the steps of: (a) isolating a sample from the mammal, the sample having a nuclear fraction; and (b) measuring the
10 amount of a polypeptide that is recognized by an antibody that specifically binds an IAP polypeptide or an antibody that specifically binds an NAIP polypeptide in the nuclear fraction of the sample relative an amount from a control sample, an increase in the amount from the sample being an indication that the mammal has a poor prognosis. In preferred embodiments of this aspect of the invention, the sample is from a tissue of the mammal, the
15 sample including a cell that is proliferating in the proliferative disease, and the control sample is from the tissue, the control sample consisting of healthy cells. In another embodiment, the sample and the control sample are from the mammal.

In various embodiments of the seventh aspect of the invention, the biological activity is the level of expression of the polypeptide (measured, for example, by assaying the amount
20 of the polypeptide present in the cell); wherein the biological activity is the level of expression of an mRNA molecule encoding the polypeptide; or wherein the biological activity is an apoptosis-inhibiting activity. In another embodiment, the polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP. In other embodiments, the polypeptide is NAIP, XIAP, HIAP-1, or HIAP-2. In
25 another embodiment, the amount is measured by immunological methods.

In an eighth aspect, the invention features a method for treating a mammal diagnosed as having a proliferative disease that includes the steps of: (a) measuring the amount of an IAP or NAIP polypeptide in a first sample from a tissue from the mammal, the first sample including a cell that is proliferating in the proliferative disease; (b) measuring the amount of
30 the polypeptide in a second sample from the tissue, the second sample consisting of healthy cells; (c) detecting an increase in the amount of the polypeptide in the first sample to the

- 7 -

amount of the polypeptide in the second sample; and (d) treating the mammal with a compound that decreases the biological activity of the polypeptide. Preferably, the first sample and the second sample are from the mammal.

In various embodiments of the eighth aspect of the invention, the biological activity is the level of expression of the polypeptide (measured, for example, by assaying the amount of the polypeptide present in the cell); wherein the biological activity is the level of expression of an mRNA molecule encoding the polypeptide; or wherein the biological activity is an apoptosis-inhibiting activity. In another embodiment, the polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP. In other
10 embodiments, the polypeptide is NAIP, XIAP, HIAP-1, or HIAP-2.

In a ninth embodiment, the invention features the use of a compound that decreases the biological activity an IAP polypeptide or a NAIP polypeptide for the manufacture of a medicament for the enhancement of apoptosis.

In various embodiments of the ninth aspect of the invention, the biological activity is the level of expression of the polypeptide (measured, for example, by assaying the amount of the polypeptide present in the cell); wherein the biological activity is the level of expression of an mRNA molecule encoding the polypeptide; or wherein the biological activity is an apoptosis-inhibiting activity. In another embodiment, the polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP. In other
20 embodiments, the polypeptide is NAIP, XIAP, HIAP-1, or HIAP-2.

In a tenth aspect, the invention features a kit for diagnosing a mammal for the presence of a proliferative disease or an increased likelihood of developing a proliferative disease, the kit comprising an oligonucleotide that hybridizes to a nucleic acid sequence that encodes an IAP polypeptide or a NAIP polypeptide.

25 In various embodiments of the tenth aspect of the invention, the biological activity is the level of expression of the polypeptide (measured, for example, by assaying the amount of the polypeptide present in the cell); wherein the biological activity is the level of expression of an mRNA molecule encoding the polypeptide; or wherein the biological activity is an apoptosis-inhibiting activity. In another embodiment, the polypeptide is selected from the
30 group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP. In other embodiments, the polypeptide is NAIP, XIAP, HIAP-1, or HIAP-2.

- 8 -

In an eleventh aspect, the invention features a transgenic mammal, the mammal having an elevated level of biological activity of an IAP polypeptide or a NAIP polypeptide.

In various embodiments of the eleventh aspect of the invention, the biological activity is the level of expression of the polypeptide (measured, for example, by assaying the amount
5 of the polypeptide present in the cell); wherein the biological activity is the level of expression of an mRNA molecule encoding the polypeptide; or wherein the biological activity is an apoptosis-inhibiting activity. In another embodiment, the polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP. In other embodiments, the polypeptide is NAIP, XIAP, HIAP-1, or HIAP-2.

10 By "IAP gene" is meant a gene encoding a polypeptide having at least one BIR domain and is capable of modulating (inhibiting or enhancing) apoptosis in a cell or tissue when provided by other intracellular or extracellular delivery methods (see, *e.g.*, the U.S.S.N.s 08/511,485, 08/576,965, and PCT/IB96/01022). In preferred embodiments the IAP gene is a gene having about 50% or greater nucleotide sequence identity to at least one
15 of the IAP amino acid encoding sequences of Figs. 1-6 (SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, and SEQ ID NO: 13) or portions thereof, or has a ring zinc finger domain. Preferably, the region of sequence over which identity is measured is a region encoding at least one BIR domain and a ring zinc finger domain. Mammalian IAP genes include nucleotide sequences isolated from any mammalian source. Preferably, the
20 mammal is a human. The term "IAP gene" is meant to encompass any member of the family of genes that encode inhibitors of apoptosis. An IAP gene may encode a polypeptide that has at least 20%, preferably at least 30%, and most preferably at least 50% amino acid sequence identity with at least one of the conserved regions of one of the IAP members described herein (*i.e.*, either the BIR or ring zinc finger domains from the human or murine XIAP,
25 HIAP-1, or HIAP-2). Representative members of the IAP gene family include, without limitation, the human and murine XIAP, HIAP-1, or HIAP-2 genes.

By "a virus vector" is meant a functional or attenuated virus that is capable of delivering to a virus-infected cell a nucleic acid molecule. Preferably, the virus vector has been genetically engineered according to standard molecular biology techniques to bear a

- 9 -

heterologous nucleic acid molecule. Virus vectors include, without limitation, adenoviruses, retroviruses, baculoviruses, cytomegaloviruses (CMV), and vaccinia viruses.

By "IAP protein" or "IAP polypeptide" is meant a polypeptide, or fragment thereof, encoded by an IAP gene.

5 By "NAIP gene" and "NAIP polypeptide" is meant the NAIP genes, fragments thereof, and polypeptides encoded by the same described in UK9601108.5 filed January 19, 1996 and PCT Application No. PCT/IB97/00142 (claiming priority from UK9601108.5) filed January 17, 1997.

By "BIR domain" is meant a domain having the amino acid sequence of the
10 consensus sequence: Xaal-Xaal-Xaal-Arg-Leu-Xaal-Thr-Phe-Xaal-Xaal-Trp-Pro-Xaa2-Xaal-Xaal-Xaa2-Xaa2-Xaal-Xaal-Xaal-Xaal-Leu-Ala-Xaal-Ala-Gly-Phe-Tyr-Tyr-Xaal-Gly-Xaal-Xaal-Asp-Xaal-Val-Xaal-Cys-Phe-Xaal-Cys-Xaal-Xaal- Xaal-Xaal-Xaal-Xaal-Trp-Xaal-Xaal-Xaal-Asp-Xaal-Xaal-Xaal- Xaal-Xaal-His-Xaal-Xaal-Xaal-Xaal-Pro-Xaal-Cys-Xaal-Phe-Val, wherein Xaal is any amino acid and Xaa2 is any amino acid or is absent (SEQ ID
15 NO: 2). Preferably, the sequence is substantially identical to one of the BIR domain sequences provided for XIAP, HIAP-1, or HIAP-2 herein.

By "ring zinc finger" or "RZF" is meant a domain having the amino acid sequence of the consensus sequence: Glu-Xaal-Xaal-Xaal-Xaal-Xaal-Xaal-Xaa2-Xaal-Xaal-Xaal-Cys-Lys-Xaa3-Cys-Met-Xaal-Xaal-Xaal-Xaal-Xaal-Xaa3-Xaal-Phe-Xaal-Pro-Cys-Gly-His-Xaal-
20 Xaal-Xaal-Cys-Xaal-Xaal-Cys-Ala- Xaal-Xaal-Xaal-Xaal-Xaal-Cys-Pro-Xaal-Cys, wherein Xaal is any amino acid, Xaa2 is Glu or Asp, and Xaa3 is Val or Ile (SEQ ID NO:1).

Preferably, the sequence is substantially identical to the RZF domains provided herein for the human or murine XIAP, HIAP-1, or HIAP-2.

By "enhancing apoptosis" is meant increasing the number of cells which apoptose in
25 a given cell population. Preferably, the cell population is selected from a group including ovarian cancer cells, breast cancer cells, pancreatic cancer cells, T cells, neuronal cells, fibroblasts, or any other cell line known to proliferate in a laboratory setting. It will be appreciated that the degree of apoptosis enhancement provided by an apoptosis enhancing compound in a given assay will vary, but that one skilled in the art can determine the
30 statistically significant change in the level of apoptosis which identifies a compound which enhances apoptosis otherwise limited by an IAP. Preferably, "enhancing apoptosis" means

- 10 -

that the increase in the number of cells undergoing apoptosis is at least 25%, more preferably the increase is 50%, and most preferably the increase is at least one-fold. Preferably, the sample monitored is a sample of cells which normally undergo insufficient apoptosis (*i.e.*, cancer cells).

5 By "proliferative disease" is meant a disease which is caused by or results in inappropriately high levels of cell division, inappropriately low levels of apoptosis, or both. For example, cancers such as lymphoma, leukemia, melanoma, ovarian cancer, breast cancer, pancreatic cancer, and lung cancer are all examples of proliferative disease. A neoplasm (*i.e.*, any abnormal proliferation of cells, malignant or benign), is also a proliferative disease of the
10 invention.

By a "cell proliferating in a proliferative disease" is meant a cell whose abnormal proliferation contributes to the disease. Preferably, the cell expresses the antigen PCNA.

By "polypeptide" is meant any chain of more than two amino acids, regardless of post-translational modification such as glycosylation or phosphorylation.

15 By "IAP or NAIP biological activity" is meant any activity known to be caused *in vivo* or *in vitro* by a NAIP or an IAP polypeptide. Preferred biological activities of IAP and NAIP polypeptides are those described herein, and include, without limitation, a level of expression of the polypeptide that is normal for that cell type, a level of expression of the mRNA that is normal for that cell type, an ability to block apoptosis, and an ability to be
20 cleaved.

By a "compound that decreases the biological activity" is meant a compound that decreases any activity known to be caused *in vivo* or *in vitro* by a NAIP polypeptide or an IAP polypeptide. Preferred compounds include, without limitation, an antisense nucleic acid molecule that is complementary to the coding strand of nucleic acid molecule that encodes an
25 IAP or a NAIP polypeptide; an antibody, such as a neutralizing antibody, that specifically binds to an IAP or a NAIP polypeptide; and a negative regulator of an IAP or a NAIP polypeptide, such as a polypeptide fragment that includes the ring zing finger of an IAP polypeptide, a polypeptide fragment that has no more than two BIR domains, or nucleic acid molecules encoding these polypeptide fragments.

30 By "substantially identical" is meant a polypeptide or nucleic acid exhibiting at least 50%, preferably 85%, more preferably 90%, and most preferably 95% homology to a

- 11 -

reference amino acid or nucleic acid sequence. For polypeptides, the length of comparison sequences will generally be at least 16 amino acids; preferably at least 20 amino acids, more preferably at least 25 amino acids, and most preferably 35 amino acids. For nucleic acids, the length of comparison sequences will generally be at least 50 nucleotides, preferably at least 60 nucleotides, more preferably at least 75 nucleotides, and most preferably 110 nucleotides.

Sequence identity is typically measured using sequence analysis software with the default parameters specified therein (e.g., Sequence Analysis Software Package of the Genetics Computer Group, University of Wisconsin Biotechnology Center, 1710 University Avenue, Madison, WI 53705). This software program matches similar sequences by assigning degrees of homology to various substitutions, deletions, and other modifications. Conservative substitutions typically include substitutions within the following groups: glycine, alanine, valine, isoleucine, leucine; aspartic acid, glutamic acid, asparagine, glutamine; serine, threonine; lysine, arginine; and phenylalanine, tyrosine.

By "substantially pure polypeptide" is meant a polypeptide that has been separated from the components that naturally accompany it. Typically, the polypeptide is substantially pure when it is at least 60%, by weight, free from the proteins and naturally-occurring organic molecules with which it is naturally associated. Preferably, the polypeptide is an IAP polypeptide that is at least 75%, more preferably at least 90%, and most preferably at least 99%, by weight, pure. A substantially pure IAP polypeptide may be obtained, for example, by extraction from a natural source (e.g. a fibroblast, neuronal cell, or lymphocyte) by expression of a recombinant nucleic acid encoding an IAP polypeptide, or by chemically synthesizing the protein. Purity can be measured by any appropriate method, e.g., by column chromatography, polyacrylamide gel electrophoresis, or HPLC analysis.

A protein is substantially free of naturally associated components when it is separated from those contaminants which accompany it in its natural state. Thus, a protein which is chemically synthesized or produced in a cellular system different from the cell from which it naturally originates will be substantially free from its naturally associated components.

Accordingly, substantially pure polypeptides include those derived from eukaryotic organisms but synthesized in *E. coli* or other prokaryotes.

- 12 -

By "substantially pure DNA" is meant DNA that is free of the genes which, in the naturally-occurring genome of the organism from which the DNA of the invention is derived, flank the gene. The term therefore includes, for example, a recombinant DNA which is incorporated into a vector; into an autonomously replicating plasmid or virus; or into the genomic DNA of a prokaryote or eukaryote; or which exists as a separate molecule (*e.g.*, a cDNA or a genomic or cDNA fragment produced by PCR or restriction endonuclease digestion) independent of other sequences. It also includes a recombinant DNA which is part of a hybrid gene encoding additional polypeptide sequence.

By "transformed cell" is meant a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a DNA molecule encoding (as used herein) an IAP polypeptide.

By "transgene" is meant any piece of DNA which is inserted by artifice into a cell, and becomes part of the genome of the organism which develops from that cell. Such a transgene may include a gene which is partly or entirely heterologous (*i.e.*, foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism.

By "transgenic" is meant any cell which includes a DNA sequence which is inserted by artifice into a cell and becomes part of the genome of the organism which develops from that cell. As used herein, the transgenic organisms are generally transgenic mammalian (*e.g.*, rodents such as rats or mice) and the DNA (transgene) is inserted by artifice into the nuclear genome.

By "transformation" is meant any method for introducing foreign molecules into a cell. Lipofection, calcium phosphate precipitation, retroviral delivery, electroporation, and biolistic transformation are just a few of the teachings which may be used. For example, biolistic transformation is a method for introducing foreign molecules into a cell using velocity driven microprojectiles such as tungsten or gold particles. Such velocity-driven methods originate from pressure bursts which include, but are not limited to, helium-driven, air-driven, and gunpowder-driven techniques. Biolistic transformation may be applied to the transformation or transfection of a wide variety of cell types and intact tissues including, without limitation, intracellular organelles (*e.g.*, and mitochondria and chloroplasts), bacteria, yeast, fungi, algae, animal tissue, and cultured cells.

- 13 -

By "positioned for expression" is meant that the DNA molecule is positioned adjacent to a DNA sequence which directs transcription and translation of the sequence (*i.e.*, facilitates the production of, *e.g.*, an IAP polypeptide, a recombinant protein or a RNA molecule).

By "reporter gene" is meant a gene whose expression may be assayed; such genes
5 include, without limitation, glucuronidase (GUS), luciferase, chloramphenicol transacetylase (CAT), and β -galactosidase.

By "promoter" is meant minimal sequence sufficient to direct transcription. Also included in the invention are those promoter elements which are sufficient to render promoter-dependent gene expression controllable for cell type-specific, tissue-specific or
10 inducible by external signals or agents; such elements may be located in the 5' or 3' regions of the native gene.

By "operably linked" is meant that a gene and one or more regulatory sequences are connected in such a way as to permit gene expression when the appropriate molecules (*e.g.*, transcriptional activator proteins are bound to the regulatory sequences).

15 By "conserved region" is meant any stretch of six or more contiguous amino acids exhibiting at least 30%, preferably 50%, and most preferably 70% amino acid sequence identity between two or more of the IAP family members, (*e.g.*, between human HIAP-1, HIAP-2, and XIAP). Examples of preferred conserved regions are shown (as boxed or designated sequences) in Figures 5-7 and Tables 1 and 2, and include, without limitation,
20 BIR domains and ring zinc finger domains.

By "detectably-labelled" is meant any means for marking and identifying the presence of a molecule, *e.g.*, an oligonucleotide probe or primer, a gene or fragment thereof, or a cDNA molecule. Methods for detectably-labelling a molecule are well known in the art and include, without limitation, radioactive labelling (*e.g.*, with an isotope such as ^{32}P or ^{35}S)
25 and nonradioactive labelling (*e.g.*, chemiluminescent labelling, *e.g.*, fluorescein labelling).

By "antisense," as used herein in reference to nucleic acids, is meant a nucleic acid sequence, regardless of length, that is complementary to a region on the coding strand of nucleic acid molecule (*e.g.*, genomic DNA, cDNA, or mRNA) that encodes an IAP or a NAIP polypeptide. The region of the nucleic acid molecule encoding an IAP or a NAIP
30 polypeptide that the antisense molecule is complementary to may be a region within the coding region, a region upstream of the coding region, a region downstream of the coding

- 14 -

region, or a region within an intron, where the nucleic acid molecule is genomic DNA. Preferably, the antisense nucleic acid is capable of enhancing apoptosis when present in a cell which normally does not undergo sufficient apoptosis and/or is between 8 and 25 nucleotides in length. Preferably, the increase is at least 10%, relative to a control, more preferably 25%,
5 and most preferably 1-fold or more. It will be understood that antisense nucleic acid molecules may have chemical modifications known in the art of antisense design to enhance antisense compound efficiency.

By "purified antibody" is meant antibody which is at least 60%, by weight, free from proteins and naturally occurring organic molecules with which it is naturally associated.
10 Preferably, the preparation is at least 75%, more preferably 90%, and most preferably at least 99%, by weight, antibody, *e.g.*, an IAP specific antibody. A purified antibody may be obtained, for example, by affinity chromatography using recombinantly-produced protein or conserved motif peptides and standard techniques.

By "specifically binds" is meant an antibody that recognizes and binds a protein but
15 that does not substantially recognize and bind other molecules in a sample, *e.g.*, a biological sample, that naturally includes protein.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiments thereof, and from the claims.

Brief Description of the Drawings

20 Fig. 1 is the human XIAP cDNA sequence (SEQ ID NO: 3) and the XIAP polypeptide sequence (SEQ ID NO: 4).

Fig. 2 is the human HIAP-1 cDNA sequence (SEQ ID NO: 5) and the HIAP-1 polypeptide sequence (SEQ ID NO: 6).

Fig. 3 is the human HIAP-2 cDNA sequence (SEQ ID NO: 7) and the HIAP-2
25 polypeptide sequence (SEQ ID NO: 8).

Fig. 4 is the murine XIAP (also referred to as "MIAP-3" or "m-XIAP") cDNA sequence (SEQ ID NO: 9) and encoded murine XIAP polypeptide sequence (SEQ ID NO:
10).

- 15 -

Fig. 5 is the murine HIAP-1 (also referred to as "MIAP-1" or "m-HIAP-1") cDNA sequence (SEQ ID NO: 11) and the encoded murine HIAP-1 polypeptide sequence (SEQ ID NO: 12).

Fig. 6 is the murine HIAP-2 (also referred to as "MIAP-2" or "m-HIAP-2") cDNA
5 sequence (SEQ ID NO: 13) and the encoded murine HIAP-2 polypeptide (SEQ ID NO: 14).

Fig. 7 is a photograph of a Northern blot illustrating human HIAP-1 and HIAP-2 mRNA expression in human tissues.

Fig. 8 is a photograph of a Northern blot illustrating human HIAP-2 mRNA expression in human tissues.

10 Fig. 9 is a photograph of a Northern blot illustrating human XIAP mRNA expression in human tissues.

Figs. 10A - 10D are graphs depicting suppression of apoptosis by XIAP, HIAP-1, HIAP-2, BCL-2, SMN, and 6-MYC.

Fig. 11 is a photograph of an agarose gel containing cDNA fragments that were
15 amplified, with HIAP 1-specific primers, from RNA obtained from Raji, Ramos, EB-3, Burkitt's lymphoma cells, and Jiyoye cells, and cells from normal placenta.

Fig. 12 is a photograph of a Western blot containing protein extracted from Jurkat and astrocytoma cells stained with an anti-XIAP antibody. The position and size of a series of marker proteins is indicated.

20 Fig. 13 is a photograph of a Western blot containing protein extracted from Jurkat cells following treatment as described in Example XII. The blot was stained with a rabbit polyclonal anti-XIAP antibody. Lane 1, negative control; lane 2, anti-Fas antibody; lane 3, anti-Fas antibody and cycloheximide; lane 4, TNF- α ; lane 5, TNF- α and cycloheximide.

Fig. 14 is a photograph of a Western blot containing protein extracted from HeLa
25 cells following exposure to anti-Fas antibodies. The blot was stained with a rabbit polyclonal anti-XIAP antibody. Lane 1, negative control; lane 2, cycloheximide; lane 3, anti-Fas antibody; lane 4, anti-Fas antibody and cycloheximide; lane 5, TNF- α ; lane 6, TNF- α and cycloheximide.

Figs. 15A and 15B are photographs of Western blots stained with rabbit polyclonal
30 anti-XIAP antibody. Protein was extracted from HeLa cells (Fig. 15A) and Jurkat cells (Fig. 15B) immediately, 1, 2, 3, 5, 10, and 22 hours after exposure to anti-Fas antibody.

- 16 -

Figs. 16A and 16B are photographs of Western blots stained with an anti-CPP32 antibody (Fig. 16A) or a rabbit polyclonal anti-XIAP antibody (Fig. 16B). Protein was extracted from Jurkat cells immediately, 3 hours, or 7 hours after exposure to an anti-Fas antibody. In addition to total protein, cytoplasmic and nuclear extracts are shown.

5 Fig. 17 is a photograph of a polyacrylamide gel following electrophoresis of the products of an *in vitro* XIAP cleavage assay.

Figs. 18 and 19 shows the increased level of HIAP-1 and HIAP-2 mRNA, respectively, in breast cancer cell lines having p53 mutations (lanes 5-7). The bottom portion of the figure shows the control.

10 Fig. 20 shows the influence of Taxol on DNA fragmentation in Cisplatin-sensitive (right) and resistant (left) human ovarian epithelial cancer cells.

Fig. 21 shows the influence of Cisplatin on DNA fragmentation in sensitive (right) and resistant (left) human ovarian epithelial cancer cells.

Fig. 22 shows the effects of Taxol on XIAP and HIAP-2 protein levels in Cisplatin
15 sensitive (right) and resistant (left) human ovarian epithelial cancer cells.

Figs. 23A and 23B show the influence of Taxol and TGF β on HIAP-2 mRNA levels in Cisplatin sensitive (right) and resistant (left) human epithelial cancer cells.

Figs. 24A and 24B show the effect of TGF β on XIAP protein expression (Fig. 24A) and DNA fragmentation (Fig. 24B) in Cisplatin-sensitive (OV2008) and cisplatin-resistant
20 (C13) cells.

Fig. 25 is a series of bar graphs showing the effect of XIAP and HIAP-2 down-regulation on ovarian epithelial cancer cell viability and number. The top two panels show dead cells as a percentage of total cell population. The bottom two panels illustrate total cell number at the end of the infection period. Data represents the mean \pm SEM of four
25 experiments. **p<0.01, ***p<0.001 (compared to vector control).

Fig. 26A is a set of photographs showing the influence of XIAP down-regulation on whole cell morphology (phase contrast; black arrows indicate cell detachment) in OV2008 cells after 60 hours of adenovirus infection with vector only (left) or adenoviral antisense XIAP (right). MOI=5 (1X; "a" and "b"); magnification 400X.

30 Fig. 26B is a series of photographs ("a" through "d") showing the influence of XIAP down-regulation on nuclear morphology (Hoechst staining; white arrows show nuclear

- 17 -

fragmentation) in OV2008 cells after 60 hours of adenovirus infection with vector only ("a" and "c") or adenoviral antisense XIAP ("b" and "d"). MOI=5 (1X; "a" and "b") and MOI=10 (2X; "c" and "d"); magnification 400X.

Fig. 26C is a bar graph showing the influence of XIAP down-regulation on the extent of apoptosis in OV2008 cells after 60 hours of no treatment, adenovirus infection with vector only, or adenovirus infection with antisense XIAP. Data represents the mean \pm SEM of three to four experiments. MOI=5 (1X) and MOI=10 (2X); * p <0.05, ** p <0.01 (compared to vector control).

Fig. 26D is a representative Western blotting analysis showing effective XIAP antisense infection in OV2008 cells after 60 hours of no treatment, adenovirus infection with vector only, or adenovirus infection with antisense XIAP. Lanes are, from left to right: control, vector (1X), vector (2X), antisense XIAP (1X), and antisense XIAP (2X). MOI=5 (1X) and MOI=10 (2X).

Fig. 26E is a bar graph showing changes in XIAP protein content in OV2008 cells after 60 hours of no treatment, adenovirus infection with vector only, or adenovirus infection with antisense XIAP, as analyzed densitometrically, using a Molecular Dynamic Phosphoimager. Data represents the mean \pm SEM of three to four experiments. MOI=5 (1X) and MOI=10 (2X); * p <0.05, ** p <0.01 (compared to vector control).

Fig. 27A is a series of photographs showing effects of cisplatin-induced apoptosis (at 0 and 30 μ M cisplatin in a 24 hour culture) the nuclear morphology of cisplatin-sensitive cells (OV2008; left two photographs) and cisplatin-resistant cells (C13; right two photographs), using Hoechst staining, magnification 400X; arrows show fragmented nuclei.

Fig. 27B is a set of photographs showing agarose gel immobilized electrophoretically resolved apoptotic low molecular weight DNA fragmentation from cisplatin treated OV2008 and C13 cells.

Fig. 27C is a line graph showing a concentration-response study of apoptosis in OV2008 and C13 cells following 24 hours of culture in 0, 10, 20, and 30 μ M cisplatin. Data represents the mean \pm SEM of three experiments. ** p <0.01 (compared to control).

Fig. 28A is a series of representative Western blotting analyses showing concentration-dependent inhibition of XIAP and HIAP-2 protein expression in cisplatin-sensitive (OV2008) and cisplatin-resistant (C13) ovarian epithelial cancer cells following 24

- 18 -

hour culture with 0, 10, 20, and 30 μ M cisplatin. Equal amounts of solubilized proteins (20-60 μ g/lane, depending on the individual experiment) were analyzed by Western blot using anti-human XIAP or anti-HIAP-2 antibodies.

Fig. 28B is a panel of bar graphs showing the changes in XIAP (left two graphs) and HIAP-2 (right two graphs) protein content as analyzed densitometrically, using the Molecular Dynamics Phosphoimager, for cisplatin-treated (24 hours at indicated concentration) OV2008 cells (upper two graphs) and C13 cells (lower two graphs). Data represents the mean \pm SEM of three experiments. * p <0.05, ** p <0.01 (compared to control).

Fig. 29A is a series of representative Western blotting analyses showing concentration-dependent inhibition of XIAP and HIAP-2 protein expression in cisplatin-sensitive (OV2008) and cisplatin-resistant (C13) ovarian epithelial cancer cells following 6, 12, or 24 hours of culture with or without 30 μ M cisplatin. Equal amounts of solubilized proteins (20-60 μ g/lane, depending on the individual experiment) were analyzed by Western blot using anti-human XIAP or anti-HIAP-2 antibodies.

Fig. 29B is a panel of bar graphs showing the changes in XIAP (left two graphs) and HIAP-2 (right two graphs) protein content as analyzed densitometrically, using the Molecular Dynamics Phosphoimager, for OV2008 cells (white bars) and C13 cells (black bars) cultured with or without 30 μ M cisplatin for 6, 12, or 24 hours. Data represents the mean \pm SEM of three experiments. * p <0.05, ** p <0.01 (compared to control).

Fig. 30A is a series of representative Western blotting analyses showing concentration-dependent inhibition of XIAP and HIAP-2 protein expression in cisplatin-sensitive (A2780s) and cisplatin-resistant (A2780cp) ovarian epithelial cancer cells following hours of culture with or without 30 μ M cisplatin. Equal amounts of solubilized proteins (40-60 μ g/lane, depending on the individual experiment) were analyzed by Western blot using anti-human XIAP or anti-HIAP-2 antibodies.

Fig. 30B is a panel of bar graphs showing the changes in XIAP (top graph) and HIAP-2 (bottom graph) protein content as analyzed densitometrically, using the Molecular Dynamics Phosphoimager, for A2780s cells (left) and A2780cp cells (right) cultured with (black bars) or without (white bars) 30 μ M cisplatin for 24 hours. Data represents the mean \pm SEM of three experiments. ** p <0.01 (compared to control).

- 19 -

Fig. 31A is set of photographs ("a" through "d") showing the effects of XIAP overexpression on the apoptotic action of cisplatin (30 μ M) on nuclear morphology of cisplatin-sensitive OV2008 cells after 48 hours of infection of these cells with adenoviral sense XIAP cDNA or vector only (control). At a magnification of 400X. "a", vector (no cisplatin); "b", sense XIAP (no cisplatin); "c", vector plus cisplatin-treatment; "d", sense XIAP plus cisplatin treatment.

Fig. 31B is a graph showing the percentage of total cell population undergoing apoptosis of 30 μ M cisplatin-treated OV2008 cells following 48 hours of infection of these cells with adenoviral sense XIAP cDNA or vector only (control). Data represent mean \pm SEM of three experiments. * p <0.05, *** p <0.001 (compared to vector control); \pm p <0.01, $\pm\pm$ p <0.001 (compared to vector plus cisplatin group).

Fig. 31C is a representative Western blotting analysis showing changes in XIAP protein content in OV2008 cells following infection with adenoviral sense XIAP cDNA or vector only (control) with or without treatment with 30 μ M cisplatin. Lanes are, from left to right: control, vector, vector plus cisplatin, sense XIAP, and sense XIAP plus cisplatin.

Fig. 31D is a graph showing the changes in XIAP protein content in OV2008 cells following infection with adenoviral sense XIAP cDNA or vector only (control) with or without treatment with 30 μ M cisplatin, as analyzed densitometrically, using the Molecular Dynamic Phosphoimager. Data represent mean \pm SEM of three experiments. * p <0.05, *** p <0.001 (compared to vector control); \pm p <0.01, $\pm\pm$ p <0.001 (compared to vector + cisplatin group).

Figs. 32A-32D are a series of photographs showing the *in situ* detection of apoptosis (using TUNEL) and immunolocalization of PCNA, XIAP and HIAP-2 in human ovarian surface epithelial tumour tissue. Fig. 32A indicates the *in situ* TUNEL localization of apoptotic cells. Figs. 32B, 32C, and 32D represent immuno-reactivates for PCNA, XIAP and HIAP-2, respectively. The regions of tumor shown in the circle and the rectangle in each of Figs. 32A-32D was TUNEL-positive and TUNEL-negative, respectively. Magnification is 400X.

Detailed Description

- 20 -

Previously, we have provided a novel family of inhibitors of apoptosis, the IAPs, and an additional related anti-apoptotic protein, NAIP. Here we provide identification of cancer types in which dysregulation of the IAPs and NAIP is apparent. Our results are of paramount importance and provide diagnostics, prognostics, treatments, and drug screens aimed at the
5 detection and effective treatment of cancer.

Cancer Screening

We initially studied IAP and NAIP expression levels in a variety of normal tissues and cancer cell lines using commercially available northern blots. Elevated XIAP, HIAP-1 and HIAP-2 mRNA was noted in a surprising number of cancer lines of diverse lineage,
10 including colorectal cancer, lymphoma, leukemia, and melanoma cell lines. In contrast, BCL-2 mRNA was elevated in only a single cell line. Although this result reinforced the importance of the IAPs and NAIP in cancer, the question remained as to whether the individual cancer cell lines on the blot were representative of the cancer type. As a result, we screened panels of cancer cell lines of particular tumor type by northern blot and quantitative
15 RT-PCR analysis in order to ascertain the frequency of IAP and NAIP dysregulation. The results are summarized as follows:

Burkitt's Lymphoma.

We studied both the frequency and consequences of IAP upregulation in Burkitt's lymphoma. Elevated levels of HIAP-1 and HIAP-2 have been found in the vast majority of
20 the Burkitt's cell lines examined. Furthermore, those Burkitt's lines expressing low levels of HIAP-1 are transcriptionally activated by Epstein-Barr virus (EBV) infection.

Breast Adenocarcinoma.

A key observation was made in this survey, in which a correlation was observed between drug resistance, p53 status, and HIAP-1 and HIAP-2 expression. Four of the cell
25 lines possessed wild-type p53, while three possessed documented p53 mutations that correlated with resistance to the anti-cancer drug adriamycin. Significantly, the three lines which were relatively more drug resistant also displayed elevated HIAP-1 and HIAP-2

- 21 -

mRNA levels. These results indicate that one of the ways that p53 controls apoptosis is through regulation of these genes.

Ovarian Carcinoma.

mRNA *in situ* analysis suggest a role for NAIP in the developmental biology of the
5 ovary. Overexpression of HIAP-2 and XIAP mRNA has also been documented in some ovarian cancer cell lines.

Pancreatic Cancer.

Approximately 25% of the pancreatic cancer cell lines tested to date demonstrate HIAP-1 and HIAP-2 mRNA elevation.

10 *Summary of Cancer Panels.*

To date, a significant fraction of cancer cell lines of each type examined display elevated IAP levels. Increased NAIP levels are also implicated in cancer. Our results indicate that HIAP-1 and HIAP-2 tend to be the most frequently and dramatically upregulated. The apparent coordinate regulation of both genes was surprising given that the
15 normal tissue distribution of these proteins is very different. Our observations are strengthened by the fact that HIAP-1 and HIAP-2 reside in tandem array on chromosome 11q23, a site frequently rearranged in lymphomas and leukemias.

Transcriptional regulation of the IAPs in cancer cell lines.

Our experiments have established a correlation between p53 status and transcriptional
20 overexpression of HIAP-1 and HIAP-2. This provides an important new way in which to enhance apoptosis, particularly in view of the fact that the mechanism by which p53 controls cell fate remains largely unknown. It has previously been documented that wild-type p53 negatively down-regulates BCL-2, and positively upregulates the BCL-2 antagonist BAX. In some cancer cell types, mutation of p53 causes a two-fold effect; namely, the upregulation of
25 BCL-2, and down regulation of BAX, both of which contribute to the anti-apoptotic phenotype. While not wishing to bind ourselves to a particular theory, we believe that wild-type p53 also transcriptionally suppresses HIAP-1 and HIAP-2. DNA damage that includes

- 22 -

the increase in wild-type levels p53 levels would therefore result in decreased HIAP-1 and HIAP-2 in normal cells, resulting in apoptosis. Mutations in the p53 gene would therefore result in a loss of transcriptional control of these IAP genes. As a result, p53 mutant cancer cells would display constitutively high levels of HIAP-1 and HIAP-2, rendering the cells resistant to anti-cancer therapies. The p53/HIAP-1 and HIAP-2 correlations may be extended to the other cancer cell line panels. One may directly demonstrate p53 regulation of the IAPs using transfection assays and northern blot analysis.

Accordingly, we predict that cancer cells having p53 mutations (p53*) will have increased IAP levels resulting in a poor response to chemotherapeutics. Because IAP levels may be assessed more readily than the presence of a p53* mutation, our discovery also provides an important improvement in cancer diagnosis and prognosis (see below).

Transgenic Mice

We have constructed a number of IAP and NAIP transgenic mouse expression vectors, including T-cell, B-cell, and neuronal specific promoter constructs. Founder mice have been identified and are viable, and, for most of these constructs, we have developed breeding colonies. These mice will likely be prone to cancers of the tissue types in which the promoter is active. Thus the mice provide an excellent resource for testing the efficacy of anti-sense oligonucleotides and for screening for apoptosis-enhancing cancer therapeutics. Standard mouse drug screening models and gene delivery protocols may be employed to utilize the mice for this purpose.

Diagnostic/Prognostic Reagents

There is a relative lack of diagnostic and prognostic tests which clinical oncologists may utilize in determining the appropriate degree of intervention in the treatment of cancer. Mutation of the p53 gene remains one of the best prognostic indicators in cancer biology. However, the number of different mutations identified to date is great and the mutations are scattered throughout the gene. In addition, many mutations in p53 result in an inappropriate stabilization of the protein, which allows detection at the protein level rather than at the mRNA level. Mutations which alter the transactivation/repression activities of the protein are not necessarily apparent at either the mRNA or protein levels. On the other hand, if IAP

- 23 -

and NAIP expression levels correlate with p53 mutation they may provide more valuable prognostic information and assist in the determination of which patients require more aggressive treatment or which patients are, perhaps, not treatable with currently approved therapies. This latter class of patients may be identified as ideal candidates for clinical testing of new cancer therapeutics, particularly those which decrease IAP levels or act in a manner independent of the anti-apoptotic pathway.

Thus, the invention provides at least two assays for prognosis and diagnosis. Semi-quantitative RT-PCR based assays may be used to assay for IAP and/or NAIP gene or protein expression levels. Alternatively, monoclonal antibodies may be incorporated into an ELISA (enzyme-linked immunosorbent assay) -type assay for direct determination of protein levels.

Therapeutic Products

For IAP or NAIP-related therapies, one may employ the paradigms utilized for BCL-2 and RAS antisense development, although, in contrast to RAS antisense, accommodation of mutations is not required. Most useful are antisense constructs which enhance apoptosis at least 10%, preferably by enhancing degradation of the RNA in the nucleus.

In addition to the antisense approaches described herein, the invention features small molecule screening assays which may be used to identify lead compounds that negatively regulate the IAPs or NAIP. For example, compounds which enhance apoptosis in the presence of IAP overexpression or which decrease the level of IAP biological activity may be detected and are useful cancer therapeutics.

Molecules that are found, by the methods described herein, to effectively modulate IAP gene expression or polypeptide activity may be tested further in standard animal cancer models. If they continue to function successfully in an *in vivo* setting, they may be used as therapeutics to either inhibit or enhance apoptosis, as appropriate.

25 *Manipulation of cancer chemotherapeutic drug resistance using an antisense oligonucleotide and fragment approaches.*

We have documented that overexpression of the IAPs renders cell lines resistant to serum growth factor withdrawal, tumor necrosis factor alpha (TNF) and menadione exposure, all of which are treatments that normally induce apoptosis. Herein we describe the extension

- 24 -

of these studies to cancer cell lines using apoptotic triggers used in clinical situations, such as doxorubicin, adriamycin, and methotrexate. Our findings have led up to design antisense RNA therapeutics. Rapid screening of multiple cell lines for apoptotic response has been made feasible through the generation of a series of sense and antisense adenoviral IAP and
5 NAIP expression vectors, as well as control lacZ viruses. One may now show enhanced drug resistance or enhanced drug sensitivity using these expression constructs. In addition, antisense adenovirus constructs have been developed and used to test reversal of the drug resistant phenotype of appropriate cell lines.

We have surveyed cancer cell lines with the objective of identifying tumor types in
10 which IAP or NAIP overexpression is apparent or altered and these results are described both above and in the Examples below. Concomitant to this research, we have designed a series of antisense oligonucleotides to various regions of each of the IAPs. After testing in an assay system, *i.e.*, with the adenoviral vectors system, these oligonucleotides, as well as antisense oligonucleotides to various regions of NAIP, may be used to enhance drug
15 sensitivity. Animal modeling of the effectiveness of antisense IAP and NAIP oligonucleotides may also be employed as a step in testing and appropriate transgenic mammals for this are described above and also generally available in the art.

The following describes some of the testing systems which may be employed.

Anti-Cancer Gene Therapy

20 Retroviral vectors, adenoviral vectors, adeno-associated viral vectors, or other viral vectors with the appropriate tropism for cells requiring enhanced apoptosis (for example, breast cancer and ovarian cancer cells) may be used as an oligonucleotide transfer delivery system for a therapeutic constructs.

Alternatively, standard non-viral delivery methods may be used. Numerous vectors
25 useful for viral delivery are generally known (Miller, A.D., Human Gene Therapy 1: 5-14, 1990; Friedman, T., Science 244: 1275-1281, 1989; Eglitis and Anderson, BioTechniques 6: 608-614, 1988; Tolstoshev and Anderson, Curr. Opin. Biotech. 1: 55-61, 1990; Cornetta *et al.*, Prog. Nucl. Acid Res. and Mol. Biol. 36: 311-322, 1987; Anderson, W. F., Science 226: 401-409, 1984; Moen, R. C., Blood Cells 17: 407-416, 1991; Miller *et al.*, BioTechniques 7:

- 25 -

980-990, 1989; Le Gal La Salle *et al.*, Science 259: 988-990, 1993; and Johnson, Chest 107: 77S-83S, 1995).

Retroviral vectors are particularly well developed and have been used in clinical settings (Rosenberg *et al.*, New Engl. J. Med. 323: 570-578, 1990; Anderson *et al.*, U.S.

5 Patent No. 5,399,346).

Non-viral approaches may also be employed for the introduction of therapeutic nucleic acid molecules (*e.g.*, oligonucleotides) into cells otherwise predicted to undergo apoptosis. For example, IAP may be introduced into a neuron or a T cell by lipofection (Felgner *et al.*, Proc. Natl. Acad. Sci. USA 84: 7413-7417, 1987; Ono *et al.*, Neurosci. Lett. 10 117: 259-263, 1990; Brigham *et al.*, Am. J. Med. Sci. 298: 278-281, 1989; Staubinger *et al.*, Meth. Enz. 101: 512-527, 1983), asialorsonuroid-polylysine conjugation (Wu *et al.*, J. Biol. Chem. 263: 14621-14624, 1988; Wu *et al.*, J. Biol. Chem. 264: 16985-16987, 1989); direct deliver in saline; or, less preferably, microinjection under surgical conditions (Wolff *et al.*, Science 247: 1465-1468, 1990).

15 For any of the methods of application described above, the therapeutic nucleic acid construct is preferably applied to the site of the needed apoptosis event (for example, by injection). However, it may also be applied to tissue in the vicinity of the predicted apoptosis event, to a blood vessel supplying the cells predicted to require enhanced apoptosis, or orally.

In the constructs described, nucleic acid expression can be directed from any suitable 20 promoter (*e.g.*, the human cytomegalovirus (CMV), simian virus 40 (SV40), or metallothionein promoters), and regulated by any appropriate mammalian regulatory element. For example, if desired, enhancers known to preferentially direct gene expression in ovarian cells, breast tissue, neural cells, T cells, or B cells may be used to direct expression. The enhancers used could include, without limitation, those that are characterized as tissue- 25 or cell-specific in their expression. Alternatively, if a clone used as a therapeutic construct, regulation may be mediated by the cognate regulatory sequences or, if desired, by regulatory sequences derived from a heterologous source, including any of the promoters or regulatory elements described above.

Anti-cancer therapy is also accomplished by direct administration of the therapeutic 30 sense IAP nucleic acid or antisense IAP nucleic acid (*e.g.*, oligonucleotides) to a cell that is expected to require enhanced apoptosis. The nucleic acid molecule may be produced and

- 26 -

isolated by any standard technique, but is most readily produced by *in vitro* transcription using an IAP related nucleic acid under the control of a high efficiency promoter (*e.g.*, the T7 promoter), or, by organic synthesis techniques (for, *e.g.*, oligonucleotides).

Administration of IAP antisense nucleic acid to malignant cells can be carried out by any of
5 the methods for direct nucleic acid administration described above, or any method otherwise known in the art.

Another therapeutic approach within the invention involves administration of recombinant IAP protein fragments or IAP antibodies, either directly to the site where enhanced apoptosis is desirable (for example, by injection) or systemically (for example, by
10 any conventional recombinant protein administration technique).

The dosage of a NAIP or an IAP protein, a polypeptide fragment thereof, a mutant thereof, or antibodies that specifically bind NAIP or an IAP polypeptide depends on a number of factors, including the size and health of the individual patient, but, generally, between 0.1 mg and 500 mg inclusive are administered per day to an adult in any
15 pharmaceutically acceptable formulation.

Administration of IAP and NAIP Polypeptides, Nucleic Acids, and Inhibitors of IAP or NAIP Synthesis or Function

An IAP or NAIP mutant protein or protein fragment, a nucleic acid molecule encoding the same, a nucleic acid molecule encoding an IAP or NAIP antisense nucleic acid,
20 or a inhibitor of an IAPs or NAIP may be administered within a pharmaceutically-acceptable diluent, carrier, or excipient, in unit dosage form. Conventional pharmaceutical practice may be employed to provide suitable formulations or compositions to administer the compounds to patients suffering from a disease that is caused by excessive cell proliferation.

Administration may begin before the patient is symptomatic.

25 Any appropriate route of administration may be employed, for example, administration may be parenteral, intravenous, intraarterial, subcutaneous, intramuscular, intracranial, intraorbital, ophthalmic, intraventricular, intrathecal, intracapsular, intracisternal, intraperitoneal, intranasal, aerosol, suppository, or oral administration. For example, therapeutic formulations may be in the form of liquid solutions or suspensions; for oral

- 27 -

administration, formulations may be in the form of tablets or capsules; and for intranasal formulations, in the form of powders, nasal drops, or aerosols.

Methods well known in the art for making formulations are found, for example, in Remington's Pharmaceutical Sciences, (18th edition), ed. A. Gennaro, 1990, Mack Publishing Company, Easton, PA. Formulations for parenteral administration may, for example, contain excipients, sterile water, or saline, polyalkylene glycols such as polyethylene glycol, oils of vegetable origin, or hydrogenated naphthalenes. Biocompatible, biodegradable lactide polymer, lactide/glycolide copolymer, or polyoxyethylene-polyoxypropylene copolymers may be used to control the release of the compounds. Other potentially useful parenteral delivery systems for IAP or NAIP modulatory compounds include ethylene-vinyl acetate copolymer particles, osmotic pumps, implantable infusion systems, and liposomes. Formulations for inhalation may contain excipients, for example, lactose, or may be aqueous solutions containing, for example, polyoxyethylene-9-lauryl ether, glycocholate and deoxycholate, or may be oily solutions for administration in the form of nasal drops, or as a gel.

If desired, treatment with an IAP or NAIP mutant proteins or IAP or NAIP fragments, related genes, or other modulatory compounds may be combined with more traditional therapies for the proliferative disease such as surgery or chemotherapy.

Detection of Conditions Involving Insufficient Apoptosis

IAP and NAIP polypeptides and nucleic acid sequences find diagnostic use in the detection or monitoring of conditions involving insufficient levels of apoptosis, *i.e.*, proliferative disease. For example, increased expression of IAPs or NAIP, alterations in localization, and IAP or NAIP cleavage correlate with inhibition of apoptosis and cancer in humans. Accordingly, an increase in the level of IAP or NAIP production may provide an indication of a proliferative condition or a predisposition to such a condition. Levels of IAP or NAIP expression may be assayed by any standard technique. For example, IAP or NAIP expression in a biological sample (*e.g.*, a biopsy sample) may be monitored by standard Northern blot analysis or may be aided by PCR (see, *e.g.*, Ausubel *et al.*, Current Protocols in Molecular Biology, John Wiley & Sons, New York, 1994; PCR Technology: Principles and

- 28 -

Applications for DNA Amplification, H.A. Ehrlich, Ed., Stockton Press, NY; Yap *et al.*, Nucl. Acids. Res. 19: 4294, 1991).

Alternatively, a biological sample obtained from a patient may be analyzed for one or more mutations in the IAP or NAIP sequences or p53 sequences using a mismatch detection
5 approach. Generally, these techniques involve PCR amplification of nucleic acid from the patient sample, followed by identification of the mutation (*i.e.*, mismatch) by either altered hybridization, aberrant electrophoretic gel migration, binding or cleavage mediated by mismatch binding proteins, or direct nucleic acid sequencing. Any of these techniques may be used to facilitate mutant IAP or NAIP detection, and each is well known in the art;
10 examples of particular techniques are described, without limitation, in Orita *et al.*, Proc. Natl. Acad. Sci. USA 86: 2766-2770, 1989; Sheffield *et al.*, Proc. Natl. Acad. Sci. USA 86: 232-236, 1989).

In yet another approach, immunoassays are used to detect or monitor IAP or NAIP protein in a biological sample. IAP or NAIP-specific polyclonal or monoclonal antibodies
15 (produced as described above) may be used in any standard immunoassay format (*e.g.*, ELISA, Western blot, or RIA) to measure IAP or NAIP polypeptide levels from cancerous control cells. These levels would be compared to wild-type IAP or NAIP levels, with a decrease in IAP production relative to a wild-type cell indicating a condition involving increased apoptosis and a decrease relative to a known cancer cell indicating a decreased
20 likelihood of an IAP or NAIP-related cancer. Examples of immunoassays are described, *e.g.*, in Ausubel *et al.*, *supra*. Immunohistochemical techniques may also be utilized for IAP or NAIP detection. For example, a tissue sample may be obtained from a patient, sectioned, and stained for the presence of IAP or NAIP using an anti-IAP or anti-NAIP antibodies and any standard detection system (*e.g.*, one which includes a secondary antibody conjugated to
25 horseradish peroxidase). General guidance regarding such techniques can be found in, *e.g.*, Bancroft and Stevens (Theory and Practice of Histological Techniques, Churchill Livingstone, 1982) and Ausubel *et al.* (*supra*).

In one preferred example, a combined diagnostic method may be employed that begins with an evaluation of IAP or NAIP protein production (for example, by
30 immunological techniques or the protein truncation test (Hogerrorst *et al.*, Nature Genetics 10:208-212, 1995)) and also includes a nucleic acid-based detection technique designed to

- 29 -

identify more subtle IAP or NAIP alterations, *e.g.*, mutations. As described above, a number of mismatch detection assays are available to those skilled in the art, and any preferred technique may be used. Mutations in IAP or NAIP may be detected that either result in enhanced IAP or NAIP expression or alterations in IAP or NAIP biological activity. In a variation of this combined diagnostic method, IAP or NAIP biological activity is measured as anti-apoptotic activity using any appropriate apoptosis assay system (for example, those described above).

Mismatch detection assays also provide an opportunity to diagnose an IAP-mediated or an NAIP-mediated predisposition to diseases caused by insufficient apoptosis. For example, a patient heterozygous for an IAP or a NAIP mutation may show no clinical symptoms and yet possess a higher than normal probability of developing one or more types of proliferative diseases. Given this diagnosis, a patient may take precautions to minimize their exposure to adverse environmental factors (for example, UV exposure or chemical mutagens) and to carefully monitor their medical condition (for example, through frequent physical examinations). This type of IAP or NAIP diagnostic approach may also be used to detect IAP or NAIP mutations in prenatal screens. The IAP or NAIP diagnostic assays described above may be carried out using any biological sample (for example, any biopsy sample or bodily fluid or tissue) in which IAP or NAIP is normally expressed. Identification of a mutant IAP or NAIP gene may also be assayed using these sources for test samples.

Alternatively, an alteration in IAP or NAIP activity, particularly as part of a diagnosis for predisposition to IAP-associated or NAIP-associated proliferative disease, may be tested using a nucleic acid sample from any cell, for example, by mismatch detection techniques. Preferably, the DNA sample is subjected to PCR amplification prior to analysis.

The following examples are meant to illustrate, not limit, the invention.

EXAMPLE 1: ELEVATED IAP LEVELS IN CANCER CELL LINES

In order to specifically demonstrate the utility of IAP gene sequences as diagnostics and prognostics for cancer, a Human Cancer Cell Line Multiple Tissue Northern Blot (Clontech, Palo Alto, CA; #7757-1) was probed. This Northern blot contained approximately 2 µg of poly A⁺ RNA per lane from eight different human cell lines: (1) promyelocytic leukemia HL-60, (2) HeLa cell S3, (3) chronic myelogenous leukemia K-562, (4)

- 30 -

lymphoblastic leukemia MOLT-4, (5) Burkitt's lymphoma Raji, (6) colorectal adenocarcinoma SW480, (7) lung carcinoma A549, and (8) melanoma G361. As a control, a Human Multiple Tissue Northern Blot (Clontech, Palo Alto, CA; #7759-1) was probed. This Northern blot contained approximately 2 µg of poly A⁺ RNA from eight different human tissues: (1) spleen, (2) thymus, (3) prostate, (4) testis, (5) ovary, (6) small intestine, (7) colon, and (8) peripheral blood leukocytes.

The Northern blots were hybridized sequentially with: (1) a 1.6 kb probe to the XIAP coding region, (2) a 375 bp HIAP-2 specific probe corresponding to the 3' untranslated region, (3) a 1.3 kb probe to the coding region of HIAP-1, which cross-reacts with HIAP-2, (4) a 1.0 kb probe derived from the coding region of BCL-2, and (5) a probe to β-actin, which was provided by the manufacturer. Hybridization was carried out at 50°C overnight, according to the manufacturer's suggestion. The blot was washed twice with 2X SSC, 0.1% SDS at room temperature for 15 minutes and then with 2X SSC, 0.1% SDS at 50°C.

All cancer lines tested showed increased IAP expression relative to samples from non-cancerous control tissues (Table 1). Expression of XIAP was particularly high in HeLa (S-3), chronic myelogenous leukemia (K-562), colorectal adenocarcinoma (SW-480), and melanoma (G-361) lines. Expression of HIAP-1 was extremely high in Burkitt's lymphoma, and was also elevated in colorectal adenocarcinoma. Expression of HIAP-2 was particularly high in chronic myelogenous leukemia (K-562) and colorectal adenocarcinoma (SW-480). Expression of BCL-2 was upregulated only in HL-60 leukemia cells.

TABLE 1

NORTHERN BLOT IAP RNA LEVELS IN CANCER CELLS*

| | XIAP | HIAP-1 | HIAP- 2 |
|------------------------------------|------|--------|---------|
| Promyelocytic Leukemia HL-60 | + | + | + |
| Hela S-3 | + | + | + |
| Chronic Myelogenous Leukemia K-562 | +++ | + | +++ |
| 5 Lymphoblastic Leukemia MOLT-4 | +++ | + | + |
| Burkitt's Lymphoma Raji | + | +(x10) | + |
| Colorectal Adenocarcinoma SW-480 | +++ | +++ | +++ |
| Lung Carcinoma A-549 | + | + | + |
| 10 Melanoma G-361 | +++ | + | + |

- 10 *Levels are indicated by a (+) and are the approximate increase in RNA levels relative to Northern blots of RNA from non-cancerous control cell lines. A single plus indicates an estimated increase of at least 1-fold

These observations indicate that upregulation of the anti-apoptotic IAP genes may be a widespread phenomenon in proliferative diseases, perhaps occurring much more frequently than upregulation of BCL-2. Furthermore, upregulation may be necessary for the establishment or maintenance of the transformed state of cancerous cells.

In order to pursue the observation described above, *i.e.*, that HIAP-1 is overexpressed in the Raji Burkitt's lymphoma cell line, RT-PCR analysis was performed in multiple Burkitt's lymphoma cell lines. Total RNA was extracted from cells of the Raji, Ramos, EB-20 3, and Jiyoye cell lines, and as a positive control, from normal placental tissue. The RNA was reverse transcribed, and amplified by PCR with the following set of oligonucleotide primers:

5'-AGTGC GGG TTTT TATTATGTG-3' (SEQ ID NO: 15) and
 5'-AGATGACCACAAGGAATAAACACTA-3' (SEQ ID NO: 16), which selectively
 25 amplify a hiap-1 cDNA fragment. RT-PCR was conducted using a Perkin Elmer 480 Thermocycler to carry out 35 cycles of the following program: 94°C for 1 minute, 50°C for 1.5 minutes, and 72°C for 1 minute. The PCR reaction product was electrophoresed on an agarose gel and stained with ethidium bromide. Amplified cDNA fragments of the appropriate size were clearly visible in all lanes containing Burkitt's lymphoma samples, but

- 32 -

absent in the lanes containing the normal placental tissue sample, and absent in lanes containing negative control samples, where template DNA was omitted from the reaction (Fig. 11).

EXAMPLE 2: IAPs IN BREAST CANCER

5 The following data relate to the regulation and role of IAPs in cancer cells. Figs. 18 and 19 show data demonstrating that HIAP-1 and HIAP-2 are both upregulated in breast cancer cell lines that contain mutant p53. The lanes contain 20 µg of total RNA from the following lines: 1. MCF-7 (clone 1, wt p53); 2. MCF-7 (clone 2, wt p53); 3. MCF-7 (American Type Culture Collection, wt p53); 4. MCF-7 (parental line, California, wt p53); 5. 10 MCF-7 (California, adriamycin resistant variant, mutant p53); 6. MDA MB 231 (ATCC, mutant p53, codon 280); 7. T47-D (ATCC, mutant p53, codon 194); 8. ZR-75 (ATCC, wt p53). The amount of RNA loaded on each gel was controlled for by hybridization with glyceral phosphate dehydrogenase (GAPDH).

EXAMPLE 3: IAPS IN OVARIAN CANCER

15 *Overview.*

Epithelial ovarian cancer is the leading cause of death from gynecologic malignancy. Although clinical and histologic prognostic factors such as tumor grade and surgical stage are well understood, the biologic process that leads to uncontrolled cellular growth is less clear. The control of cell numbers during tissue growth is thought to be the results of a balance of 20 cell proliferation and cell death. An aberration in this natural homeostasis likely contributes to malignant cellular transformation.

Recent studies on ovarian cancer cell biology have suggested that the deregulation of apoptosis may be one of the underlying pathologic mechanism in this disease. However, the molecular mechanisms involved in its regulation is poorly understood and the role and 25 regulation of the IAP genes in ovarian cell transformation have not been examined previously. Ovarian epithelial cancer is in part a result of suppressed apoptosis of ovarian surface epithelial cells. The effectiveness of certain chemotherapeutic agents rests on their ability to induce cell death. The loss of responsiveness of the cells to these agents is due to a desensitization of the apoptotic process to these agents. The regulation of ovarian epithelial

- 33 -

cell apoptosis involves changes in the expression of IAP genes and post-translational modification/processing of the IAP gene products.

We have conducted experiments and now believe that IAPs play a key role in maintaining the normal growth of ovarian surface epithelial cells and that the overexpression
5 of these genes leads to cellular transformation. Furthermore, we have discovered that the effectiveness of chemotherapeutic agents in the treatment of this form of malignancy rests upon their ability to suppress the expression of the IAP genes. By seeking to control the regulation of the IAP genes in human ovarian epithelial cancer cells we have provided a rational approach for the development of new chemotherapeutics for patients both responsive
10 and resistant to current cancer drugs. Similarly, assays designed to detect compounds which decrease IAP biological activity provide a rational method for drug discovery.

Methods.

a) Human Ovarian Epithelial Cancer Cell Culture

Cisplatin-sensitive (OV2008) and cisplatin-resistant (C13) human ovarian epithelial
15 cells were cultured in a chemically-defined medium at 37°C for up to 48 hours in the presence or absence of TGFβ (20 ng/ml), taxol (0 - 1.0 μM) or cisplatin (0 - 30 μM). At the end of the culture period, cells were either fixed for immunocytochemistry and TUNEL analyses, or snap frozen for subsequent extraction for IAP mRNA and proteins analyses.

b) Identification of Cell Death

20 For nuclear staining, human ovarian epithelial cancer cells were fixed (4% formalin in PBS; 10 min., room temp.), washed in PBS, resuspended in Hoescht 33248 stain (0.1 μg/ml PBS, 10 min) washed again and spotted onto slides for microscopy. Nuclear staining was observed and photographed using a Zeiss fluorescent microscope equipped with an FITC filter. Apoptotic cells were identified by typical nuclear morphology, and counted using
25 randomly selected fields and numbered photographic slides to avoid bias during counting.

For quantitation of DNA ladders, cellular DNA was extracted using the Qiagen Blood kit (Qiagen Inc., Chatsworth, CA). DNA was quantified by ethidium bromide fluorescence. DNA (0.5 μg) was then end labelled by incubating (20 min., room temp.) with Klenow enzyme (2 U in 10 mM Tris plus 5 mM MgCl₂) and 0.1 μCi [α³²P]dCTP. Unincorporated

- 34 -

nucleotides were removed with the Qiagen nucleotide removal kit and samples were resolved by Tris-acetate-EDTA agarose (1.8%) gel electrophoresis. The gel was then dried (2 hours, no heat) and exposed to a Bio-Rad phosphorimager screen to densitometrically quantify low molecular weight DNA (<15 kilo base-pairs), and subsequently to X-ray film at -80°C.

5 For *in situ* TUNEL labelling of apoptotic cells to identify cell death, the *in situ* cell death detection kit (Boehringer-Mannheim, Indianapolis, IN) was used, according to manufacturer's instructions. Slides prepared for histology were treated (20 min. at 37°C) with terminal transferase in the presence of FITC-conjugated dUTP.

c) Western Blot Analyses for IAPs

10 Protein extracts were prepared from human surface epithelial cancer cells sonicated (8 sec/cycle, 3 cycles) on ice in sucrose buffer (0.25 M sucrose, 0.025 M NaCl, 1 mM EGTA and 15 mM Tris-HCl pH 6.8, supplemented with 1 mM PMSF, 2 µg/ml of leupeptin and 5 µg/ml of aprotinin. The sonicates were centrifuged at 13,000xg for 10 min., the supernatants were collected and stored at -20°C until electrophoretic analyses were performed. Protein
15 concentration was determined by Bio-Rad Protein Assay. Proteins (10-30 µg) were resolved by one-dimensional SDS-PAGE, and electrophoretically transferred to nitrocellulose membrane. Membranes were blocked with 5% non-fat milk, and subsequently incubated with rabbit polyclonal antibody for IAP [anti-human HIAP-2ΔE (960529; 1:1000 dilution), anti-human NAIP E1.0 (951015; 1:1000 dilution) or anti-human XIAP (1:1000 dilution)]
20 diluted in TBST (10 mM Tris-buffered saline, 0.1% Tween-20, pH7.5) containing 5% milk. An ECL kit was used to visualize immunopositive protein (Amersham Intl., Arlington Heights, IL).

d) Northern Blots for IAP mRNAs

Total RNA from ovarian surface epithelial cancer cells by using RNeasy Kit
25 (Qiagen). The RNA samples (10-15 µg) were quantified spectrophotometrically and size-fractionated by electrophoresis on formaldehyde-agarose gels (1.1%) containing 1 µg/ml ethidium bromide to confirm even loading of RNA samples and adequate separation of 28S and 18S ribosomal bands. The RNAs bands were blotted onto a nylon membrane and cross-linked by UV light. Membranes were prehybridized in 50% formamide, saline sodium citrate

- 35 -

(SSC; 750 mM NaCl, 75 mM sodium citrate), 1X Denhardt's solution, 1% SDS, 4 mM EDTA and 100 µg/ml sheared salmon sperm DNA for 4 hours at 42°C. Hybridization was performed overnight at 42 °C with 20 million cpm of ³²P-labelled IAP cDNA probes (rat NAIP, rat XIAP or human HIAP-2) added to the prehybridization buffer. The membranes
5 were then washed twice with SSC (300 mM NaCl, 30 mM sodium citrate) in 0.1% SDS for 20 min at room temperature and twice with SSC (30 mM NaCl, 3 mM sodium citrate) in 0.1% SDS for 20 min at 55°C and exposed to X-ray film at -80°C for visualization. Densitometric analysis of various IAPs and 28S rRNA band was performed with the Image Analysis Systems from Bio-Rad Laboratories. Data were normalized by the respective 28S
10 and expressed as a percentage of the control (defined as 100%).

Results

We observed the following.

1. Cisplatin induced a concentration-dependent increase in the incidence of apoptosis in cisplatin-sensitive (OV2008) but to a lesser extent in -resistant (C13) human ovarian
15 epithelial cells in vitro (Fig. 20). Similarly, Taxol also induced apoptosis in OV2008 cells, but to a lesser extent in the C13 cells (Fig. 21).
2. Basal XIAP and HIAP-2 protein contents were markedly higher in cisplatin-sensitive than -resistant cells. Taxol (0.04-1.0 µM) decreased XIAP and HIAP-2 protein levels in a concentration-dependent manner, the response being more pronounced in sensitive than
20 resistant cells (Fig. 22). A lower molecular weight (approx. 45 kDa) immunoreactive fragment of HIAP-2 was also evident in both the sensitive and resistant cells. The content of this fragment was increased in the C13 cells but decreased in OV2008 cells by Taxol (Fig. 22).
3. Whereas Taxol (0.2 µM) marked suppressed HIAP-2 mRNA abundance in cisplatin-
25 sensitive cells (approx. 80%), it was ineffective in the resistant cells (Fig. 23).
4. TGFβ (20ng/ml) induced apoptosis in OV2008 but not in C13. Although its influence on XIAP protein content in cisplatin-resistant cells was only marginal, it markedly suppressed

- 36 -

the protein level of this IAP in the cisplatin-sensitive cells (Fig. 24A, 24B). TGF β (20 ng/ml) also decreased HIAP-2 mRNA in OV2008 but not C13 cells (Fig. 23).

Significant observations and possible applications.

Induction of apoptosis in human ovarian epithelial cancer cell by Taxol was
5 accompanied by suppressed IAP gene expression. Eventual loss of sensitivity of the cells to the chemotherapeutic agent may be associated with the decreased ability of the cell to express IAP genes. In drug-resistant cells, the decreased HIAP-2 protein content (in the face of an absence of noticeable change in HIAP-2 mRNA abundance) in the presence of Taxol was accompanied an increase in the intensity of a 45 kDa immunoreactive HIAP-2 protein band.
10 These observations lead us to believe that the 45 kDa protein is a proteolytic product of HIAP-2 and plays a role in the development of drug resistance. In addition, the sensitivity of the IAP family in these ovarian cancer cells to Taxol suggest possible novel sites for gene targeting in the development of new chemotherapeutic agents for the treatment of human ovarian epithelial cell cancer.

15 **EXAMPLE 4: Accumulation of a 26 kDa Cleavage Protein in Astrocytoma Cells**

Identification of a 26 kDa Cleavage Protein

A total protein extract was prepared from Jurkat and astrocytoma cells by sonicating them (X3 for 15 seconds at 4°C) in 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 1 mM PMSF, 1 µg/ml aprotinin, and 5 mM benzamidine. Following sonication, the samples were
20 centrifuged (14,000 RPM in a micro centrifuge) for five minutes. 20 µg of protein was loaded per well on a 10% SDS-polyacrylamide gel, electrophoresed, and electroblotted by standard methods to PVDF membranes. Western blot analysis, performed as described previously, revealed that the astrocytoma cell line (CCF-STTG1) abundantly expressed an anti-xiap reactive band of approximately 26 kDa, despite the lack of an apoptotic trigger
25 event (Fig. 12). In fact, this cell line has been previously characterized as being particularly resistant to standard apoptotic triggers.

A 26 kDa XIAP-reactive band was also observed under the following experimental conditions. Jurkat cells (a transformed human T cell line) were induced to undergo apoptosis by exposure to an anti-Fas antibody (1 µg/ml). Identical cultures of Jurkat cells were

- 37 -

exposed either to: (1) anti-Fas antibody and cycloheximide (20 µg/ml), (2) tumor necrosis factor alpha (TNF-α, at 1,000 U/ml), or (3) TNF-α and cycloheximide (20 µg/ml). All cells were harvested 6 hours after treatment began. In addition, as a negative control, anti-Fas antibody was added to an extract after the cells were harvested. The cells were harvested in
5 SDS sample buffer, electrophoresed on a 12.5% SDS polyacrylamide gel, and electroblotted onto PVDF membranes using standard methods. The membranes were immunostained with a rabbit polyclonal anti-XIAP antibody at 1:1000 for 1 hour at room temperature. Following four 15 minute washes, a goat anti-rabbit antibody conjugated to horse-radish peroxidase was applied at room temperature for 1 hour. Unbound secondary antibody was washed away, and
10 chemiluminescent detection of XIAP protein was performed. The Western blot revealed the presence of the full-length, 55 kDa XIAP protein, both in untreated and treated cells. In addition, a novel, approximately 26 kDa XIAP-reactive band was also observed in apoptotic cell extracts, but not in the control, untreated cell extracts (Fig. 13).

Cleavage of XIAP occurs in a variety of cell types, including other cancer cell lines
15 such as HeLa. The expression of the 26 kDa XIAP cleavage product was demonstrated in HeLa cells as follows. HeLa cells were treated with either: (1) cyclohexamide (20 µg/ml), (2) anti-Fas antibody (1 µg/ml), (3) anti-Fas antibody (1 µg/ml) and cyclohexamide (20 µg/ml), (4) TNFα (1,000 U/ml), or (5) TNFα (1,000 U/ml) and cyclohexamide (20 µg/ml). All cells were harvested 18 hours after treatment began. As above, anti-Fas antibody was
20 added to an extract after the cells were harvested. HeLa cells were harvested, and the Western blot was probed under the same conditions as used to visualize XIAP-reactive bands from Jurkat cell samples. A 26 kDa XIAP band was again seen in the apoptotic cell preparations (Fig. 14). Furthermore, the degree of XIAP cleavage correlated positively with cellular exposure to apoptotic triggers. Treatment of HeLa cells with cycloheximide or
25 TNFα alone caused only minor apoptosis, and little cleavage product was observed. If the cells were treated with the anti-Fas antibody, a greater amount of cleavage product was apparent. These data indicate that XIAP is cleaved in more than one cell type and in response to more than one type of apoptotic trigger.

Time Course of Expression

- 38 -

The time course over which the 26 kDa cleavage product accumulates was examined by treating HeLa and Jurkat cells with anti-Fas antibody (1 µg/ml) and harvesting them either immediately, or 1, 2, 3, 5, 10, or 22 hours after treatment. Protein extracts were prepared and Western blot analysis was performed as described above. Both types of cells accumulated increasing quantities of the 26 kDa cleavage product over the time course examined (Figs. 15A and 15B).

Subcellular Localization of the 26 kDa XIAP Cleavage Product

In order to determine the subcellular location of the 26 kDa cleavage product, Jurkat cells were induced to undergo apoptosis by exposure to anti-Fas antibody (1 µg/ml) and were then harvested either immediately, 3 hours, or 7 hours later. Total protein extracts were prepared, as described above, from cells harvested at each time point. In order to prepare nuclear and cytoplasmic cell extracts, apoptotic Jurkat cells were washed with isotonic Tris buffered saline (pH 7.0) and lysed by freezing and thawing five times in cell extraction buffer (50 mM PIPES, 50 mM KCl, 5 mM EGTA, 2 mM MgCl₂, 1 mM DTT, and 20 µM cytochalasin B). Nuclei were pelleted by centrifugation and resuspended in isotonic Tris (pH 7.0) and frozen at -80°C. The cytoplasmic fraction of the extract was processed further by centrifugation at 60,000 RPM in a TA 100.3 rotor for 30 minutes. Supernatants were removed and frozen at -80°C. Samples of both nuclear and cytoplasmic fractions were loaded on a 12.5% SDS-polyacrylamide gel, and electroblotted onto PVDF membranes. Western blot analysis was then performed using either an anti-CPP32 antibody (Transduction Laboratories Lexington, KY; Fig. 16A) or the rabbit anti-XIAP antibody described above (Fig. 16B).

The anti-CPP32 antibody, which recognizes the CPP32 protease (also known as YAMA or Apopain) partitioned almost exclusively in the cytoplasmic fraction. The 55 kDa XIAP protein localized exclusively in the cytoplasm of apoptotic cells, in agreement with the studies presented above, where XIAP protein in normal, healthy COS cells was seen to localize, by immunofluorescence microscopy, to the cytoplasm. In contrast, the 26 kDa cleavage product localized exclusively to the nuclear fraction of apoptotic Jurkat cells.

Taken together, these observations suggest that the anti-apoptotic component of XIAP could be the 26 kDa cleavage product, which exerts its influence within the nucleus.

- 39 -

In vitro Cleavage of XIAP protein and Characterization of the Cleavage Product

For this series of experiments, XIAP protein was labeled with ^{35}S using the plasmid pcDNA3-6myc-XIAP, T7 RNA polymerase, and a coupled transcription/translation kit (Promega, Madison, WI) according to the manufacturer's instructions. Radioactively labeled

5 XIAP protein was separated from unincorporated methionine by column chromatography using Sephadex G-50TM. In addition, extracts of apoptotic Jurkat cells were prepared following treatment with anti-Fas antibody (1 $\mu\text{g}/\text{ml}$) for three hours. To prepare the extracts, the cells were lysed in Triton X-100 buffer (1% Triton X-100, 25 mM Tris HCl) on ice for two hours and then microcentrifuged for 5 minutes. The soluble extract was retained

10 (and was labeled TX100). Cells were lysed in cell extraction buffer with freeze/thawing. The soluble cytoplasmic fraction was set aside (and labeled CEB). Nuclear pellets from the preparation of the CEB cytoplasmic fraction were solubilized with Triton X-100 buffer, microcentrifuged, and the soluble fractions, which contains primarily nuclear DNA, was retained (and labeled CEB-TX100). Soluble cell extract was prepared by lysing cells with

15 NP-40 buffer, followed by microcentrifugation for 5 minutes (and was labeled NP-40). *In vitro* cleavage was performed by incubating 16 μl of each extract (CEB, TX-100, CEB-TX100, and NP-40) with 4 μl of *in vitro* translated XIAP protein at 37°C for 7 hours. Negative controls, containing only TX100 buffer or CEB buffer were also included. The proteins were separated on a 10% SDS-polyacrylamide gel, which was dried and exposed to

20 X-ray film overnight.

In vitro cleavage of XIAP was apparent in the CEB extract. The observed molecular weight of the cleavage product was approximately 36 kDa (Fig. 17). The 10 kDa shift in the size of the cleavage product indicates that the observed product is derived from the amino-terminus of the recombinant protein, which contains six copies of the myc epitope (10 kDa).

25 It thus appears that the cleavage product possesses at least two of the BIR domains, and that it is localized to the nucleus.

EXAMPLE 5: CHARACTERIZATION OF IAP ACTIVITY AND INTRACELLULAR LOCALIZATION STUDIES

The ability of IAPs to modulate apoptosis can be defined *in vitro* systems in which

30 alterations of apoptosis can be detected. Mammalian expression constructs carrying IAP

- 40 -

cDNAs, which are either full-length truncated, or antisense constructs can be introduced into cell lines such as CHO, NIH 3T3, HL60, Rat-1, or Jurkat cells. In addition, SF21 insect cells may be used, in which case the IAP gene is preferentially expressed using an insect heat shock promoter. Following transfection, apoptosis can be induced by standard methods, which include serum withdrawal, or application of staurosporine, menadione (which induces apoptosis via free radical formation), or anti-Fas antibodies. As a control, cells are cultured under the same conditions as those induced to undergo apoptosis, but either not transfected, or transfected with a vector that lacks an IAP insert. The ability of each IAP related construct to inhibit or enhance apoptosis upon expression can be quantified by calculating the survival index of the cells, *i.e.*, the ratio of surviving transfected cells to surviving control cells. These experiments can confirm the presence of apoptosis inhibiting activity and, as discussed below, can also be used to determine the functional region(s) of an IAP which may be employed to achieve enhancement of apoptosis. These assays may also be performed in combination with the application of additional compounds in order to identify compounds that enhance apoptosis via IAP expression.

EXAMPLES 6: CELL SURVIVAL FOLLOWING TRANSFECTION WITH IAP CONSTRUCTS AND INDUCTION OF APOPTOSIS

Specific examples of the results obtained by performing various apoptosis suppression assays are shown in Figs. 10A to 10D. For example, CHO cell survival following transfection with one of six constructs and subsequent serum withdrawal is shown in Fig. 10A. The cells were transfected using Lipofectace™ with 2 µg of one of the following recombinant plasmids: pCDNA36myc-xiap (XIAP), pCDNA3-6myc-hiap-1 (HIAP-1), pCDNA3-6myc-hiap-2 (HIAP-2), pCDNA3-bcl-2 (BCL-2), pCDNA3-HA-smn (SMN), and pCDNA3-6myc (6-myc). Oligonucleotide primers were synthesized to allow PCR amplification and cloning of the XIAP, HIAP-1, and HIAP-2 ORFs in pCDNA3 (Invitrogen). Each construct was modified to incorporate a synthetic myc tag encoding six repeats of the peptide sequence MEQKLISEEDL (SEQ ID NO: 17), thus allowing detection of myc-IAP fusion proteins via monoclonal anti-myc antiserum (Egan *et al.*, Nature 363: 45-51, 1993). Triplicate samples of cell lines in 24-well dishes were washed 5 times with serum free media and maintained in serum free conditions during the course of the experiment.

- 41 -

Cells that excluded trypan blue, and that were therefore viable, were counted with a hemocytometer immediately, 24 hours, 48 hours, and 72 hours, after serum withdrawal. Survival was calculated as a percentage of the initial number of viable cells. In this experiment and those presented in Figs. 10B and 10D, the percentage of viable cells shown represents the average of three separate experiments performed in triplicate, +/- average deviation.

The survival of CHO cells following transfection (with each one of the six constructs described above) and exposure to menadione is shown in Fig. 10B. The cells were plated in 24-well dishes, allowed to grow overnight, and then exposed to 20 μ M menadione for 1.5 hours (Sigma Chemical Co., St. Louis, MO). Triplicate samples were harvested at the time of exposure to menadione and 24 hours afterward, and survival was assessed by trypan blue exclusion.

The survival of Rat-1 cells following transfection (with each one of the six constructs described above) and exposure to staurosporine is shown in Fig. 10C. Rat-1 cells were transfected and then selected in medium containing 800 μ g/ml G418 for two weeks. The cell line was assessed for resistance to staurosporine-induced apoptosis (1 μ M) for 5 hours. Viable cells were counted 24 hours after exposure to staurosporine by trypan blue exclusion. The percentage of viable cells shown represents the average of two experiments, +/- average deviation.

The Rat-1 cell line was also used to test the resistance of these cells to menadione (Fig. 10D) following transfection with each of the six constructs described above. The cells were exposed to 10 μ M menadione for 1.5 hours, and the NUMBER of viable cells was counted 18 hours later.

EXAMPLE 7: COMPARISON OF CELL SURVIVAL FOLLOWING TRANSFECTION WITH FULL-LENGTH VS. PARTIAL IAP CONSTRUCTS

In order to investigate the mechanism whereby human IAPs, including XIAP, HIAP-1, and HIAP-2, afford protection against cell death, expression vectors were constructed that contained either: (1) full-length IAP cDNA (as described above), (2) a portion of an IAP gene that encodes the BIR domains, but not the RZF, or (3) a portion of an IAP gene that encodes the RZF, but not the BIR domains. Human and murine XIAP cDNAs were tested by

- 42 -

transient or stable expression in HeLa, Jurkat, and CHO cell lines. Following transfection, apoptosis was induced by serum withdrawal, application of menadione, or application of an anti-Fas antibody. Cell death was then assessed, as described above, by trypan blue exclusion. As a control for transfection efficiency, the cells were co-transfected with a β -gal expression construct. Typically, approximately 20% of the cells were successfully transfected.

When CHO cells were transiently transfected, constructs containing full-length human or mouse xiap cDNAs conferred modest but definite protection against cell death. In contrast, the survival of CHO cells transfected with constructs encoding only the BIR domains (*i.e.*, lacking the RZF domain) was markedly enhanced 72 hours after serum deprivation. Furthermore, a large percentage of cells expressing the BIR domains were still viable after 96 hours, at which time no viable cells remained in the control, *i.e.* non-transfected, cell cultures, and less than 5% of the cells transfected with the vector only, *i.e.*, lacking a cDNA insert, remained viable. Deletion of any of the BIR domains results in the complete loss of apoptotic suppression, which is reflected by a decrease in the percentage of surviving CHO cells to control levels within 72 hours of serum withdrawal.

Stable pools of transfected CHO cells, which were maintained for several months under G418 selection, were induced to undergo apoptosis by exposure to 10 μ M menadione for 2 hours. Among the CHO cells tested were those that were stably transfected with: (1) full-length murine XIAP cDNA (MIAP), (2) full-length XIAP cDNA (XIAP), (3) full-length BCL-2 cDNA (BCL-2), (4) cDNA encoding the three BIR domains (but not the RZF) of murine XIAP (BIR), and (5) cDNA encoding the RZF (but not BIR domains) of M-XIAP (RZF). Cells that were non-transfected (CHO) or transfected with the vector only (pcDNA3), served as controls for this experiment. Following exposure to 10 μ M menadione, the transfected cells were washed with phosphate buffered saline (PBS) and cultured for an additional 24 hours in menadione-free medium. Cell death was assessed, as described above, by trypan blue exclusion. Less than 10% of the non-transfected or vector-only transfected cells remained viable at the end of the 24 hour survival period. Cells expressing the RZF did not fare significantly better. However, expression of full-length murine XIAP, human XIAP, or BCL-2, and expression of the BIR domains, enhanced cell survival. When the concentration of menadione was increased from 10 μ M to 20 μ M (with all other conditions

of the experiment being the same as when 10 μ M menadione was applied), the percentage of viable CHO cells that expressed the BIR domain cDNA construct was higher than the percentage of viable cells that expressed either full-length murine XIAP or BCL-2.

EXAMPLE 8: ANALYSIS OF THE SUBCELLULAR LOCATION OF EXPRESSED
RZF AND BIR DOMAINS

The assays of cell death described above indicate that the RZF acts as a negative regulator of the anti-apoptotic function of IAPs. One way in which the RZF, and possibly other IAP domains, may exert their regulatory influence is by altering the expression of genes, whose products function in the apoptotic pathway.

10 In order to determine whether the subcellular locations of expressed RZF and BIR domains are consistent with roles as nuclear regulatory factors, COS cells were transiently transfected with the following four constructs, and the expressed polypeptide was localized by immunofluorescent microscopy: (1) pcDNA3-6myc-XIAP, which encodes all 497 amino acids of SEQ ID NO: 4, (2) pcDNA3-6myc-m-XIAP, which encodes all 496 amino acids of
15 mouse XIAP (SEQ ID NO: 10), (3) pcDNA3-6myc-mxiap-BIR, which encodes amino acids 1 to 341 of m-XIAP, and (4) pcDNA3-6myc-mxiap-RZF, which encodes amino acids 342-496 of murine XIAP. The cells were grown on multi-well tissue culture slides for 12 hours, and then fixed and permeabilized with methanol. The constructs used (here and in the cell death assays) were tagged with a human Myc epitope tag at the N-terminus. Therefore, a
20 monoclonal anti-Myc antibody and a secondary goat anti-mouse antibody, which was conjugated to FITC, could be used to localize the expressed products in transiently transfected COS cells. Full-length XIAP and MIAP were located in the cytoplasm, with accentuated expression in the peri-nuclear zone. The same pattern of localization was observed when the cells expressed a construct encoding the RZF domain (but not the BIR
25 domains). However, cells expressing the BIR domains (without the RZF) exhibited, primarily, nuclear staining. The protein expressed by the BIR domain construct appeared to be in various stages of transfer to the nucleus.

These observations are consistent with the fact that, as described below, XIAP is cleaved within T cells that are treated with anti-Fas antibodies (which are potent inducers of

apoptosis), and its N-terminal domain is translocated to the nucleus. As noted in Example 2, HIAP-2 appears to undergo a similar cleavage event.

EXAMPLE 9: TESTING OF ANTISENSE OLIGONUCLEOTIDES:

1. *Complete panel of adenovirus constructs.* The panel may consist of approximately four
5 types of recombinant virus. A) Sense orientation viruses for each of the IAP or NAIP open
reading frames: XIAP, HIAP-1, HIAP-2, and NAIP. These viruses are designed to
massively overexpress the recombinant protein in infected cells. B) Antisense orientation
viruses in which the viral promoter drives the synthesis of an mRNA of opposite polarity to
the IAP mRNA, thereby shutting off host cell synthesis of the targeted protein coding region.
10 XIAP, HIAP-1, HIAP-2, and NAIP "antisense" constructs required. C) Sub-domain
expression viruses. These constructs express only a partial IAP protein in infected cells. Our
results indicate that deletion of the zinc finger of XIAP renders the protein more potent in
protecting cell against apoptotic triggers. This data also indicates that expression of the zinc
finger alone will indicate apoptosis by functioning as a dominant-negative repressor of XIAP
15 function. XIAP-ΔZF and XIAP-ΔBIR viruses required. D) Control viruses. Functional
analysis of the IAPs requires suitable positive and negative controls for comparison. BCL-2
sense, BCL-2 antisense, p53 sense, and Lac Z (negative control) viruses may be utilized.
2. *Confirmation of recombinant adenovirus function.* Verification of the sense adenovirus
function involves infection of tissue culture cells and determination of protein expression
20 levels. We have performed western blot analysis of several of the recombinant adenoviruses,
including NAIP, XIAP and XIAP-ΔRZF. The remaining viruses may be readily
assessed for protein expression using the polyclonal IAP antibodies. Functional analysis of
the antisense viruses may be done at the RNA level using either northern blots of total RNA
harvested from infected tissue culture cells or ribonuclease protection assays. Western blot
25 analysis of infected cells will be used to determine whether the expressed antisense RNA
interferes with IAP expression in the host cell.
3. *Documentation that IAP overexpression results in increased drug resistance.* We have
optimized cell death assays to allow high through-put of samples with minimal sample

- 45 -

variation. Testing of the sense IAP adenoviruses for their ability to alter drug sensitivity of breast and pancreatic adenocarcinoma cell lines may be accomplished as follows. Cancer cell lines are infected with the recombinant viruses, cultured for 5 days, then subdivided into 24 well plates. Triplicate cell receive increasing concentrations of the anti-cancer drug under investigation. Samples are harvested at 24, 48, and 72 hours post exposure, and assayed for the number of viable cells in the well. The dose response curve is then compared to uninfected and control virus (both positive and negative) infected cells. One may document a dramatic increase in the relative resistance of the cancer cell lines when infected with the sense viruses, confirming our hypothesis that overexpression of the IAP proteins contributes to the anti-apoptotic phenotype of cancer cells. Initial experiments utilize the chemotherapeutic drugs doxorubicin and adriamycin.

4. Documentation that antisense IAP overexpression results in increased drug sensitivity.

Having confirmed that IAP overexpression renders cancer cell more resistant to chemotherapeutic drugs, one may examine whether the antisense adenoviruses render the same cells more sensitive. The effectiveness of antisense IAP viruses relative to antisense BCL-2 virus will also be assessed as a crucial milestone.

5. Identification of antisense oligonucleotides. Concomitant to the adenovirus work, we have designed a series of antisense oligonucleotides to various regions of each of the IAPs. A generally accepted model of how antisense oligonucleotides function proposes that the formation of RNA/DNA duplexes in the nucleus activates cellular RnaseH enzymes which then enzymatically degrade the mRNA component of the hybrid. Virtually any region of the mRNA can be targeted, and therefore choosing an appropriate sequence to target is somewhat empirical. Many factors, including secondary structure of the target mRNA and the binding affinity of the targeted sequence determine whether a particular oligonucleotide will be effective, necessitating several oligonucleotides for each IAP. Five oligonucleotides have been made for each IAP mRNA based on the available computer algorithms for predicting binding affinities and mRNA secondary structures. These and other oligonucleotides may be tested for their ability to target their respective mRNAs for degradation using northern blot analysis.

- 46 -

6. *Optimization of oligonucleotides.* A secondary round of oligonucleotides may be made when effective target regions have been identified. These oligonucleotides target sequences in the immediate vicinity of the most active antisense oligonucleotides identified using methods such as those provided above. A second round of testing by northern blot analysis
5 may be required.

7. *Testing antisense oligonucleotides in vitro.* Following successful identification and optimization of targeting oligonucleotides, one may test these in the tissue culture model system using the optimal cell lines such as those described in the cancer survey described herein. Experimental procedures may parallel those used in the recombinant antisense
10 adenovirus work. Negative control oligonucleotides with miss-match sequences are used to establish base line or non-specific effects. Assisted transfection of the oligonucleotides using cationic lipid carriers may be compared to unassisted transfection. Confirmation of the effectiveness of specific antisense oligonucleotides prompts synthesis of oligonucleotides with modified phosphodiester linkages, such as phosphorothioate or methylimino substituted
15 oligonucleotides. These may also be tested *in vitro*.

8. *Animal modeling of antisense oligonucleotide therapies.*
Animal modeling of the effectiveness of the antisense IAP approach is described here. Cell lines are routinely assessed for their tumorigenic potential in "nude" mice, a hairless strain of mouse that is immunocompromised (lacks a functional thymus), and thus extremely
20 susceptible to developing tumors. In the nude mouse assay, cancer cells are grown in tissue culture and then injected under the skin at multiple sites. The frequency with which these cells give rise to palpable tumors within a defined period of time provides an index of the tumorigenic potential of the cell line in the absence of interference by a functional immune system. Preliminary assessment of an antisense IAP therapeutic involves injection of cancer
25 cells infected with the recombinant adenoviruses (sense, antisense, and control viruses) under the skin, and the tumorigenic index compared to that of untreated cells. One may also use this model to assess the effectiveness of systemic administration of antisense oligonucleotides in increasing the efficacy of anti-cancer drugs in the nude mouse model. Phosphorothioate or methylimino substituted oligonucleotides will be assessed at this stage.

This type of antisense oligonucleotide has demonstrated enhanced cell permeability and slower clearance rates from the body in experimental animal models.

EXAMPLE 10: ADDITIONAL APOPTOSIS ASSAYS

Specific examples of apoptosis assays are also provided in the following references.

- 5 Assays for apoptosis in lymphocytes are disclosed by: Li *et al.*, Science 268: 429-431, 1995; Gibellini *et al.*, Br. J. Haematol. 89: 24-33, 1995; Martin *et al.*, J. Immunol. 152: 330-342, 1994; Terai *et al.*, J. Clin Invest. 87: 1710-1715, 1991; Dhein *et al.*, Nature 373: 438-441, 1995; Katsikis *et al.*, J. Exp. Med. 1815: 2029-2036, 1995; Westendorp *et al.*, Nature 375: 497-500, 1995; DeRossi *et al.*, Virology 198: 234-244, 1994.
- 10 Assays for apoptosis in fibroblasts are disclosed by: Vossbeck *et al.*, Int. J. Cancer 61: 92-97, 1995; Goruppi *et al.*, Oncogene 9: 1537-1544, 1994; Fernandez *et al.*, Oncogene 9: 2009-2017, 1994; Harrington *et al.*, EMBO J., 13: 3286-3295, 1994; Itoh *et al.*, J. Biol. Chem. 268: 10932-10937, 1993.

Assays for apoptosis in neuronal cells are disclosed by: Melino *et al.*, Mol. Cell.

- 15 Biol. 14: 6584-6596, 1994; Rosenbaum *et al.*, Ann. Neurol. 36: 864-870, 1994; Sato *et al.*, J. Neurobiol. 25: 1227-1234, 1994; Ferrari *et al.*, J. Neurosci. 1516: 2857-2866, 1995; Talley *et al.*, Mol. Cell. Biol. 15: 2359-2366, 1995; Walkinshaw *et al.*, J. Clin. Invest. 95: 2458-2464, 1995.

Assays for apoptosis in insect cells are disclosed by: Clem *et al.*, Science 254: 1388-

- 20 1390, 1991; Crook *et al.*, J. Virol. 67: 2168-2174, 1993; Rabizadeh *et al.*, J. Neurochem. 61: 2318-2321, 1993; Birnbaum *et al.*, J. Virol. 68: 2521-2528, 1994; Clem *et al.*, Mol. Cell. Biol. 14: 5212-5222, 1994.

EXAMPLE 11: CONSTRUCTION OF A TRANSGENIC ANIMAL

Characterization of IAP and NAIP genes provided information that necessary for

- 25 generation IAP and NAIP transgenic animal models to be developed by homologous recombination (for knockouts) or transfection (for expression of IAP or NAIP fragments, antisense nucleic acids, or increased expression of wild-type or mutant IAPs or NAIP). Such a model may be a mammalian animal, e.g., a mouse, and is useful for the identification of

- 48 -

cancer therapeutics alone or in combination with cancer inducing cells or agents, or when such mice are crossed with mice genetically predisposed to cancers.

The preferred transgenic animal overexpression in IAP or NAIP and has a predisposition to cancer. This mouse is particularly useful for the screening of potential
5 cancer therapeutics.

EXAMPLE 12: IAP OR NAIP PROTEIN EXPRESSION

IAP and NAIP genes and fragments thereof (*i.e.*, RZF fragments) may be expressed in both prokaryotic and eukaryotic cell types. If an IAP or NAIP fragment enhances apoptosis, it may be desirable to express that protein under control of an inducible promoter.

10 In general, IAPs and NAIP, and fragments thereof, may be produced by transforming a suitable host cell with all or part of the IAP-encoding or NAIP-encoding cDNA fragment that has been placed into a suitable expression vector.

Those skilled in the art of molecular biology will understand that a wide variety of expression systems may be used to produce the recombinant protein. The precise host cell
15 used is not critical to the invention, although cancer cells are preferable. The IAP protein may be produced in a prokaryotic host (*e.g.*, *E. coli*) or in a eukaryotic host (*e.g.*, *S. cerevisiae*, insect cells such as Sf21 cells, or mammalian cells such as COS-1, NIH 3T3, or HeLa cells, or other highly proliferative cell types). These cells are publically available, for example, from the American Type Culture Collection, Rockville, MD; see also Ausubel *et al.*, *supra*).
20 *al.*, *supra*). The method of transduction and the choice of expression vehicle will depend on the host system selected. Transformation and transfection methods are described, *e.g.*, in Ausubel *et al.* (*supra*), and expression vehicles may be chosen from those provided, *e.g.*, in Cloning Vectors: A Laboratory Manual (P.H. Pouwels *et al.*, 1985, Supp. 1987).

Polypeptides of the invention, particularly short IAP fragments, can also be produced
25 by chemical synthesis (*e.g.*, by the methods described in Solid Phase Peptide Synthesis, 2nd ed., 1984 The Pierce Chemical Co., Rockford, IL). These general techniques of polypeptide expression and purification can also be used to produce and isolate useful IAP fragments or analogs, as described herein.

EXAMPLE 13: ANTI-IAP AND ANTI-NAIP ANTIBODIES

- 49 -

In order to generate IAP-specific and NAIP-specific antibodies, an IAP or NAIP coding sequence (*e.g.*, amino acids 180-276) can be expressed as a C-terminal fusion with glutathione S-transferase (GST; Smith *et al.*, Gene 67: 31-40, 1988). The fusion protein can be purified on glutathione-Sepharose beads, eluted with glutathione, and cleaved with thrombin (at the engineered cleavage site), and purified to the degree required to successfully immunize rabbits. Primary immunizations can be carried out with Freund's complete adjuvant and subsequent immunizations performed with Freund's incomplete adjuvant. Antibody titres are monitored by Western blot and immunoprecipitation analyses using the thrombin-cleaved IAP fragment of the GST-IAP and GST-NAIP fusion proteins. Immune sera are affinity purified using CNBr-Sepharose-coupled IAP protein. Antiserum specificity is determined using a panel of unrelated GST proteins (including GSTp53, Rb, HPV-16 E6, and E6-AP) and GST-trypsin (which was generated by PCR using known sequences).

As an alternate or adjunct immunogen to GST fusion proteins, peptides corresponding to relatively unique hydrophilic regions of IAP or NAIP may be generated and coupled to keyhole limpet hemocyanin (KLH) through an introduced C-terminal lysine. Antiserum to each of these peptides is similarly affinity purified on peptides conjugated to BSA, and specificity is tested by ELISA and Western blotting using peptide conjugates, and by Western blotting and immunoprecipitation using IAP or NAIP expressed as a GST fusion protein.

Alternatively, monoclonal antibodies may be prepared using the IAP or NAIP proteins described above and standard hybridoma technology (see, *e.g.*, Kohler *et al.*, Nature 256: 495, 1975; Kohler *et al.*, Eur. J. Immunol. 6: 511, 1976; Kohler *et al.*, Eur. J. Immunol. 6:292, 1976; Hammerling *et al.*, In Monoclonal Antibodies and T Cell Hybridomas, Elsevier, New York, NY, 1981; Ausubel *et al.*, *supra*). Once produced, monoclonal antibodies are also tested for specific IAP or NAIP recognition by Western blot or immunoprecipitation analysis (by the methods described in Ausubel *et al.*, *supra*).

Antibodies that specifically recognize IAPs or NAIP or fragments thereof, such as those described herein containing one or more BIR domains (but not a ring zinc finger domain), or that contain a ring zinc finger domain (but not a BIR domain) are considered useful in the invention. They may, for example, be used in an immunoassay to monitor IAP or NAIP expression levels or to determine the subcellular location of an IAP or NAIP (or

- 50 -

fragment thereof) produced by a mammal. Antibodies that inhibit the 26 kDa IAP cleavage product described herein (which contains at least one BIR domain) may be especially useful in inducing apoptosis in cells undergoing undesirable proliferation.

Preferably, antibodies of the invention are produced using IAP or NAIP sequence that
5 does not reside within highly conserved regions, and that appears likely to be antigenic, as analyzed by criteria such as those provided by the Peptide structure program (Genetics Computer Group Sequence Analysis Package, Program Manual for the GCG Package, Version 7, 1991) using the algorithm of Jameson and Wolf (CABIOS 4: 181, 1988). Specifically, these regions, which are found between BIR1 and BIR2 of all IAPs, are: from
10 amino acid 99 to amino acid 170 of HIAP-1, from amino acid 123 to amino acid 184 of HIAP-2, and from amino acid 116 to amino acid 133 of either XIAP or m-XIAP. These fragments can be generated by standard techniques, *e.g.*, by the PCR, and cloned into the pGEX expression vector (Ausubel *et al.*, *supra*). Fusion proteins are expressed in *E. coli* and purified using a glutathione agarose affinity matrix as described in Ausubel *et al.* (*supra*). In
15 order to minimize the potential for obtaining antisera that is non-specific, or exhibits low-affinity binding to IAP, two or three fusions are generated for each protein, and each fusion is injected into at least two rabbits. Antisera are raised by injections in series, preferably including at least three booster injections.

20 **EXAMPLE 14: IDENTIFICATION OF MOLECULES THAT MODULATE THE
EXPRESSION OR BIOLOGICAL ACTIVITY OF AN IAP OR NAIP GENE**

IAP and NAIP cDNAs facilitate the identification of molecules that decrease IAP or NAIP expression or otherwise enhance apoptosis normally blocked by these polypeptides. Such compounds are highly useful as, for example, chemotherapeutic agents to destroy a cancer cell, or to reduce the growth of a cancer cell, where the cancer cell is one, as is
25 described herein, with an elevated level of an IAP or NAIP polypeptide.

In one approach, candidate molecules are added, in varying concentration, to the culture medium of cells expressing IAP or NAIP mRNA. IAP or NAIP expression is then measured, for example, by Northern blot analysis (Ausubel *et al.*, *supra*) using an IAP or NAIP cDNA, or cDNA fragment, as a hybridization probe. The level of IAP or NAIP
30 expression in the presence of the candidate molecule is compared to the level of IAP or NAIP

- 51 -

expression in the absence of the candidate molecule, all other factors (*e.g.*, cell type and culture conditions) being equal.

The effect of candidate molecules on IAP- or NAIP-mediated apoptosis may, instead, be measured at the level of protein or the level of polypeptide fragments of IAP or NAIP polypeptides using the general approach described above with standard polypeptide detection techniques, such as Western blotting or immunoprecipitation with an IAP or NAIP-specific antibodies (for example, the antibodies described herein).

Compounds that modulate the level of a IAP or NAIP polypeptide may be purified, or substantially purified, or may be one component of a mixture of compounds such as an extract or supernatant obtained from cells (Ausubel *et al.*, *supra*). In an assay of a mixture of compounds, IAP or NAIP polypeptide expression is tested against progressively smaller subsets of the compound pool (*e.g.*, produced by standard purification techniques such as HPLC or FPLC) until a single compound or minimal number of effective compounds is demonstrated to modulate IAP or NAIP expression.

Compounds may also be screened for their ability to modulate the biological activity of an IAP or NAIP polypeptide by, for example, an ability to enhance IAP- or NAIP-mediated apoptosis. In this approach, the degree of apoptosis in the presence of a candidate compound is compared to the degree of apoptosis in its absence, under equivalent conditions. Again, the screen may begin with a pool of candidate compounds, from which one or more useful modulator compounds are isolated in a step-wise fashion. Apoptosis activity may be measured by any standard assay, for example, those described herein.

Another method for detecting compounds that modulate the expression or biological activity of an IAP or a NAIP polypeptide is to screen for compounds that interact physically with a given IAP polypeptide. These compounds may be detected by adapting two hybrid systems known in the art. These systems detect protein interactions using a transcriptional activation assay and are generally described by Gyuris *et al.* (Cell 75: 791-803, 1993) and Field *et al.* (Nature 340: 245-246, 1989), and are commercially available from Clontech (Palo Alto, CA). In addition, PCT Publication WO 95/28497 describes a two hybrid system in which proteins involved in apoptosis, by virtue of their interaction with BCL-2, are detected. A similar method may be used to identify proteins and other compounds that interact with IAP or NAIP polypeptides.

Compounds or molecules that function as modulators of IAP-mediated cell death may include peptide and non-peptide molecules such as those present in cell extracts, mammalian serum, or growth medium in which mammalian cells have been cultured. In addition, compounds previously known for their abilities to modulate apoptosis in cancer cells may be tested for an ability to modulate expression of an IAP molecule.

TABLE 2

OLIGONUCLEOTIDE PRIMERS FOR THE SPECIFIC
RT-PCR AMPLIFICATION OF IAP GENES

| 10 | IAP Gene | Forward Primer (nucleotide position*) | Reverse Primer (nucleotide position*) | Size of Product (bp) |
|----|----------|--|--|-------------------------|
| | h-XIAP | p2415 (876-896) | p2449 (1291-1311) | 435 |
| | m-XIAP | p2566 (458-478) | p2490 (994-1013) | 555 |
| | h-HIAP 1 | p2465 (827-847) | p2464 (1008-1038) | 211 |
| | m-HIAP 1 | p2687 (747-767) | p2684 (1177-1197) | 450 |
| 15 | HIAP2 | p2595 (1562-1585) | p2578 (2339-2363) | 801& 618@ |
| | m-HIAP2 | p2693 (1751-1772) | p2734 (2078-2100) | 349 |

* Nucleotide position as determined from Figs. 1-4 for each IAP gene

& PCR product size of hiap2a

@ PCR product size of hiap2b

20 **EXAMPLE 15: ROLE OF IAPs IN HUMAN OVARIAN CANCER RESISTANCE TO CISPLATIN**

Ovarian epithelial cancer cell apoptosis has been demonstrated to be involved in cisplatin-induced cell death (Havrilesky *et al.*, Obstet. Gynecol. 85: 1007-1010, 1995; Anthoney *et al.*, Cancer Res. 56: 1374-1381, 1996). The action of cisplatin is thought to involve the formation of inter and intra-strand DNA crosslinks (Sherman *et al.*, Science 230: 412-417, 1985) although the events leading to cell death after cisplatin treatment is unclear. If IAPs are indeed key elements in the regulation of apoptosis in ovarian cancer cells, one

- 53 -

would expect that down-regulation of this anti-apoptotic protein would result in cell death. To test this, cisplatin-sensitive human ovarian surface epithelial cells (OV2008) were infected with either adenoviral XIAP antisense, adenoviral HIAP-2 antisense, or the empty vector with LacZ (as control) for up to 60 hours, at which time changes in cell morphology, apoptotic cell number, cell viability, and total cell number were determined. The full length sense and antisense constructs of XIAP and HIAP-2 were prepared as briefly described hereafter. To construct the adenoviruses, the open reading frame for XIAP and HIAP-2 were PCR amplified with primers corresponding to the amino and carboxy terminus. These PCR products were cloned in the pCR2.1 vector (InvitroGen, Carlsbad, CA), and sequenced. The ORFs were then excised with EcoRI digestion, blunt ended with Klenow fragment, and ligated into SmaI digested pAdex1CAwt cosmid DNA. Packaging was performed with Promega (Madison, WI) cosmid packaging extracts and used to infect *E. coli*. Colonies were picked and screened for the presence of the insert in both the sense and antisense orientation relative to the chicken B-actin (CA) promoter. CsCl purified cosmid DNA was co-transfected with wild-type adenovirus DNA, which contains the terminal protein complexed to the ends of the DNA. Wild type adenovirus DNA was cut with NsiI such that only homologous recombinant with the cosmid DNA generated infectious adenovirus DNA. The final recombinant adenovirus contains a linear, double stranded genome of 44,820 bp plus the insert size (approximately 1,500 for XIAP, approximately 1,800 for HIAP-2).

Cisplatin-sensitive (OV2008) and cisplatin-resistant (C13) ovarian epithelial cancer cells were infected with adenovirus [multiplicity of infection (MOI) = 5 (1X); MOI = 10 (2X)] containing antisense XIAP or HIAP-2 cDNA, or vector (control) for 60 hours. Cells were then trypsinized and total cell number was determined with haemocytometry while cell viability was determined by the trypan blue dye exclusion test. XIAP antisense infection of OV2008 cells significantly increased the percentage of dead cells compared to control (vector, $p < 0.001$), as determined by trypan blue exclusion tests (Fig. 25, top left panel). Although there appeared also to be a slight increase in percentage of dead cells with HIAP-2 antisense infection of OV2008 cells, it was not statistically significant (Fig. 25, top left panel; $p > 0.05$). Infection of the cisplatin-resistant variant of OV2008 cells (C13) with antisense of XIAP but not of Hisp-2 also significantly, though to a lesser extent, decreased cell viability (Fig. 25, top right panel). The cell death induced in both OV2008 and C13 by XIAP

antisense was also accompanied by decreases in total cell number, with the effect of the antisense infection being more pronounced in the cisplatin-sensitive cells (Fig.25, bottom two panels).

In addition, 60 hours of adenoviral XIAP antisense infection of OV2008 decreased XIAP protein content and induced extensive cell detachment, as is shown in Fig. 26A (black arrows in left "b" photograph). Nuclear fragmentation (Fig. 26B, white arrows in photographs "b" and "d") and increased the number of apoptotic cells as well as the abundance of apoptotic bodies (Fig 26B: photographs "b" and "d" compared to "a" and "c") is also induced in OV2008 cells following 60 hours of infection with adenovirus XIAP antisense. For nuclear staining, cells were fixed in 4% formalin (in PBS, room temp., 10 min.) and washed in PBS. The washed cells were then resuspended in Hoechst staining solution (0.1 µg Hoechst 33248/ml PBS, 10 min.), washed again, and spotted onto slides for microscopy. Nuclear staining was observed and photographed using a Zeiss fluorescence microscope. Cells with typical apoptotic nuclear morphology were identified and counted, using randomly selected fields and numbered photographic slides to avoid bias during counting. Analysis of variance indicated that there was highly significant effects of the antisense on XIAP protein content ($p<0.001$; Fig. 26D and 26E) and apoptosis ($p<0.001$; Fig 26C). Indeed, infection of these cells with a higher titre of the adenoviral anti-sense (MOI=10 (2X)) further increased the number of cells undergoing apoptosis (Fig. 26C).

To study whether IAP expression is the target for the chemotherapeutic action of cisplatin, OV2008 cells were cultured in the absence and presence of cisplatin (10-30 µM) for 24 hours, apoptosis and XIAP and HIAP-2 expression were assessed morphologically and by Western analysis, respectively. Like adenoviral XIAP antisense infection, the presence of cisplatin induced morphologic feature of apoptosis in OV2008 cells, including decreased cell volume, chromatin condensation and nuclear fragmentation (Fig. 27A, left two photographs), and apoptotic low molecular weight DNA fragmentation (Fig. 27B), and was accompanied by decreased IAP expression (Figs. 28A and 28B). The increase of apoptotic cell number in response to cisplatin was also concentration-dependent and was significant (50% vs. 2%; $p<0.05$) even at a concentration of 10 µM cisplatin (Fig. 27C).

As shown in Figs. 28A and 28B, although both XIAP and HIAP-2 are present in the cisplatin-sensitive human ovarian surface epithelial cancer cell line OV2008 (protein sizes

- 55 -

55kDa and 68 kDa, respectively), their expression were down-regulated by cisplatin in a concentration-dependent manner. XIAP appearing more responsive to the anti-cancer agent. While XIAP protein content was decreased by almost 80% ($p<0.01$) in the presence 20 μ M cisplatin, the decrease of HIAP-2 protein content was not suppressed by cisplatin (Figs. 28A and 28B).

The expression of XIAP and HIAP-2 in C13, the cisplatin-resistant variant of OV2008, was not suppressed by cisplatin (Figs. 28A and 28B), and no morphologic and biochemical changes characteristic of apoptosis could be detected (Figs. 27A and 27B). Although XIAP and HIAP-2 contents in C13 appeared to be higher in the presence of the anti-cancer agent, the differences were statistically non-significant ($p>0.05$). Time course experiments on IAP expression demonstrated that the suppression of XIAP and HIAP-2 protein levels in OV2008 by cisplatin was time-dependent; a significant decrease was observed between 12-24 hours of culture (Figs. 29A and 29B). Expression of XIAP and HIAP-2 in C13 cells was not influenced by cisplatin, irrespective of the duration of treatment.

To determine if the observed XIAP responses in OV2008 and C13 cells were specific to this pair of cell lines, the influence of cisplatin in vitro on XIAP and HIAP-2 protein content in another cisplatin-sensitive ovarian surface epithelial cancer cell line (A2780s) and its cisplatin-resistant variant (A2780cp) was studied (Figs. 30A and 30B). Interestingly, whereas HIAP-2 expression in both the sensitive and resistant cells was not significantly altered by the presence of the cisplatin (30 μ M; Fig. 30B), XIAP protein content was decreased in A2780s (as in OV2008 cells) and not significantly altered in A2780cp (as in C13 cells) in the presence of the chemotherapeutic agent. Taken together, these data suggest that the apoptotic responsiveness of ovarian cancer cells to cisplatin may be related to the ability of the chemotherapeutic agent to down-regulate XIAP expression and that HIAP-2 may play a minor or no role in cisplatin-induced apoptosis.

To determine if XIAP expression is indeed the an important determinant in chemoresistance in human ovarian surface epithelial cancer, the influence of cisplatin on XIAP protein content and apoptosis in OV2008 cells following adenoviral XIAP sense infection was investigated. While cisplatin reduced XIAP protein content in OV2008 cells infected with the empty vector (Figs. 31C and 31D, vector plus cisplatin), overexpression of the protein with adenoviral sense XIAP cDNA 48 hrs prior to treatment with the

- 56 -

chemotherapeutic agent *in vitro* attenuated the cisplatin effects not only on XIAP protein expression (Figs. 31C and 31D) but also apoptotic nuclear fragmentation (Fig. 31A, "d" compared to "c") and number of apoptotic cells (Fig. 31B), suggesting that XIAP may be an important element in human ovarian epithelial cancer chemoresistance.

5 The *in vitro* studies with ovarian epithelial cancer cell lines strongly suggest an important role of IAPs, particularly of XIAP, in the control of apoptosis and tumor progression in human ovarian cancer. To determine if indeed IAPs are expressed in ovarian carcinoma and thus of clinical relevance, XIAP and HIAP-2 were immunolocalized in human ovarian surface epithelial tumors obtained as pathological samples from patients during
10 surgical debulking, using polyclonal antibodies (rabbit polyclonal anti-XIAP and HIAP-2 antibodies were prepared by immunization with human XIAP and HIAP-2 GST fusion protein) against human XIAP and HIAP-2, respectively (Figs. 32C and 32D, respectively). In addition, *in situ* TUNEL (described in Gavrieli *et al.*, J. Cell. Biol. 119: 493-501, 1992) and immunohistochemistry for PCNA (proliferating cell nuclear antigen: an auxiliary protein
15 of DNA polymerase α highly expressed as the G1/S interphase) were performed to examine if and how the expression of these IAPs relates to epithelial cell apoptosis and /or proliferation. Ovarian epithelial tumors exhibited considerable cellular heterogeneity (Fig. 32A) and PCNA positive cells were evident throughout the nucleus in the tumor section (Fig. 32B). In general, most of the cells were TUNEL negative (Fig. 32A), and the expression of
20 XIAP and HIAP-2 was highly correlated to the proliferative state of the cells and inversely related to epithelial cell death. XIAP and HIAP-2 immunoreactivity (Figs. 32C and 32D, respectively) specifically localized in the cytoplasm or the perinuclear region was highest in proliferatively active cells (PCNA positive) and was low or absent in apoptotic cells (TUNEL positive) occasionally found in the tumor specimens.

25 EXAMPLE 16: ADDITIONAL CANCER THERAPIES

Given the increased proliferation rate of cancer cells, it is preferable in anti-cancer therapeutic regimes to initiate treatment with an anti-cancer agent that will successfully inhibit the growth of the particular cancer of interest. One method to detect such an agent is
30 to excise proliferative cells from the cancer of interest, and determine the level of expression

- 57 -

and/or level of biological activity of each individual IAP or NAIP polypeptide, and compare these levels to the levels of these polypeptide in a similar cell type from an unaffected individual. For example, if an human female individual has breast cancer (or a neoplasm suspected of being cancerous), cells from the cancer collected, for example, during a biopsy
5 of the cancer, can be isolated and, if necessary, propagated in culture. The cells can then be analyzed for level of expression and/or level of biological activity of all of the IAP and NAIP polypeptides in the cell. The expression levels and/or biological activity levels of these polypeptides from the proliferating cells can be compared to the levels of expression and/or biological activity of these polypeptides from normal, healthy cells from a human female
10 individual. Preferably, the comparison is made between on affected (*i.e.*, abnormally proliferating) and healthy cells of the same individual (*e.g.*, cells taken from healthy breast tissue from the individual being tested. The level of expression and/or biological activity of each polypeptide in the affected cells is compared to its counterpart in the healthy cells. Any increase in any (or all) of the IAP or NAIP polypeptides is detected. The cancer is then
15 treated with a compound that decreases expression level or biological activity level of each particular elevated IAP or NAIP polypeptide. Methods for identifying such compounds are described above (see, *e.g.*, Example 14).

It will be understood that the individual undergoing such analysis and treatment may have already received treatment with an anti-cancer therapeutic agent. It will also be
20 understood that, in addition to targeting the levels of expression and/or biological activities of IAP and NAIP polypeptides, the anti-cancer compounds may also target these levels for other apoptosis-inhibiting polypeptides, such as BCL-2. For example, an individual with breast cancer whose proliferating cells have an increased level of XIAP compared to the level of XIAP in healthy breast cells may be treated with a compound (*e.g.*, cisplatin) plus a
25 compound that targets another IAP polypeptide, or that targets an NAIP polypeptide or a non-related apoptosis-inhibiting polypeptide, such as BCL-2).

One rapid method to determine expression levels of IAP and NAIP polypeptides is an ELISA assay using antibodies that specifically binds each of these polypeptides. Other methods include quantitative PCR and the various apoptosis assays described herein.

- 58 -

**EXAMPLE 17: ASSIGNMENT OF XIAP, HIAP-1, AND HIAP-2 TO
CHROMOSOMES XQ25 AND 11Q22-23 BY FLUORESCENCE IN SITU
HYBRIDIZATION (FISH)**

Fluorescence *in situ* hybridization (FISH) was used to identify the chromosomal
5 location of XIAP, HIAP-1 and HIAP-2.

A total of 101 metaphase spreads were examined with the XIAP probe, as described
above. Symmetrical fluorescent signals on either one or both homologs of chromosome
Xq25 were observed in 74% of the cells analyzed. Following staining with HIAP-1 and
HIAP-2 probes, 56 cells were analyzed and doublet signals in the region 11q22-23 were
10 observed in 83% of cells examined. The XIAP gene was mapped to Xq25 while the HIAP-1
and HIAP-2 genes were mapped at the border of 11q22 and 11q23 bands.

These experiments confirmed the location of the XIAP gene on chromosome Xq25.
No highly consistent chromosomal abnormalities involving band Xq25 have been reported so
far in any malignancies. However, deletions within this region are associated with a number
15 of immune system defects including X-linked lymphoproliferative disease (Wu *et al.*,
Genomics 17:163-170, 1993).

Cytogenetic abnormalities of band 11q23 have been identified in more than 50% of
infant leukemias regardless of the phenotype (Martinez-Climet *et al.*, Leukaemia 9: 1299-
1304, 1995). Rearrangements of the MLL Gene (mixed lineage leukemia or myeloid
20 lymphoid leukemia; Ziemann-van der Poel *et al.*, Proc. Natl. Acad. Sci. USA 88: 10735-
10739, 1991) have been detected in 80% of cases with 11q23 translocation, however patients
whose rearrangements clearly involved regions other than the MLL gene were also reported
(Kobayashi *et al.*, Blood 82: 547-551, 1993). Thus, the IAP genes may follow the BCL-2
paradigm, and would therefore play an important role in cancer transformation.

25 **Incorporation by Reference**

The following documents and all the references referred to herein are incorporated by
reference: U.S.S.N. 08/511,485, filed August 4, 1995; U.S.S.N. 08/576,956, filed December
22, 1995; PCT/IB96/01022, filed August 5, 1996; U.S.S.N. 60/017,354, filed April 26, 1996;
U.S.S.N. 60/030,931, filed November 15, 1996; U.S.S.N. 60/030,590, filed November 14,
30 1996; U.S.P.N. 5,576,208, issued November 19, 1996; and PCT Application No.

- 59 -

1B97/00142, filed January 17, 1997 claiming priority from UK 9601108.5, filed January 19, 1996.

Other Embodiments

In other embodiments, the invention includes use of any protein which is substantially
5 identical to a mammalian IAP polypeptides (Figs. 1-6; SEQ ID Nos: 3-14); such homologs
include other substantially pure naturally-occurring mammalian IAP proteins as well as
allelic variants; natural mutants; induced mutants; DNA sequences which encode proteins
and also hybridize to the IAP DNA sequences of Figs. 1-6 (SEQ ID NOS: 3-14) under high
stringency conditions or, less preferably, under low stringency conditions (*e.g.*, washing at
10 2X SSC at 40°C with a probe length of at least 40 nucleotides); and proteins specifically
bound by antisera directed to a IAP polypeptide. The term also includes chimeric
polypeptides that include a IAP portion.

The invention further includes use of analogs of any naturally-occurring IAP
polypeptide. Analogs can differ from the naturally-occurring IAP protein by amino acid
15 sequence differences, by post-translational modifications, or by both. Analogs of the
invention will generally exhibit at least 85%, more preferably 90%, and most preferably
95% or even 99% identity with all or part of a naturally occurring IAP amino acid sequence.
The length of sequence comparison is at least 15 amino acid residues, preferably at least 25
amino acid residues, and more preferably more than 35 amino acid residues. Modifications
20 include *in vivo* and *in vitro* chemical derivatization of polypeptides, *e.g.*, acetylation,
carboxylation, phosphorylation, or glycosylation; such modifications may occur during
polypeptide synthesis or processing or following treatment with isolated modifying enzymes.
Analogues can also differ from the naturally-occurring IAP polypeptide by alterations in
primary sequence. These include genetic variants, both natural and induced (for example,
25 resulting from random mutagenesis by irradiation or exposure to ethanemethylsulfate or by
site-specific mutagenesis as described in Sambrook, Fritsch and Maniatis, Molecular
Cloning: A Laboratory Manual, 2nd ed., CSH Press, 1989, or Ausubel *et al.*, *supra*). Also
included are cyclized peptides, molecules, and analogs which contain residues other than L-
amino acids, *e.g.*, D-amino acids or nonnaturally occurring or synthetic amino acids, *e.g.*, B
30 or y amino acids. In addition to full-length polypeptides, the invention also includes IAP

- 60 -

polypeptide fragments. As used herein, the term "fragment," means at least 20 contiguous amino acids, preferably at least 30 contiguous amino acids, more preferably at least 50 contiguous amino acids, and most preferably at least 60 to 80 or more contiguous amino acids. Fragments of IAP polypeptides can be generated by methods known to those skilled in the art or may result from normal protein processing (e.g., removal of amino acids from the nascent polypeptide that are not required for biological activity or removal of amino acids by alternative mRNA splicing or alternative protein processing events).

Preferable fragments or analogs used according to the methods of the invention are those which facilitate specific detection of an IAP nucleic acid or amino acid sequence in a sample to be diagnosed. Particularly useful IAP fragments for this purpose include, without limitation, the amino acid fragments shown in Table 2.

The methods of the invention may use antibodies prepared by a variety of methods. For example, the IAP or NAIP polypeptide, or antigenic fragments thereof, can be administered to an animal in order to induce the production of polyclonal antibodies. Alternatively, antibodies used as described herein may be monoclonal antibodies, which are prepared using hybridoma technology (see, e.g., Kohler *et al.*, Nature 256: 495-497, 1975; Kohler *et al.*, Eur. J. Immunol. 6: 511-519, 1976; Kohler *et al.*, Eur. J. Immunol. 6: 292-295, 1976; Hammerling *et al.*, In Monoclonal Antibodies and T Cell Hybridomas, Elsevier, NY, 1981; Harlow and Lane, Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, 1988). The invention features use of antibodies that specifically bind human or murine IAP or NAIP polypeptides, or fragments thereof. In particular the invention features "neutralizing" antibodies. By "neutralizing" antibodies is meant antibodies that interfere with any of the biological activities of IAP or NAIP polypeptides, particularly the ability of IAPs to inhibit apoptosis. The neutralizing antibody may reduce the ability of IAP polypeptides to inhibit polypeptides by, preferably 50%, more preferably by 70%, and most preferably by 90% or more. Any standard assay of apoptosis, including those described herein, by those incorporated by reference and those in the art, may be used to assess neutralizing antibodies.

In addition to intact monoclonal and polyclonal anti-IAP antibodies, the invention features use of various genetically engineered antibodies, humanized antibodies, and antibody fragments, including F(ab')₂, Fab', Fab, Fv and sFv fragments. Antibodies can be

- 61 -

humanized by methods known in the art, *e.g.*, monoclonal antibodies with a desired binding specificity can be commercially humanized (Scotgene, Scotland; Oxford Molecular, Palo Alto, CA). Fully human antibodies, such as those expressed in transgenic animals, are also features of the invention (Green *et al.*, Nature Genetics 7:13-21, 1994).

- 5 Ladner (U.S. Patent Nos. 4,946,778 and 4,704,692) describes methods for preparing single polypeptide chain antibodies. Ward *et al.* (Nature 341: 544-546, 1989) describe the preparation of heavy chain variable domains, which they term "single domain antibodies," which have high antigen-binding affinities. McCafferty *et al.* (Nature 348: 552-554, 1990) show that complete antibody V domains can be displayed on the surface of fd bacteriophage,
- 10 that the phage bind specifically to antigen, and that rare phage (one in a million) can be isolated after affinity chromatography. Boss *et al.* (U.S. Patent No. 4,816,397) describe various methods for producing immunoglobulins, and immunologically functional fragments thereof, which include at least the variable domains of the heavy and light chain in a single host cell. Cabilly *et al.* (U.S. Patent No. 4,816,567) describe methods for preparing chimeric
- 15 antibodies.

- 62 -

SEQUENCE LISTING

(1) GENERAL INFORMATION

(i) APPLICANT: University of Ottawa

5 (ii) TITLE OF THE INVENTION: DETECTION AND MODULATION OF
IAPS AND NAIP FOR THE DIAGNOSIS
AND TREATMENT OF PROLIFERATIVE
DISEASE

(iii) NUMBER OF SEQUENCES: 17

10 (iv) CORRESPONDENCE ADDRESS:
(A) ADDRESSEE: Clark & Elbing LLP
(B) STREET: 176 Federal Street
(C) CITY: Boston
(D) STATE: MA
15 (E) COUNTRY: USA
(F) ZIP: 02110

(v) COMPUTER READABLE FORM:
(A) MEDIUM TYPE: Diskette
(B) COMPUTER: IBM Compatible
20 (C) OPERATING SYSTEM: DOS
(D) SOFTWARE: FastSEQ for Windows Version 2.0

(vi) CURRENT APPLICATION DATA:
(A) APPLICATION NUMBER: PCT/---
(B) FILING DATE: 13-FEB-1998
25 (C) CLASSIFICATION:

(vi) PRIOR APPLICATION DATA:
(A) APPLICATION NUMBER: 08/800,929
(B) FILING DATE: 13-FEB-1997
(C) CLASSIFICATION:

- 63 -

(viii) ATTORNEY/AGENT INFORMATION:

- (A) NAME: Bieker-Brady, Kristina
(B) REGISTRATION NUMBER: 39,109
(C) REFERENCE/DOCKET NUMBER: 07891/009WO2

5 (ix) TELECOMMUNICATION INFORMATION:

- (A) TELEPHONE: 617-428-0200
(B) TELEFAX: 617-428-7045
(C) TELEX:

(2) INFORMATION FOR SEQ ID NO:1:

10 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 46 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

15 (ii) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(ix) FEATURE:

- (D) OTHER INFORMATION: Xaa at positions 2, 3, 4, 5,
6, 7, 9, 10, 11, 17, 18, 19, 20, 21, 23, 25, 30, 31, 32, 34, 35,
20 38, 39, 40, 41, 42, and 45 may be any amino acid. Xaa at position
8 is Glu or Asp. Xaa at positions 14 & 22 is Val or Ile.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Glu Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Lys Xaa Cys Met
1 5 10 15
25 Xaa Xaa Xaa Xaa Xaa Xaa Phe Xaa Pro Cys Gly His Xaa Xaa Xaa
20 25 30
Cys Xaa Xaa Cys Ala Xaa Xaa Xaa Xaa Xaa Cys Pro Xaa Cys
35 40 45

(2) INFORMATION FOR SEQ ID NO:2:

- 64 -

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 68 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(ix) FEATURE:

(D) OTHER INFORMATION: Xaa at positions 1, 2, 3,

6, 9, 10, 14, 15, 18, 19, 20, 21, 24, 30, 32, 33, 35, 37, 40,
 42, 43, 44, 45, 46, 47, 49, 50, 51, 53, 54, 55, 56, 57, 59, 60,
 61, 62, 64 and 66 may be any amino acid. Xaa at positions 13, 16 and
 17 may be any amino acid or may be absent.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

15 Xaa Xaa Xaa Arg Leu Xaa Thr Phe Xaa Xaa Trp Pro Xaa Xaa Xaa Xaa
 1 5 10 15
 Xaa Xaa Xaa Xaa Xaa Leu Ala Xaa Ala Gly Phe Tyr Tyr Xaa Gly Xaa
 20 25 30
 Xaa Asp Xaa Val Xaa Cys Phe Xaa Cys Xaa Xaa Xaa Xaa Xaa Trp
 20 35 40 45
 Xaa Xaa Xaa Asp Xaa Xaa Xaa Xaa Xaa His Xaa Xaa Xaa Xaa Pro Xaa
 50 55 60
 Cys Xaa Phe Val
 65

25 (2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 5232 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

30

- 65 -

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

| | | |
|----|--|------|
| | GAAAAGGTGG ACAAGTCCTA TTTTCAAGAG AAGATGACTT TTAACAGTTT TGAAGGATCT | 60 |
| | AAACTTTGTG TACCTGCAGA CATCAATAAG GAAGAAGAAT TTGTAGAAGA GTTTAATAGA | 120 |
| 5 | TTAAAACTT TTGCTAATTT TCCAAGTGGT AGTCCTGTTT CAGCATCAAC ACTGGCACGA | 180 |
| | GCAGGGTTTC TTTATACTGG TGAAGGAGAT ACCGTGCGGT GCTTTAGTTG TCATGCAGCT | 240 |
| | GTAGATAGAT GGCAATATGG AGACTCAGCA GTTGAAGAC ACAGGAAAGT ATCCCCAAAT | 300 |
| | TGCAGATTTA TCAACGGCTT TTATCTTGAA AATAGTGCCA CGCAGTCTAC AAATTCTGGT | 360 |
| | ATCCAGAATG GTCAGTACAA AGTTGAAAAC TATCTGGGAA GCAGAGATCA TTTTGCCTTA | 420 |
| 10 | GACAGGCCAT CTGAGACACA TGCAGACTAT CTTTGTAGAA CTGGGCAGGT TGTAGATATA | 480 |
| | TCAGACACCA TATACCCGAG GAACCCTGCC ATGTATAGTG AAGAAGCTAG ATTAAAGTCC | 540 |
| | TTTCAGAACT GGCCAGACTA TGCTCACCTA ACCCCAAGAG AGTTAGCAAG TGCTGGACTC | 600 |
| | TACTACACAG GTATTGGTGA CCAAGTGCAG TGCTTTTGTT GTGGTGGAAA ACTGAAAAAT | 660 |
| | TGGGAACCTT GTGATCGTGC CTGGTCAGAA CACAGGCGAC ACTTTCCTAA TTGCTTCTTT | 720 |
| 15 | GTTTTGGGCC GGAATCTTAA TATTCGAAGT GAATCTGATG CTGTGAGTTC TGATAGGAAT | 780 |
| | TTCCCAAATT CAACAAATCT TCCAAGAAAT CCATCCATGG CAGATTATGA AGCACGGATC | 840 |
| | TTTACTTTTG GGACATGGAT ATACTCAGTT AACAAGGAGC AGCTTGCAAG AGCTGGATTT | 900 |
| | TATGCTTTAG GTGAAGGTGA TAAAGTAAAG TGCTTTCCTT GTGGAGGAGG GCTAACTGAT | 960 |
| | TGGAAGCCCA GTGAAGACCC TTGGGAACAA CATGCTAAAT GGTATCCAGG GTGCAAAATAT | 1020 |
| 20 | CTGTTAGAAC AGAAGGGACA AGAATATATA AACAATATTC ATTAACTCA TTCACTTGAG | 1080 |
| | GAGTGTCTGG TAAGAACTAC TGAGAAAACA CCATCACTAA CTAGAAGAAT TGATGATACC | 1140 |
| | ATCTTCCAAA ATCCTATGGT ACAAGAAGCT ATACGAATGG GGTTCAAGTT CAAGGACATT | 1200 |
| | AAGAAAATAA TGGAGGAAAA AATTCAGATA TCTGGGAGCA ACTATAAATC ACTTGAGGTT | 1260 |
| | CTGGTTGCAG ATCTAGTGAA TGCTCAGAAA GACAGTATGC AAGATGAGTC AAGTCAGACT | 1320 |
| 25 | TCATTACAGA AAGAGATTAG TACTGAAGAG CAGCTAAGGC GCCTGCAAGA GGAGAAAGCTT | 1380 |
| | TGCAAAATCT GTATGGATAG AAATATTGCT ATCGTTTTTG TTCCTTGTGG ACATCTAGTC | 1440 |
| | ACTTGTAAC AATGTGCTGA AGCAGTTGAC AAGTGTCCCA TGTGCTACAC AGTCATTACT | 1500 |
| | TTCAAGCAAA AAATTTTAT GTCTTAATCT AACTCTATAG TAGGCATGTT ATGTTGTTCT | 1560 |
| | TATTACCCTG ATTGAATGTG TGATGTGAAC TGACTTTAAG TAATCAGGAT TGAATTCCAT | 1620 |
| 30 | TAGCATTTGC TACCAAGTAG GAAAAAAAT GTACATGGCA GTGTTTTAGT TGGCAATATA | 1680 |
| | ATCTTTGAAT TTCTTGATTT TTCAGGGTAT TAGCTGTATT ATCCATTTTT TTTACTGTTA | 1740 |
| | TTTAATTGAA ACCATAGACT AAGAATAAGA AGCATCATAC TATAACTGAA CACAATGTGT | 1800 |
| | ATTCATAGTA TACTGATTTA ATTTCTAAGT GTAAGTGAAT TAATCATCTG GATTTTTTAT | 1860 |
| | TCTTTTCAGA TAGGCTTAAC AAATGGAGCT TTCTGTATAT AAATGTGGAG ATTAGAGTTA | 1920 |
| 35 | ATCTCCCCAA TCACATAATT TGTTTTGTGT GAAAAGGAA TAAATTGTTT CATGCTGGTG | 1980 |

- 66 -

| | | | | | | | |
|----|------------|-------------|-------------|------------|------------|-------------|------|
| | GAAAGATAGA | GATTGTTTTT | AGAGGTGGT | TGTTCTGTTT | TAGGATTCTG | TCCATTTTCT | 2040 |
| | TTTAAAGTTA | TAAACACGTA | CTTGTGCGAA | TTATTTTTTT | AAAGTGATTT | GCCATTTTTG | 2100 |
| | AAAGCGTATT | TAATGATAGA | ATACTATCGA | GCCAACATGT | ACTGACATGG | AAAGATGTCA | 2160 |
| | AAGATATGTT | AAGTGTAATA | TGCAAGTGGC | AAAACACTAT | GTATAGTCTG | AGCCAGATCA | 2220 |
| 5 | AAGTATGTAT | GTTTTTAATA | TGCATAGAAC | AAAAGATTG | GAAAGATATA | CACCAAACCTG | 2280 |
| | TTAAATGTGG | TTTCTCTTCG | GGGAGGGGGG | GATTGGGGGA | GGGGCCCCAG | AGGGGTTTTA | 2340 |
| | TAGGGGCCTT | TTCACTTTCT | ACTTTTTTCA | TTTTGTCTG | TTCGAATTTT | TTATAAGTAT | 2400 |
| | GTATTACTTT | TGTAATCAGA | ATTTTITAGAA | AGTATTTTGC | TGATTTAAAG | GCTTAGGCAT | 2460 |
| | GTTCAAACGC | CTGCAAAACT | ACTTATCACT | CAGCTTTAGT | TTTTCTAATC | CAAGAAGGCA | 2520 |
| 10 | GGGCAGTTAA | CCTTTTTGGT | GCCAATGTGA | AATGTAAATG | ATTTTATGTT | TTTCCTGCTT | 2580 |
| | TGTGGATGAA | AAATATTTCT | GAGTGGTAGT | TTTTTGACAG | GTAGACCATG | TCTTATCTTG | 2640 |
| | TTTCAAAATA | AGTATTTCTG | ATTTTGTAATA | ATGAAATATA | AAATATGTCT | CAGATCTTCC | 2700 |
| | AATTAATTAG | TAAGGATTCA | TCCTTAATCC | TTGCTAGTTT | AAGCCTGCCT | AAGTCACTTT | 2760 |
| | ACTAAAAGAT | CTTTGTAAAC | TCAGTATTTT | AAACATCTGT | CAGCTTATGT | AGGTAAAAGT | 2820 |
| 15 | AGAAGCATGT | TTGTACACTG | CTTGTAGTTA | TAGTGACAGC | TTCCATGTT | GAGATTCTCA | 2880 |
| | TATCATCTTG | TATCTTAAAG | TTTCATGTGA | GTTTTTACCG | TTAGGATGAT | TAAGATGTAT | 2940 |
| | ATAGGACAAA | ATGTTAAGTC | TTTCCTCTAC | CTACATTTGT | TTTCTTGGCT | AGTAATAGTA | 3000 |
| | GTAGATACTT | CTGAAATAAA | TGTTCTCTCA | AGATCCTTAA | AACCTCTTGG | AAATTATAAA | 3060 |
| | AATATTGGCA | AGAAAAGAAG | AATAGTTGTT | TAAATATTTT | TTAAAAACA | CTGAATAAG | 3120 |
| 20 | AATCAGTAGG | GTATAAACTA | GAAGTTTAAA | AATGCCTCAT | AGAACGTCCA | GGGTTTACAT | 3180 |
| | TACAAGATTC | TCACAACAAA | CCCATTGTAG | AGGTGAGTAA | GGCATGTTAC | TACAGAGGAA | 3240 |
| | AGTTTGAGAG | TAAACTGTGA | AAAAATTATA | TTTTTGTTGT | ACTTTCTAAG | AGAAAGAGTA | 3300 |
| | TTGTTATGTT | CTCCTAACTT | CTGTTGATTA | CTACTTTAAG | TGATATTCAT | TTAAACATT | 3360 |
| | GCAAATTTAT | TTTATTTATT | TAATTTTCTT | TTTGAGATGG | AGTCTTGCTT | GTCACCCAGG | 3420 |
| 25 | CTGGAGTGCA | GTGGAGTGAT | CTCTGCTCAC | TGCAACCTCC | GCCTTCTGGG | TTCAAGCGAT | 3480 |
| | TCTCGTGCCT | CAGCTTCCTG | AGTAGCTGGA | ATTACAGGCA | GGTGCCACCA | TGCCCCACTA | 3540 |
| | ATTTTTTTTT | ATTTTITAGTA | GAGACGGGGT | TTCACCATGT | TGGCCAGGCT | GGTATCAAAC | 3600 |
| | TCCTGACCTC | AAGAGATCCA | CTCGCCTTGC | CCTCCCAAAG | TGCTGGGATT | ACAGGCTTGA | 3660 |
| | GCCACCACGC | CCGGCTAAAA | CATTGCAAAAT | TTAAATGAGA | GTTTTAAAAA | TTAAATAATG | 3720 |
| 30 | ACTGCCCTGT | TTCTGTTTTA | GTATGTAAAT | CCTCAGTTCT | TCACCTTTGC | ACTGTCTGCC | 3780 |
| | ACTTAGTTTG | GTTATATAGT | CATTAACTTG | AATTTGGTCT | GTATAGTCTA | GACTTTAAAT | 3840 |
| | TTAAAGTTTT | CTACAAGGGG | AGAAAAGTGT | TAAAATTTTT | AAAATATGTT | TTCCAGGACA | 3900 |
| | CTTCACTTCC | AAGTCAGGTA | GGTAGTTCAA | TCTAGTTGTT | AGCCAAGGAC | TCAAGGACTG | 3960 |
| | AATTGTTTTA | ACATAAGGCT | TTTCCTGTTT | TGGGAGCCGC | ACTTCATTAA | AATTCTTCTA | 4020 |
| 35 | AAACTTGTAT | GTTTAGAGTT | AAGCAAGACT | TTTTTTCTTC | CTCTCCATGA | GTTGTGAAAT | 4080 |
| | TTAATGCACA | ACGCTGATGT | GGCTAACAAG | TTTATTTTAA | GAATTGTTTA | GAAATGCTGT | 4140 |
| | TGCTTCAGGT | TCTTAAAAATC | ACTCAGCACT | CCAACCTCTA | ATCAAATTTT | TGGAGACTTA | 4200 |

- 67 -

ACAGCATTTG TCTGTGTTTG AACTATAAAA AGCACCGGAT CTTTCCATC TAATCCGCA 4260
 AAAATTGATC ATTTGCAAAG TCAAACTAT AGCCATATCC AAATCTTTTC CCCCTCCCAA 4320
 GAGTTCTCAG TGTCTACATG TAGACTATTC CTTTCTGTA TAAAGTTCAC TCTAGGATTT 4380
 CAAGTCACCA CTTATTTTAC ATTTTAGTCA TGCAAAGATT CAAGTAGTTT TGCAATAAGT 4440
 5 ACTTATCTTT ATTTGTAATA ATTTAGTCTG CTGATCAAAA GCATTGTCTT AATTTTGTAG 4500
 AACTGGTTTT AGCATTTACA AACTAAATTC CAGTTAATTA ATTAATAGCT TTATATTGCC 4560
 TTTCTGCTA CATTTGGTTT TTTCCCTGT CCCTTTGATT ACGGGCTAAG GTAGGGTAAG 4620
 AXXGGGTGTA GTGAGTGTAT ATAATGTGAT TTGGCCCTGT GTATTATGAT ATTTTGTAT 4680
 TTTTGTGTT ATATTATTA CATTCAGTA GTTGTTTTTT GTGTTTCCAT TTTAGGGGAT 4740
 10 AAAATTTGTA TTTTGAATA TGAATGGAGA CTACCGCCCC AGCATTAGTT TCACATGATA 4800
 TACCCTTAA ACCCGAATCA TTGTTTTATT TCCTGATTAC ACAGGTGTTG AATGGGGAAA 4860
 GGGGCTAGTA TATCAGTAGG ATATACTATG GGATGTATAT ATATCATTGC TGTTAGAGAA 4920
 ATGAAATAAA ATGGGGCTGG GCTCAGTGGC TCACGCCTGT AATCCCAGCA CTTGGGGAGG 4980
 CTGAGGCAGG TGGATCACGA GGTGAGGAGA TCGAGACCAT CCTGGCTAAC ACGGTGAAAC 5040
 15 CCCGTCTCTA CTAAAAACA GAAAATTAGC CGGGCGTGGT GCGGGGCGCC TGTAGTCCCA 5100
 GCTACTCGGG AGGCTGAGGC AGGAGAATGG TGTGAACCCG GGAGGCAGAG CTTGCAGTGA 5160
 GCCGAGATCT CGCCACTGCA CTCCAGCCTG GGCAACAGAG CAAGACTCTG TCTCAAAAAA 5220
 AAAAAAAAAA AG 5232

(2) INFORMATION FOR SEQ ID NO:4:

20 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 497 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Thr Phe Asn Ser Phe Glu Gly Ser Lys Thr Cys Val Pro Ala Asp
 1 5 10 15
 Ile Asn Lys Glu Glu Glu Phe Val Glu Glu Phe Asn Arg Leu Lys Thr
 20 25 30
 Phe Ala Asn Phe Pro Ser Gly Ser Pro Val Ser Ala Ser Thr Leu Ala
 35 40 45

- 68 -

Arg Ala Gly Phe Leu Tyr Thr Gly Glu Gly Asp Thr Val Arg Cys Phe
 50 55 60
 Ser Cys His Ala Ala Val Asp Arg Trp Gln Tyr Gly Asp Ser Ala Val
 65 70 75 80
 5 Gly Arg His Arg Lys Val Ser Pro Asn Cys Arg Phe Ile Asn Gly Phe
 85 90 95
 Tyr Leu Glu Asn Ser Ala Thr Gln Ser Thr Asn Ser Gly Ile Gln Asn
 100 105 110
 Gly Gln Tyr Lys Val Glu Asn Tyr Leu Gly Ser Arg Asp His Phe Ala
 10 115 120 125
 Leu Asp Arg Pro Ser Glu Thr His Ala Asp Tyr Leu Leu Arg Thr Gly
 130 135 140
 Gln Val Val Asp Ile Ser Asp Thr Ile Tyr Pro Arg Asn Pro Ala Met
 145 150 155 160
 15 Tyr Cys Glu Glu Ala Arg Leu Lys Ser Phe Gln Asn Trp Pro Asp Tyr
 165 170 175
 Ala His Leu Thr Pro Arg Glu Leu Ala Ser Ala Gly Leu Tyr Tyr Thr
 180 185 190
 Gly Ile Gly Asp Gln Val Gln Cys Phe Cys Cys Gly Gly Lys Leu Lys
 20 195 200 205
 Asn Trp Glu Pro Cys Asp Arg Ala Trp Ser Glu His Arg Arg His Phe
 210 215 220
 Pro Asn Cys Phe Phe Val Leu Gly Arg Asn Leu Asn Ile Arg Ser Glu
 225 230 235 240
 25 Ser Asp Ala Val Ser Ser Asp Arg Asn Phe Pro Asn Ser Thr Asn Leu
 245 250 255
 Pro Arg Asn Pro Ser Met Ala Asp Tyr Glu Ala Arg Ile Phe Thr Phe
 260 265 270
 Gly Thr Trp Ile Tyr Ser Val Asn Lys Glu Gln Leu Ala Arg Ala Gly
 30 275 280 285
 Phe Tyr Ala Leu Gly Glu Gly Asp Lys Val Lys Cys Phe His Cys Gly
 290 295 300
 Gly Gly Leu Thr Asp Trp Lys Pro Ser Glu Asp Pro Trp Glu Gln His
 305 310 315 320
 35 Ala Lys Trp Tyr Pro Gly Cys Lys Tyr Leu Leu Glu Gln Lys Gly Gln
 325 330 335
 Glu Tyr Ile Asn Asn Ile His Leu Thr His Ser Leu Glu Glu Cys Leu

- 69 -

340 345 350
 Val Arg Thr Thr Glu Lys Thr Pro Ser Leu Thr Arg Arg Ile Asp Asp
 355 360 365
 Thr Ile Phe Gln Asn Pro Met Val Gln Glu Ala Ile Arg Met Gly Phe
 5 370 375 380
 Ser Phe Lys Asp Ile Lys Lys Ile Met Glu Glu Lys Ile Gln Ile Ser
 385 390 395 400
 Gly Ser Asn Tyr Lys Ser Leu Glu Val Leu Val Ala Asp Leu Val Asn
 405 410 415
 10 Ala Gln Lys Asp Ser Met Gln Asp Glu Ser Ser Gln Thr Ser Leu Gln
 420 425 430
 Lys Glu Ile Ser Thr Glu Glu Gln Leu Arg Arg Leu Gln Glu Glu Lys
 435 440 445
 Leu Cys Lys Ile Cys Met Asp Arg Asn Ile Ala Ile Val Phe Val Pro
 15 450 455 460
 Cys Gly His Leu Val Thr Cys Lys Gln Cys Ala Glu Ala Val Asp Lys
 465 470 475 480
 Cys Pro Met Cys Tyr Thr Val Ile Thr Phe Lys Gln Lys Ile Phe Met
 485 490 495
 20 Ser

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

25

- (A) LENGTH: 6669 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

30 TTGCTCTGTC ACCCAGTTTG GAGTGCAGTT ATGCAGTCTC AACTGCAAG CTCTGCCTCA 60
 TGGGCTCAAG TGAACCTCCT GCCTCAGCCT CTCAAGTAGC TGGGACCACA GGCAGGTGCC 120
 ACCATGTCTG GCTAATTTTT GAGTTTCTTT GTAGAGATGG TGTTTTGCCA AGTCACCCAG 180

- 70 -

| | | | | | | | |
|----|------------|------------|------------|-------------|------------|-------------|------|
| | TTTGAGGCTG | GTCTCAAACA | CCTGGGCTCA | AGCAATCCAT | CTACCTCAGC | CTCCCAAAGT | 240 |
| | GCTGGGATTA | CAGGAGTGAG | CCATGGCATG | AGGCCTTG TG | GGGTGTCTCT | TTTAAATGAA | 300 |
| | AGCATACTCT | TTTTACGTAT | TTGATATGAA | GGAATATCCT | TCCTTTCCAC | AAAGACAAAA | 360 |
| | ATTATCTTAT | TTTTCTCAAA | ACATATGTCC | TTTTTCTCTA | CTTTTCATTT | TTGTTACTTT | 420 |
| 5 | TGATGGACAC | ATGTGTTACA | TTGATTTCAC | TTTCTCATAA | TTCTGCTGTA | AGAAAAACAA | 480 |
| | TAGTGCCAGT | TCAATGACAA | ATAGCAACAG | TCTGTTATTG | CTAGACTGTT | ACTGTTAGTG | 540 |
| | GAGACTACCA | GAACAGTCAG | TCCCAGTGTC | AGGGAATCAA | AGAGAACATG | TTCCCTCTCT | 600 |
| | AAAGGGCACA | GCTGCTGCTC | AGCTTTAGCT | GATTGCTGCC | CTGCAGGACT | ATAGGCCCCAG | 660 |
| | TGTTGCTAGA | TCTTTTGATG | TTTCAAGAGA | AGCTTGGAAT | CTAGAATGTG | ATGGGAAGTC | 720 |
| 10 | TCTTACATTT | AAACATGTTG | GCAATTAATG | GTAAGATTTA | AAAATACTGT | GGTCCAAGAA | 780 |
| | AAAAATGGAT | TTGGAAACTG | GATTAAATTC | AAATGAGGCA | TGCAGATTAA | TCTACAGCAT | 840 |
| | GGTACAATGT | GAATTTTCTG | GTTTCTTTAA | TTGCACTGTA | ATTAGGTAAG | ATGTTAGCTT | 900 |
| | TGGGGAAGCT | AAGTCAGAG | TATGCAGAAA | CTATTATTTT | TGTAAGTTTT | CTCTAAGTAT | 960 |
| | AAATAAATTT | CAAAATAAAA | ATAAAAACTT | AGTAAAGAAC | TATAATGCAA | TTCTATGTAA | 1020 |
| 15 | GCCAAACATA | ATATGTCTTC | CAGTTTGAAA | CCTCTGGGTT | TTATTTTATT | TTATTTTATT | 1080 |
| | TTTGAGACAG | AGTCTTGCTG | TGTCACCCAG | GCTGGAGTGT | AGTGGCACTA | TTTCGGCCCA | 1140 |
| | CTGCAACCTC | CACCTCCCAG | GCTCAAATGA | TTCTCCTGCC | TCAGCCTCCG | GAGTAGCTGG | 1200 |
| | GATTACAGGC | GCGTACCACC | ACACCCAGCT | AATTTTGTGA | TTTTTAGTAG | AGATGGGGTT | 1260 |
| | TCACCATTTT | GGCCAGGCTG | GTTTTGAACT | CCTGACCTCA | AGTGATCCAC | TTGTCTTG GC | 1320 |
| 20 | CTCCCAAAT | GCTGGGATTA | CAGGCGTGAG | CCACTGCACC | AGGCAGAGGC | CTCTGTTTTT | 1380 |
| | TATCTCTTTT | TGGCCTCTAC | AGTGCCTAGT | AAAGCACCTG | ATACATGGTA | AACGATCAGT | 1440 |
| | AATTACTAGT | ACTCTATTTT | GGAGAAAATG | ATTTTTTAAA | AAGTCATTGT | GTTCCATCCA | 1500 |
| | TGAGTCGTTT | GAGTTTAAA | ACTGTCTTTT | TGTTTGTTTT | TGAACAGGTT | TACAAAGGAG | 1560 |
| | GAAAACGACT | TCTTCTAGAT | TTTTTTTCA | GTTTCTCTA | TAAATCAAAA | CATCTCAAAA | 1620 |
| 25 | TGGAGACCTA | AAATCCTTAA | AGGGACTTAG | TCTAATCTCG | GGAGGTAGTT | TTGTGCATGG | 1680 |
| | GTAAACAAAT | TAAGTATTAA | CTGGTGTTTT | ACTATCCAAA | GAATGCTAAT | TTTATAAACA | 1740 |
| | TGATCGAGTT | ATATAAGGTA | TACCATAATG | AGTTTGATTT | TGAATTTGAT | TTGTGGAAAT | 1800 |
| | AAAGGAAAAG | TGATTCTAGC | TGGGGCATAT | TGTTAAAGCA | TTTTTTTCAG | AGTTGGCCAG | 1860 |
| | GCAGTCTCCT | ACTGGCACAT | TCTCCCATT | TGTAGAATAG | AAATAGTACC | TGTGTTTGGG | 1920 |
| 30 | AAAGATTTTA | AAATGAGTGA | CAGTTATTTG | GAACAAAGAG | CTAATAATCA | ATCCACTGCA | 1980 |
| | AATTAAAGAA | ACATGCAGAT | GAAAGTTTTG | ACACATTAAA | ATACTTCTAC | AGTGACAAAG | 2040 |
| | AAAAATCAAG | AACAAAGCTT | TTTGATATGT | GCAACAAATT | TAGAGGAAGT | AAAAAGATAA | 2100 |
| | ATGTGATGAT | TGGTCAAGAA | ATTATCCAGT | TATTTACAAG | GCCACTGATA | TTTTAAACGT | 2160 |
| | CCAAAAGTTT | GTTTAAATGG | GCTGTTACCG | CTGAGAATGA | TGAGGATGAG | AATGATGGTT | 2220 |
| 35 | GAAGGTTACA | TTTTAGGAAA | TGAAGAACT | TAGAAAATTA | ATATAAAGAC | AGTGATGAAT | 2280 |
| | ACAAAGAAGA | TTTTTATAAC | AATGTGTAAA | ATTTTTGGCC | AGGGAAAGGA | ATATTGAAGT | 2340 |
| | TAGATACAAT | TACTTACCTT | TGAGGGAAAT | AATTGTTGGT | AATGAGATGT | GATGTTTCTC | 2400 |

- 71 -

| | | | | | | | |
|----|------------|-------------|------------|------------|------------|------------|------|
| | CTGCCACCTG | GAAACAAAGC | ATTGAAGTCT | GCAGTTGAAA | AGCCCAACGT | CTGTGAGATC | 2460 |
| | CAGGAAACCA | TGCTTGCAAA | CCACTGGTAA | AAAAAAAAAA | AAAAAAAAAA | AAAGCCACAG | 2520 |
| | TGACTTGCTT | ATTGGTCATT | GCTAGTATTA | TCGACTCAGA | ACCTCTTTAC | TAATGGCTAG | 2580 |
| | TAAATCATAA | TTGAGAAATT | CTGAATTTTG | ACAAGGTCTC | TGCTGTTGAA | ATGGTAAATT | 2640 |
| 5 | TATTATTTTT | TTTGTTCATGA | TAAATCTCTG | TTCAAGGTAT | GCTATCCATG | AAATAATTTC | 2700 |
| | TGACCAAAAC | TAAATTGATG | CAATTTGATT | ATCCATCTTA | GCCTACAGAT | GGCATCTGGT | 2760 |
| | AACTTTTGAC | TGTTTTAAAA | AATAAATCCA | CTATCAGAGT | AGATTTGATG | TTGGCTTCAG | 2820 |
| | AAACATTTAG | AAAAACAAAA | GTTCAAAAAT | GTTTTCAGGA | GGTGATAAGT | TGAATAACTC | 2880 |
| | TACAATGTTA | GTTCTTTGAG | GGGGACAAAA | AATTTAAAAT | CTTTGAAAGG | TCTTATTTTA | 2940 |
| 10 | CAGCCATATC | TAAATTATCT | TAAGAAAATT | TTTAACAAAG | GGAATGAAAT | ATATATCATG | 3000 |
| | ATTCTGTTTT | TCCAAAAGTA | ACCTGAATAT | AGCAATGAAG | TTCAGTTTTG | TTATTGGTAG | 3060 |
| | TTTGGGCAGA | GTCTCTTTTT | GCAGCACCTG | TTGTCTACCA | TAATTACAGA | GGACATTTC | 3120 |
| | ATGTTCTAGC | CAAGTATACT | ATTAGAATAA | AAAACTTAA | CATTGAGTTG | CTTCAACAGC | 3180 |
| | ATGAACTGA | GTCCAAAAGA | CCAAATGAAC | AAACACATTA | ATCTCTGATT | ATTTATTTTA | 3240 |
| 15 | AATAGAATAT | TTAATTGTGT | AAGATCTAAT | AGTATCATT | TACTTAAGCA | ATCATATTCC | 3300 |
| | TGATGATCTA | TGGGAAATAA | CTATTATTTA | ATTAATATTG | AAACCAGGTT | TTAAGATGTG | 3360 |
| | TTAGCCAGTC | CTGTTACTAG | TAAATCTCTT | TATTTGGAGA | GAAATTTTAG | ATTGTTTTGT | 3420 |
| | TCTCCTTATT | AGAAGGATTG | TAGAAAGAAA | AAAATGACTA | ATTGGAGAAA | AATTGGGGAT | 3480 |
| | ATATCATATT | TCCTGAATT | CAAAATGTCT | TCAGTTGTAA | ATCTTACCAT | TATTTTACGT | 3540 |
| 20 | ACCTCTAAGA | AATAAAAGTG | CTTCTAATTA | AAATATGATG | TCATTAATTA | TGAAATACTT | 3600 |
| | CTTGATAACA | GAAGTTTTTA | AATAGCCATC | TTAGAATCAG | TGAAATATGG | TAATGTATTA | 3660 |
| | TTTTCTCTCT | TTGAGTNAGG | TCTTGTGCTT | TTTNTCCTG | GCCACTAAAT | NTCACCATNT | 3720 |
| | CCAANAAGCA | AANTAAACCT | ATTCTGAATA | TTTTTGCTGT | GAAACACTTG | NCAGCAGAGC | 3780 |
| | TTCCCNCCA | TGNAGAAGC | TTCATGAGTC | ACACATTACA | TCTTTGGGTT | GATTGAATGC | 3840 |
| 25 | CACTGAAACA | TTTCTAGTAG | CCTGGAGNAG | TTGACCTACC | TGTGGAGATG | CCTGCCATTA | 3900 |
| | AATGGCATCC | TGATGGCTTA | ATACACATCA | CTCTTCTGTG | NAGGSTTTTA | ATTTTCAACA | 3960 |
| | CAGCTTACTC | TGTAGCATCA | TGTTTACATT | GTATGTATAA | AGATTATACN | AAGGTGCAAT | 4020 |
| | TGTGTATTTT | TTCTTAAAA | TGTATCAGTA | TAGGATTTAG | AATCTCCATG | TTGAAACTCT | 4080 |
| | AAATGCATAG | AAATAAAAAT | AATAAAAAAT | TTTTCATTTT | GGCTTTTCAG | CCTAGTATTA | 4140 |
| 30 | AAACTGATAA | AAGCAAAGCC | ATGCACAAAA | CTACCTCCCT | AGAGAAAGGC | TAGTCCCTTT | 4200 |
| | TCTTCCCAT | TCATTTTATT | ATGAACATAG | TAGAAAACAG | CATATTCTTA | TCAAATTTGA | 4260 |
| | TGAAAAGCGC | CAACACGTTT | GAAGTGAAT | ACGACTTGTC | ATGTGAACTG | TACCGAATGT | 4320 |
| | CTACGTATTC | CACTTTTCCT | GCTGGGGTTC | CTGTCTCAGA | AAGGAGTCTT | GCTCGTGCTG | 4380 |
| | GTTTCTATTA | CACTGGTGTG | AATGACAAGG | TCAAATGCTT | CTGTTGTGGC | CTGATGCTGG | 4440 |
| 35 | ATAACTGGAA | AAGAGGAGAC | AGTCCTACTG | AAAAGCATAA | AAAGTTGTAT | CCTAGCTGCA | 4500 |
| | GATTGCTTCA | GAGTCTAAAT | TCCGTTAACA | ACTTGGAAGC | TACCTCTCAG | CCTACTTTTC | 4560 |
| | CTTCTTCAGT | AACACATTCC | ACACACTCAT | TACTTCCGGG | TACAGAAAAC | AGTGGATATT | 4620 |

- 72 -

| | | | | | | | |
|----|------------|------------|------------|------------|-------------|------------|------|
| | TCCGTGGCTC | TTATTCAAAC | TCTCCATCAA | ATCCTGTAAA | CTCCAGAGCA | AATCAAGAAT | 4680 |
| | TTTCTGCCTT | GATGAGAAGT | TCCTACCCCT | GTCCAATGAA | TAACGAAAAT | GCCAGATTAC | 4740 |
| | TTACTTTTCA | GACATGGCCA | TTGACTTTTC | TGTCGCCAAC | AGATCTGGCA | CGAGCAGGCT | 4800 |
| | TTTACTACAT | AGGACCTGGA | GACAGAGTGG | CTTGCTTTGC | CTGTGGTGGA | AAATTGAGCA | 4860 |
| 5 | ATTGGGAACC | GAAGGATAAT | GCTATGTCAG | AACACCTGAG | ACATTTTCCC | AAATGCCCAT | 4920 |
| | TTATAGAAAA | TCAGCTTCAA | GACACTTCAA | GATACACAGT | TTCTAATCTG | AGCATGCAGA | 4980 |
| | CACATGCAGC | CCGCTTTAAA | ACATTCTTTA | ACTGGCCCTC | TAGTGTCTTA | GTTAATCCTG | 5040 |
| | AGCAGCTTGC | AAGTGCGGGT | TTTTATTATG | TGGGTAACAG | TGATGATGTC | AAATGCTTTT | 5100 |
| | GCTGTGATGG | TGGACTCAGG | TGTTGGGAAT | CTGGAGATGA | TCCATGGGTT | CAACATGCCA | 5160 |
| 10 | AGTGGTTTCC | AAGGTGTGAG | TACTTGATAA | GAATTAAAGG | ACAGGAGTTC | ATCCGTCAAG | 5220 |
| | TTCAAGCCAG | TTACCCTCAT | CTACTTGAAC | AGCTGCTATC | CACATCAGAC | AGCCCAGGAG | 5280 |
| | ATGAAAATGC | AGAGTCATCA | ATTATCCATT | TTGAACCTGG | AGAAGACCAT | TCAGAAGATG | 5340 |
| | CAATCATGAT | GAATACTCCT | GTGATTAATG | CTGCCGTGGA | AATGGGCTTT | AGTAGAAGCC | 5400 |
| | TGGTAAAACA | GACAGTTCAG | AGAAAAATCC | TAGCAACTGG | AGAGAATTAT | AGACTAGTCA | 5460 |
| 15 | ATGATCTTGT | GTTAGACTTA | CTCAATGCAG | AAGATGAAAT | AAGGGAAGAG | GAGAGAGAAA | 5520 |
| | GAGCAACTGA | GGAAAAAGAA | TCAAATGATT | TATTATTAAT | CCGGAAGAAT | AGAATGGCAC | 5580 |
| | TTTTTCAACA | TTTGACTTGT | GTAATTCCAA | TCCTGGATAG | TCTACTAACT | GCCGGAATTA | 5640 |
| | TTAATGAACA | AGAACATGAT | GTTATTAAAC | AGAAGACACA | GACGTCTTTA | CAAGCAAGAG | 5700 |
| | AACTGATTGA | TACGATTTTA | GTAAAAGGAA | ATATTGCAGC | CACTGTATTTC | AGAAACTCTC | 5760 |
| 20 | TGCAAGAAGC | TGAAGCTGTG | TTATATGAGC | ATTTATTTGT | GCAACAGGAC | ATAAAATATA | 5820 |
| | TTCCACAGAG | AGATGTTTCA | GATCTACCAG | TGGAAGAACA | ATTGCGGAGA | CTACAAGAAG | 5880 |
| | AAAGAACATG | TAAAGTGTGT | ATGGACAAAG | AAGTGTCAT | AGTGTTTATT | CCTGTGGTTC | 5940 |
| | ATCTAGTAGT | ATGCAAAGAT | TGTGCTCCTT | CTTTAAGAAA | GTGTCCTATT | TGTAGGAGTA | 6000 |
| | CAATCAAGGG | TACAGTTCGT | ACATTTCTTT | CATGAAGAAG | AACCAAAACA | TCGTCTAAAC | 6060 |
| 25 | TTTAGAATTA | ATTTATTAAA | TGTATTATAA | CTTTAACTTT | TATCCTAATT | TGTTTTCCTT | 6120 |
| | AAAATTTTTA | TTTATTTACA | ACTCAAAAAA | CATTGTTTTG | TGTAACATAT | TTATATATGT | 6180 |
| | ATCTAAACCA | TATGAACATA | TATTTTITAG | AACTAAGAG | AATGATAGGC | TTTGTTCTTT | 6240 |
| | ATGAACGAAA | AAGAGGTAGC | ACTACAAACA | CAATATTCAA | TCAAAATTTC | AGCATTATTG | 6300 |
| | AAATTGTAAG | TGAAGTAAAA | CTTAAGATAT | TTGAGTTAAC | CTTTAAGAAT | TTTAAATATT | 6360 |
| 30 | TTGGCATTGT | ACTAATACCG | GGAACATGAA | GCCAGGTGTG | GTGGTATGTG | CCTGTAGTCC | 6420 |
| | CAGGCTGAGG | CAAGAGAATT | ACTTGAGCCC | AGGAGTTTGA | ATCCATCCTG | GGCAGCATAC | 6480 |
| | TGAGACCCTG | CCTTTAAAAA | CAAACAGAAC | AAAAACAAAA | CACCAGGGAC | ACATTTCTCT | 6540 |
| | GTCTTTTTTG | ATCAGTGTCC | TATACATCGA | AGGTGTGCAT | ATATGTTGAA | TCACATTTTA | 6600 |
| | GGGACATGGT | GTTTTTATAA | AGAATTCTGT | GAGAAAAAAT | TTAATAAAGC | AACCAAAAAA | 6660 |
| 35 | AAAAAATAA | | | | | | 6669 |

(2) INFORMATION FOR SEQ ID NO:6:

- 73 -

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 604 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

5 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Asn Ile Val Glu Asn Ser Ile Phe Leu Ser Asn Leu Met Lys Ser
  1             5             10             15
10 Ala Asn Thr Phe Glu Leu Lys Tyr Asp Leu Ser Cys Glu Leu Tyr Arg
      20             25             30
Met Ser Thr Tyr Ser Thr Phe Pro Ala Gly Val Pro Val Ser Glu Arg
      35             40             45
Ser Leu Ala Arg Ala Gly Phe Tyr Tyr Thr Gly Val Asn Asp Lys Val
15      50             55             60
Lys Cys Phe Cys Cys Gly Leu Met Leu Asp Asn Trp Lys Arg Gly Asp
65             70             75             80
Ser Pro Thr Glu Lys His Lys Lys Leu Tyr Pro Ser Cys Arg Phe Val
      85             90             95
20 Gln Ser Leu Asn Ser Val Asn Asn Leu Glu Ala Thr Ser Gln Pro Thr
      100            105            110
Phe Pro Ser Ser Val Thr His Ser Thr His Ser Leu Leu Pro Gly Thr
      115            120            125
Glu Asn Ser Gly Tyr Phe Arg Gly Ser Tyr Ser Asn Ser Pro Ser Asn
25      130            135            140
Pro Val Asn Ser Arg Ala Asn Gln Glu Phe Ser Ala Leu Met Arg Ser
145            150            155            160
Ser Tyr Pro Cys Pro Met Asn Asn Glu Asn Ala Arg Leu Leu Thr Phe
      165            170            175
30 Gln Thr Trp Pro Leu Thr Phe Leu Ser Pro Thr Asp Leu Ala Arg Ala
      180            185            190
Gly Phe Tyr Tyr Ile Gly Pro Gly Asp Arg Val Ala Cys Phe Ala Cys
      195            200            205
Gly Gly Lys Leu Ser Asn Trp Glu Pro Lys Asp Asn Ala Met Ser Glu

```

- 74 -

| | | | | | | |
|----|---|-----|-----|-----|-----|-----|
| | 210 | | 215 | | 220 | |
| | His Leu Arg His Phe Pro Lys Cys Pro Phe Ile Glu Asn Gln Leu Gln | | | | | |
| | 225 | | 230 | | 235 | 240 |
| | Asp Thr Ser Arg Tyr Thr Val Ser Asn Leu Ser Met Gln Thr His Ala | | | | | |
| 5 | | 245 | | 250 | | 255 |
| | Ala Arg Phe Lys Thr Phe Phe Asn Trp Pro Ser Ser Val Leu Val Asn | | | | | |
| | | 260 | | 265 | | 270 |
| | Pro Glu Gln Leu Ala Ser Ala Gly Phe Tyr Tyr Val Gly Asn Ser Asp | | | | | |
| | | 275 | | 280 | | 285 |
| 10 | Asp Val Lys Cys Phe Cys Cys Asp Gly Gly Leu Arg Cys Trp Glu Ser | | | | | |
| | | 290 | | 295 | | 300 |
| | Gly Asp Asp Pro Trp Val Gln His Ala Lys Trp Phe Pro Arg Cys Glu | | | | | |
| | | 305 | | 310 | | 315 |
| | Tyr Leu Ile Arg Ile Lys Gly Gln Glu Phe Ile Arg Gln Val Gln Ala | | | | | |
| 15 | | 325 | | 330 | | 335 |
| | Ser Tyr Pro His Leu Leu Glu Gln Leu Leu Ser Thr Ser Asp Ser Pro | | | | | |
| | | 340 | | 345 | | 350 |
| | Gly Asp Glu Asn Ala Glu Ser Ser Ile Ile His Leu Glu Pro Gly Glu | | | | | |
| | | 355 | | 360 | | 365 |
| 20 | Asp His Ser Glu Asp Ala Ile Met Met Asn Thr Pro Val Ile Asn Ala | | | | | |
| | | 370 | | 375 | | 380 |
| | Ala Val Glu Met Gly Phe Ser Arg Ser Leu Val Lys Gln Thr Val Gln | | | | | |
| | | 385 | | 390 | | 395 |
| | Arg Lys Ile Leu Ala Thr Gly Glu Asn Tyr Arg Leu Val Asn Asp Leu | | | | | |
| 25 | | 405 | | 410 | | 415 |
| | Val Leu Asp Leu Leu Asn Ala Glu Asp Glu Ile Arg Glu Glu Glu Arg | | | | | |
| | | 420 | | 425 | | 430 |
| | Glu Arg Ala Thr Glu Glu Lys Glu Ser Asn Asp Leu Leu Leu Ile Arg | | | | | |
| | | 435 | | 440 | | 445 |
| 30 | Lys Asn Arg Met Ala Leu Phe Gln His Leu Thr Cys Val Ile Pro Ile | | | | | |
| | | 450 | | 455 | | 460 |
| | Leu Asp Ser Leu Leu Thr Ala Gly Ile Ile Asn Glu Gln Glu His Asp | | | | | |
| | | 465 | | 470 | | 475 |
| | Val Ile Lys Gln Lys Thr Gln Thr Ser Leu Gln Ala Arg Glu Leu Ile | | | | | |
| 35 | | 485 | | 490 | | 495 |
| | Asp Thr Ile Leu Val Lys Gly Asn Ile Ala Ala Thr Val Phe Arg Asn | | | | | |
| | | 500 | | 505 | | 510 |

- 75 -

Ser Leu Gln Glu Ala Glu Ala Val Leu Tyr Glu His Leu Phe Val Gln
515 520 525
Gln Asp Ile Lys Tyr Ile Pro Thr Glu Asp Val Ser Asp Leu Pro Val
530 535 540
5 Glu Glu Gln Leu Arg Arg Leu Pro Glu Glu Arg Thr Cys Lys Val Cys
545 550 555 560
Met Asp Lys Glu Val Ser Ile Val Phe Ile Pro Cys Gly His Leu Val
565 570 575
Val Cys Lys Asp Cys Ala Pro Ser Leu Arg Lys Cys Pro Ile Cys Arg
10 580 585 590
Ser Thr Ile Lys Gly Thr Val Arg Thr Phe Leu Ser
595 600

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- 15 (A) LENGTH: 3732 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

20 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

GAGCGCCCGG GCTGATCCGA GCCGAGCGGG CCGTATCTCC TTGTCGGCGC CGCTGATTC 60
CGGCTCTGCG GAGGCCTCTA GGCAGCCGCG CAGCTTCCGT GTTGCTGCG CCCGCACTGC 120
GATTTACAAC CCTGAAGAAT CTCCCTATCC CTATTTTGTC CCCCTGCAGT AATAAATCCC 180
ATTATGGAGA TCTCGAAACT TTATAAAGGG ATATAGTTTG AATTCTATGG AGTGTAATTT 240
25 TGTGTATGAA TTATATTTTT AAAACATTGA AGAGTTTCA GAAAGAAGGC TAGTAGAGTT 300
GATTACTGAT ACTTTATGCT AAGCAGTACT TTTTGGTAG TACAATATTT TGTTAGGCGT 360
TTCTGATAAC ACTAGAAAGG ACAAGTTTTA TCTTGTGATA AATTGATTAA TGTTTACAAC 420
ATGACTGATA ATTATAGCTG AATAGTCCTT AAATGATGAA CAGGTTATTT AGTTTTTAAA 480
TGCACTGTAA AAAGTGTGCT GTGGAAATTT TATGGCTAAC TAAGTTTATG GAGAAAATAC 540
30 CTTCACTTGA TCAAGAATAA TAGTGGTATA CAAAGTTAGG AAGAAAGTCA ACATGATGCT 600
GCAGGAAATG GAAACAAATA CAAATGATAT TTAACAAAGA TAGAGTTTAC AGTTTTTGAA 660
CTTTAAGCCA AATTCATTTG ACATCAAGCA CTATAGCAGG CACAGGTTCA ACAAAGCTTG 720

- 76 -

| | | | | | | | |
|----|------------|------------|------------|------------|------------|------------|------|
| | TGGGTATTGA | CTTCCCCCAA | AAGTTGTCAG | CTGAAGTAAT | TTAGCCCACT | TAAGTAAATA | 780 |
| | CTATGATGAT | AAGCTGTGTG | AACCTAGCTT | TTAAATAGTG | TGACCATATG | AAGGTTTTAA | 840 |
| | TTACTTTTGT | TTATTGGAAT | AAAATGAGAT | TTTTTGGGTT | GTCATGTTAA | AGTGCTTATA | 900 |
| | GGGAAAGAAG | CCTGCATATA | ATTTTTTACC | TTGTGGCATA | ATCAGTAATT | GGTCTGTTAT | 960 |
| 5 | TCAGGCTTCA | TAGCTTGTA | CCAATATAA | ATAAAAGGCA | TAATTTAGGT | ATTCTATAGT | 1020 |
| | TGCTTAGAAT | TTTGTTAATA | TAAATCTCTG | TGAAAAATCA | AGGAGTTTTA | ATATTTTCAG | 1080 |
| | AAGTGCATCC | ACCTTTCAGG | GCTTTAAGTT | AGTATTAAC | CAAGATTATG | AACAAATAGC | 1140 |
| | ACTTAGGTTA | CCTGAAAGAG | TTACTACAAC | CCCAAAGAGT | TGTGTTCTAA | GATGATCTT | 1200 |
| | GGTAATTCAG | AGAGATACTC | ATCCTACCTG | AATATAAACT | GAGATAAATC | CAGTAAAGAA | 1260 |
| 10 | AGTGTAGTAA | ATTCTACATA | AGAGTCTATC | ATTGATTCTT | TTTTGTGGTA | AAAATCTTAG | 1320 |
| | TTCAATGTGA | GAAATTTTCA | GTGAATGTTT | TAGCTATCAA | ACAGTACTGT | CACCTACTCA | 1380 |
| | TGCACAAAAC | TGCCTCCCAA | AGACTTTTCC | CAGGTCCCTC | GTATCAAAAC | ATTAAGAGTA | 1440 |
| | TAATGGAAGA | TAGCACGATC | TTGTCAGATT | GGACAAACAG | CAACAAACAA | AAAATGAAGT | 1500 |
| | ATGACTTTTC | CTGTGAACTC | TACAGAATGT | CTACATATTC | AACTTTCCCC | GCCGGGGTGC | 1560 |
| 15 | CTGTCTCAGA | AAGGAGTCTT | GCTCGTGCTG | GTTTTTATTA | TACTGGTGTG | AATGACAAGG | 1620 |
| | TCAAATGCTT | CTGTTGTGGC | CTGATGCTGG | ATAACTGGAA | ACTAGGAGAC | AGTCCTATTC | 1680 |
| | AAAAGCATAA | ACAGCTATAT | CCTAGCTGTA | GCTTTATTCA | GAATCTGGTT | TCAGCTAGTC | 1740 |
| | TGGGATCCAC | CTCTAAGAAT | ACGTCTCCAA | TGAGAAACAG | TTTTGCACAT | TCATTATCTC | 1800 |
| | CCACCTTGGA | ACATAGTAGC | TTGTTCAAGT | GTTCTTACTC | CAGCCTTTCT | CCAAACCTTC | 1860 |
| 20 | TTAATTCTAG | AGCAGTTGAA | GACATCTCTT | CATCGAGGAC | TAACCCCTAC | AGTTATGCAA | 1920 |
| | TGAGTACTGA | AGAAGCCAGA | TTTCTTACCT | ACCATATGTG | GCCATTAAC | TTTTTGTAC | 1980 |
| | CATCAGAATT | GGCAAGAGCT | GGTTTTTATT | ATATAGGACC | TGGAGATAGG | GTAGCCTGCT | 2040 |
| | TTGCCTGTGG | TGGGAAGCTC | AGTAACTGGG | AACCAAAGGA | TGATGCTATG | TCAGAACACC | 2100 |
| | GGAGGCATTT | TCCCAACTGT | CCATTTTGGG | AAAATTCTCT | AGAACTCTG | AGGTTTAGCA | 2160 |
| 25 | TTTCAAATCT | GAGCATGCAG | ACACATGCAG | CTCGAATGAG | AACATTTATG | TACTGGCCAT | 2220 |
| | CTAGTGTTCC | AGTTCAGCCT | GAGCAGCTTG | CAAGTGCTGG | TTTTTATTAT | GTGGGTCGCA | 2280 |
| | ATGATGATGT | CAAATGCTTT | TGTGTGATG | GTGGCTTGAG | GTGTGGGAA | TCTGGAGATG | 2340 |
| | ATCCATGGGT | AGAACATGCC | AAGTGGTTTC | CAAGGTGTGA | GTTCTTGATA | CGAATGAAAG | 2400 |
| | GCCAAGAGTT | TGTTGATGAG | ATTCAAGGTA | GATATCCTCA | TCTTCTTGAA | CAGCTGTTGT | 2460 |
| 30 | CAACTTCAGA | TACCACTGGA | GAAGAAAATG | CTGACCCACC | AATTATTCAT | TTTGGACCTG | 2520 |
| | GAGAAAGTTC | TTCAGAAGAT | GCTGTCATGA | TGAATACACC | TGTGGTTAAA | TCTGCCTTGG | 2580 |
| | AAATGGGCTT | TAATAGAGAC | CTGGTGAAAC | AAACAGTTCA | AAGTAAAATC | CTGACAACTG | 2640 |
| | GAGAGAACTA | TAAAACAGTT | AATGATATTG | TGTCAGCACT | TCTTAATGCT | GAAGATGAAA | 2700 |
| | AAAGAGAAGA | GGAGAAGGAA | AAACAAGCTG | AAGAAATGGC | ATCAGATGAT | TTGTCATTAA | 2760 |
| 35 | TTCGGAAGAA | CAGAATGGCT | CTCTTTCAAC | AATTGACATG | TGTGCTTCCT | ATCCTGGATA | 2820 |
| | ATCTTTTAAA | GGCCAATGTA | ATTAATAAAC | AGGAACATGA | TATTATTTAA | CAAAAAACAC | 2880 |
| | AGATACCTTT | ACAAGCGAGA | GAAGTATTG | ATACCATTTT | GGTTAAAGGA | AATGCTGCGG | 2940 |

- 77 -

CCAACATCTT CAAAAACTGT CTAAAAGAAA TTGACTCTAC ATTGTATAAG AACTTATTTG 3000
 TGGATAAGAA TATGAAGTAT ATCCCAACAG AAGATGTTTC AGGTCTGTCA CTGGAAGAAC 3060
 AATTGAGGAG GTTGCAAGAA GAACGAACCT GTAAAGTGTG TATGGACAAA GAAGTTTCTG 3120
 TTGTATTTAT TCCTTGTTGGT CATCTGGTAG TATGCCAGGA ATGTGCCCTT TCTCTAAGAA 3180
 5 AATGCCCTAT TTGCAGGGGT ATAATCAAGG GTACTGTTTCG TACATTTCTC TCTTAAAGAA 3240
 AAATAGTCTA TATTTTAACC TGCATAAAAA GGTCTTTAAA ATATTGTTGA ACACTTGAAG 3300
 CCATCTAAAG TAAAAAGGGA ATTATGAGTT TTTCAATTAG TAACATTCAT GTTCTAGTCT 3360
 GCTTTGGTAC TAATAATCTT GTTCTGAAA AGATGGTATC ATATATTTAA TCTTAATCTG 3420
 TTTATTTACA AGGGAAGATT TATGTTTGGT GAACTATATT AGTATGTATG TGTACCTAAG 3480
 10 GGAGTAGTGT CACTGCTTGT TATGCATCAT TTCAGGAGTT ACTGGATTG TTGTTCTTTC 3540
 AGAAAGCTTT GAATACTAAA TTATAGTGTA GAAAAGAACT GGAAACCAGG AACTCTGGAG 3600
 TTCATCAGAG TTATGGTGCC GAATTGTCTT TGGTGCTTTT CACTTGTGTT TAAAAATAAG 3660
 GATTTTCTC TTATTTCTCC CCCTAGTTTG TGAGAAACAT CTCAATAAAG TGCTTTAAAA 3720
 AGAAAAAAA AA 3732

15 (2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 618 amino acids

(B) TYPE: amino acid

(C) STRANDEDNESS: single

20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met His Lys Thr Ala Ser Gln Arg Leu Phe Pro Gly Pro Ser Tyr Gln
 1 5 10 15
 25 Asn Ile Lys Ser Ile Met Glu Asp Ser Thr Ile Leu Ser Asp Trp Thr
 20 25 30
 Asn Ser Asn Lys Gln Lys Met Lys Tyr Asp Phe Ser Cys Glu Leu Tyr
 35 40 45
 Arg Met Ser Thr Tyr Ser Thr Phe Pro Ala Gly Val Pro Val Ser Glu
 30 50 55 60
 Arg Ser Leu Ala Arg Ala Gly Phe Tyr Tyr Thr Gly Val Asn Asp Lys
 65 70 75 80

- 78 -

Val Lys Cys Phe Cys Cys Gly Leu Met Leu Asp Asn Trp Lys Leu Gly
 85 90 95
 Asp Ser Pro Ile Gln Lys His Lys Gln Leu Tyr Pro Ser Cys Ser Phe
 100 105 110
 5 Ile Gln Asn Leu Val Ser Ala Ser Leu Gly Ser Thr Ser Lys Asn Thr
 115 120 125
 Ser Pro Met Arg Asn Ser Phe Ala His Ser Leu Ser Pro Thr Leu Glu
 130 135 140
 His Ser Ser Leu Phe Ser Gly Ser Tyr Ser Ser Leu Pro Pro Asn Pro
 10 145 150 155 160
 Leu Asn Ser Arg Ala Val Glu Asp Ile Ser Ser Ser Arg Thr Asn Pro
 165 170 175
 Tyr Ser Tyr Ala Met Ser Thr Glu Glu Ala Arg Phe Leu Thr Tyr His
 180 185 190
 15 Met Trp Pro Leu Thr Phe Leu Ser Pro Ser Glu Leu Ala Arg Ala Gly
 195 200 205
 Phe Tyr Tyr Ile Gly Pro Gly Asp Arg Val Ala Cys Phe Ala Cys Gly
 210 215 220
 Gly Lys Leu Ser Asn Trp Glu Pro Lys Asp Asp Ala Met Ser Glu His
 20 225 230 235 240
 Arg Arg His Phe Pro Asn Cys Pro Phe Leu Glu Asn Ser Leu Glu Thr
 245 250 255
 Leu Arg Phe Ser Ile Ser Asn Leu Ser Met Gln Thr His Ala Ala Arg
 260 265 270
 25 Met Arg Thr Phe Met Tyr Trp Pro Ser Ser Val Pro Val Gln Pro Glu
 275 280 285
 Gln Leu Ala Ser Ala Gly Phe Tyr Tyr Val Gly Arg Asn Asp Asp Val
 290 295 300
 Lys Cys Phe Gly Cys Asp Gly Gly Leu Arg Cys Trp Glu Ser Gly Asp
 30 305 310 315 320
 Asp Pro Trp Val Glu His Ala Lys Trp Phe Pro Arg Cys Glu Phe Leu
 325 330 335
 Ile Arg Met Lys Gly Gln Glu Phe Val Asp Glu Ile Gln Gly Arg Tyr
 340 345 350
 35 Pro His Leu Leu Glu Gln Leu Leu Ser Thr Ser Asp Thr Thr Gly Glu
 355 360 365
 Glu Asn Ala Asp Pro Pro Ile Ile His Phe Gly Pro Gly Glu Ser Ser

- 79 -

| | | | | | | |
|----|---|-----|-----|-----|-----|-----|
| | 370 | | 375 | | 380 | |
| | Ser Glu Asp Ala Val Met Met Asn Thr Pro Val Val Lys Ser Ala Leu | | | | | |
| | 385 | | 390 | | 395 | 400 |
| | Glu Met Gly Phe Asn Arg Asp Leu Val Lys Gln Thr Val Leu Ser Lys | | | | | |
| 5 | | 405 | | 410 | | 415 |
| | Ile Leu Thr Thr Gly Glu Asn Tyr Lys Thr Val Asn Asp Ile Val Ser | | | | | |
| | | 420 | | 425 | | 430 |
| | Ala Leu Leu Asn Ala Glu Asp Glu Lys Arg Glu Glu Glu Lys Glu Lys | | | | | |
| | | 435 | | 440 | | 445 |
| 10 | Gln Ala Glu Glu Met Ala Ser Asp Asp Leu Ser Leu Ile Arg Lys Asn | | | | | |
| | | 450 | | 455 | | 460 |
| | Arg Met Ala Leu Phe Gln Gln Leu Thr Cys Val Leu Pro Ile Leu Asp | | | | | |
| | | 465 | | 470 | | 475 |
| | Asn Leu Leu Lys Ala Asn Val Ile Asn Lys Gln Glu His Asp Ile Ile | | | | | |
| 15 | | 485 | | 490 | | 495 |
| | Lys Gln Lys Thr Gln Ile Pro Leu Gln Ala Arg Glu Leu Ile Asp Thr | | | | | |
| | | 500 | | 505 | | 510 |
| | Ile Trp Val Lys Gly Asn Ala Ala Ala Asn Ile Phe Lys Asn Cys Leu | | | | | |
| | | 515 | | 520 | | 525 |
| 20 | Lys Glu Ile Asp Ser Thr Leu Tyr Lys Asn Leu Phe Val Asp Lys Asn | | | | | |
| | | 530 | | 535 | | 540 |
| | Met Lys Tyr Ile Pro Thr Glu Asp Val Ser Gly Leu Ser Leu Glu Glu | | | | | |
| | | 545 | | 550 | | 555 |
| | Gln Leu Arg Arg Leu Gln Glu Glu Arg Thr Cys Lys Val Cys Met Asp | | | | | |
| 25 | | 565 | | 570 | | 575 |
| | Lys Glu Val Ser Val Val Phe Ile Pro Cys Gly His Leu Val Val Cys | | | | | |
| | | 580 | | 585 | | 590 |
| | Gln Glu Cys Ala Pro Ser Leu Arg Lys Cys Pro Ile Cys Arg Gly Ile | | | | | |
| | | 595 | | 600 | | 605 |
| 30 | Ile Lys Gly Thr Val Arg Thr Phe Leu Ser | | | | | |
| | | 610 | | 615 | | |

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 2691 base pairs

(B) TYPE: nucleic acid

35

- 80 -

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

| | | |
|----|--|------|
| 5 | ATTTTTTAAA TTGATGCATT AACATTCTAA ACATTCATCT GTTTTTAAAT AGTAAAAAATT | 60 |
| | GAACCTTGCC TTGAATATGT AATGATTCAT TATAACAATT ATGCATAGTC TTTAATAATC | 120 |
| | TGCATATTTT ATGCTGCTTT CATGTTTTTC CTAATTAATG ACTTCACATG TTTAATATTT | 180 |
| | ATAATTTTTC TGTCATAGTT TCCATATTTA TATAAAATGA ATACTTAAGA TCAGTAATTC | 240 |
| | TGCTCTGTTT GTTTATATAC TATTTTCCAT CAAAAGACAA AATGGGACTG AGGTTGAGGC | 300 |
| 10 | TCGTTGCTAA AGCACTTTCC TAAAATGCAA AAGGCCCTAT GATGGATCCC TAGTACTTAT | 360 |
| | TTAAGTGAGA GAGAAACAGG CTGGGGGTGT AGGTCTGTTA GAGCATGTGT TTGGCATTAT | 420 |
| | GTGAAGCCCA AACACTAAAA AAGGAGAACA AACAAAAGCG CAGACTTTAA AACTCAAGTG | 480 |
| | GTTTGGAAT GTACGACTCT ACTGTTTAGA ATTAATGT GTCTTAGTTA TTGTGCCATT | 540 |
| | ATTTTATGT CATCACTGGA TAATATATTA GTGCTTAGTA TCAGAAATAG TCCTTATGCT | 600 |
| 15 | TTGTGTTTTG AAGTTCCTAA TGCAATGTTT TCTTTCTAGA AAAGGTGGAC AAGTCCTATT | 660 |
| | TTCCAGAGAA GATGACTTTT AACAGTTTTG AAGGAAGTAG AACTTTTGTA CTGTCAGACA | 720 |
| | CCAATAAGGA TGAAGAATTT GTAGAAGAGT TTAATAGATT AAAACATTT GCTAACTTCC | 780 |
| | CAAGTAGTAG TCCTGTTTCA GCATCAACAT TGGCGCGAGC TGGGTTTCTT TATACCGGTG | 840 |
| | AAGGAGACAC CGTGCAATGT TTCAGTTGTC ATGCGGCAAT AGATAGATGG CAGTATGGAG | 900 |
| 20 | ACTCAGCTGT TGGAAGACAC AGGAGAATAT CCCCAAATTG CAGATTTATC AATGGTTTTT | 960 |
| | ATTTTGAAAA TGGTGCTGCA CAGTCTACAA ATCCTGGTAT CCAAATGGC CAGTACAAAT | 1020 |
| | CTGAAAAC TGTTGGGAAAT AGAAATCCTT TTGCCCCTGA CAGGCCACCT GAGACTCATG | 1080 |
| | CTGATTATCT CTTGAGAACT GGACAGGTTG TAGATATTTT AGACACCATA TACCCGAGGA | 1140 |
| | ACCCTGCCAT GTGTAGTGAA GAAGCCAGAT TGAAGTCATT TCAGAACTGG CCGGACTATG | 1200 |
| 25 | CTCATTTAAC CCCAGAGAG TTAGCTAGTG CTGGCCTCTA CTACACAGGG GCTGATGATC | 1260 |
| | AAGTGCAATG CTTTGTGTGT GGGGGAAAAC TGAAAAATTG GGAACCCTGT GATCGTGCCT | 1320 |
| | GGTCAGAACA CAGGAGACAC TTTCCCAATT GCTTTTTTGT TTGGGCCGG AACGTTAATG | 1380 |
| | TTCAAGTGA ATCTGGTGTG AGTTCTGATA GGAATTTCCC AAATTCAACA AACTCTCCAA | 1440 |
| | GAAATCCAGC CATGGCAGAA TATGAAGCAC GGATCGTTAC TTTTGAACA TGGACATCCT | 1500 |
| 30 | CAGTTAACAA GGAGCAGCTT GCAAGAGCTG GATTTTATGC TTTAGGTGAA GGCGATAAAG | 1560 |
| | TGAAGTGCTT CCACTGTGGA GGAGGGCTCA CGGATTGGAA GCCAAGTGAA GACCCCTGGG | 1620 |
| | ACCAGCATGC TAAGTGCTAC CCAGGGTGCA AATACCTATT GGATGAGAAG GGGCAAGAAT | 1680 |
| | ATATAATAA TATTCATTTA ACCCATCCAC TTGAGGAATC TTGGGAAGA ACTGCTGAAA | 1740 |
| | AAACACCACC GCTAACTAAA AAAATCGATG ATACCATCTT CCAGAATCCT ATGGTGCAAG | 1800 |

- 81 -

AAGCTATACG AATGGGATTT AGCTTCAAGG ACCTTAAGAA AACAATGGAA GAAAAAATCC 1860
 AAACATCCGG GAGCAGCTAT CTATCACTTG AGGTCCTGAT TGCAGATCTT GTGAGTGCTC 1920
 AGAAAGATAA TACGGAGGAT GAGTCAAGTC AAAC TTCATT GCAGAAAGAC ATTAGTACTG 1980
 AAGAGCAGCT AAGGCGCCTA CAAGAGGAGA AGCTTTCCAA AATCTGTATG GATAGAAATA 2040
 5 TTGCTATCGT TTTTTTTCCT TGTGGACATC TGGCCACTTG TAAACAGTGT GCAGAAGCAG 2100
 TTGACAAATG TCCCATGTGC TACACCGTCA TTACGTTCAA CCAAAAAATT TTTATGTCTT 2160
 AGTGGGGCAC CACATGTTAT GTTCTCTTG CTCTAATTGA ATGTGTAATG GGAGCGAACT 2220
 TTAAGTAATC CTGCATTTGC ATTCCATTAG CATCCTGCTG TTTCCAAATG GAGACCAATG 2280
 CTAACAGCAC TGTTTCCGTC TAAACATTCA ATTTCTGGAT CTTTCGAGTT ATCAGCTGTA 2340
 10 TCATTTAGCC AGTGTTTTAC TCGATTGAAA CCTTAGACAG AGAAGCATT TATAGCTTTT 2400
 CACATGTATA TTGGTAGTAC ACTGACTTGA TTTCTATATG TAAGTGAATT CATCACCTGC 2460
 ATGTTTCATG CTTTTTGCAT AAGCTTAACA AATGGAGTGT TCTGTATAAG CATGGAGATG 2520
 TGATGGAATC TGCCCAATGA CTTTAATTGG CTTATTGTAA ACACGGAAAG AACTGCCCCA 2580
 CGCTGCTGGG AGGATAAAGA TTGTTTAGA TGCTCACTTC TGTGTTTTAG GATTCTGCCC 2640
 15 ATTTACTTGG AATTATTGG AGTTATAATG TACTTATATG ATATTTCCGA A 2691

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 496 amino acids
 (B) TYPE: amino acid
 20 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met Thr Phe Asn Ser Phe Glu Gly Thr Arg Thr Phe Val Leu Ala Asp
 25 1 5 10 15
 Thr Asn Lys Asp Glu Glu Phe Val Glu Glu Phe Asn Arg Leu Lys Thr
 20 25 30
 Phe Ala Asn Phe Pro Ser Ser Ser Pro Val Ser Ala Ser Thr Leu Ala
 35 40 45
 30 Arg Ala Gly Phe Leu Tyr Thr Gly Glu Gly Asp Thr Val Gln Cys Phe
 50 55 60
 Ser Cys His Ala Ala Ile Asp Arg Trp Gln Tyr Gly Asp Ser Ala Val

- 82 -

65 70 75 80
 Gly Arg His Arg Arg Ile Ser Pro Asn Cys Arg Phe Ile Asn Gly Phe
 85 90 95
 Tyr Phe Glu Asn Gly Ala Ala Gln Ser Thr Asn Pro Gly Ile Gln Asn
 5 100 105 110
 Gly Gln Tyr Lys Ser Glu Asn Cys Val Gly Asn Arg Asn Pro Phe Ala
 115 120 125
 Pro Asp Arg Pro Pro Glu Thr His Ala Asp Tyr Leu Leu Arg Thr Gly
 130 135 140
 10 Gln Val Val Asp Ile Ser Asp Thr Ile Tyr Pro Arg Asn Pro Ala Met
 145 150 155 160
 Cys Ser Glu Glu Ala Arg Leu Lys Ser Phe Gln Asn Trp Pro Asp Tyr
 165 170 175
 Ala His Leu Thr Pro Arg Glu Leu Ala Ser Ala Gly Leu Tyr Tyr Thr
 15 180 185 190
 Gly Ala Asp Asp Gln Val Gln Cys Phe Cys Cys Gly Gly Lys Leu Lys
 195 200 205
 Asn Trp Glu Pro Cys Asp Arg Ala Trp Ser Glu His Arg Arg His Phe
 210 215 220
 20 Pro Asn Cys Phe Phe Val Leu Gly Arg Asn Val Asn Val Arg Ser Glu
 225 230 235 240
 Ser Gly Val Ser Ser Asp Arg Asn Phe Pro Asn Ser Thr Asn Ser Pro
 245 250 255
 Arg Asn Pro Ala Met Ala Glu Tyr Glu Ala Arg Ile Val Thr Phe Gly
 25 260 265 270
 Thr Trp Ile Tyr Ser Val Asn Lys Glu Gln Leu Ala Arg Ala Gly Phe
 275 280 285
 Tyr Ala Leu Gly Glu Gly Asp Lys Val Lys Cys Phe His Cys Gly Gly
 290 295 300
 30 Gly Leu Thr Asp Trp Lys Pro Ser Glu Asp Pro Trp Asp Gln His Ala
 305 310 315 320
 Lys Cys Tyr Pro Gly Cys Lys Tyr Leu Leu Asp Glu Lys Gly Gln Glu
 325 330 335
 Tyr Ile Asn Asn Ile His Leu Thr His Pro Leu Glu Glu Ser Leu Gly
 35 340 345 350
 Arg Thr Ala Glu Lys Thr Pro Pro Leu Thr Lys Lys Ile Asp Asp Thr
 355 360 365

- 83 -

Ile Phe Gln Asn Pro Met Val Gln Glu Ala Ile Arg Met Gly Phe Ser
 370 375 380
 Phe Lys Asp Leu Lys Lys Thr Met Glu Glu Lys Ile Gln Thr Ser Gly
 385 390 395 400
 5 Ser Ser Tyr Leu Ser Leu Glu Val Leu Ile Ala Asp Leu Val Ser Ala
 405 410 415
 Gln Lys Asp Asn Thr Glu Asp Glu Ser Ser Gln Thr Ser Leu Gln Lys
 420 425 430
 Asp Ile Ser Thr Glu Glu Gln Leu Arg Arg Leu Gln Glu Glu Lys Leu
 10 435 440 445
 Ser Lys Ile Cys Met Asp Arg Asn Ile Ala Ile Val Phe Phe Pro Cys
 450 455 460
 Gly His Leu Ala Thr Cys Lys Gln Cys Ala Glu Ala Val Asp Lys Cys
 465 470 475 480
 15 Pro Met Cys Tyr Thr Val Ile Thr Phe Asn Gln Lys Ile Phe Met Ser
 485 490 495

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- 20 (A) LENGTH: 2676 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

25 TGGGAGTTCC CCGGAGCCCT GGAGGAAAGC ACCGCAGGTC TGAGCAGCCC TGAGCCGGGC 60
 AGGGTGGGG CAGTGGCTAA GGCCTAGCTG GGGACGATTT AAAGGTATCG CGCCACCCAG 120
 CCACACCCCA CAGGCCAGGC GAGGGTGCCA CCCCCGAGA TCAGAGGTCA TTGCTGGCGT 180
 TCAGAGCCTA GGAAGTGGGC TCGGTATCA GCCTAGCAGT AAAACCGACC AGAAGCCATG 240
 CACAAACTA CATCCCCAGA GAAAGACTTG TCCCTTCCCC TCCCTGTCAT CTCACCATGA 300
 30 ACATGGTTCA AGACAGCGCC TTTCTAGCCA AGCTGATGAA GAGTGCTGAC ACCTTTGAGT 360
 TGAAGTATGA CTTTTCCTGT GAGCTGTACC GATTGTCCAC GTATTGAGT TTTCCAGGG 420
 GAGTTCCTGT GTCAGAAAGG AGTCTGGCTC GTGCTGGCTT TTACTACACT GGTGCCAATG 480

- 84 -

| | | |
|----|---|------|
| | ACAAGGTCAA GTGCTTCTGC TGTGGCCTGA TGCTAGACAA CTGGAAACAA GGGGACAGTC | 540 |
| | CCATGGAGAA GCACAGAAAG TTGTACCCCA GCTGCAACTT TGTACAGACT TTGAATCCAG | 600 |
| | CCAACAGTCT GGAAGCTAGT CCTCGGCCTT CTCTTCCTTC CACGGCGATG AGCACCATGC | 660 |
| | CTTTGAGCTT TGCAAGTTCT GAGAATACTG GCTATTTCAG TGGCTCTTAC TCGAGCTTTC | 720 |
| 5 | CCTCAGACCC TGTGAAC TTCGAGCAAATC AAGATTGTCC TGCTTTGAGC ACAAGTCCCT | 780 |
| | ACCACTTTGC AATGAACACA GAGAAGGCCA GATTACTCAC CTATGAAACA TGGCCATTGT | 840 |
| | CTTTTCTGTC ACCAGCAAAG CTGGCCAAAG CAGGCTTCTA CTACATAGGA CCTGGAGATA | 900 |
| | GAGTGGCCTG CTTTGCGTGC GATGGGAAAC TGAGCAACTG GGAACGTAAG GATGATGCTA | 960 |
| | TGTCAGAGCA CCAGAGGCAT TTCCCCAGCT GTCCGTTCTT AAAAGACTTG GGTCACTCTG | 1020 |
| 10 | CTTCGAGATA CACTGTCTCT AACCTGAGCA TGCAGACACA CGCAGCCCGT ATTAGAACAT | 1080 |
| | TCTCTAACTG GCCTTCTAGT GCACTAGTTC ATTCACAGGA ACTTGCAAGT GCGGGCTTTT | 1140 |
| | ATTATACAGG ACACAGTGAT GATGTCAAGT GTTTTGTCTG TGATGGTGGG CTGAGGTGCT | 1200 |
| | GGGAATCTGG AGATGACCCC TGGGTGGAAC ATGCCAAGTG GTTTCCAAGG TGTGAGTACT | 1260 |
| | TGCTCAGAAT CAAAGGCCAA GAATTGTGCA GCCAAGTTCA AGCTGGCTAT CCTCATCTAC | 1320 |
| 15 | TTGAGCAGCT ATTATCTACG TCAGACTCCC CAGAAGATGA GAATGCAGAC GCAGCAATCG | 1380 |
| | TGCATTTTGG CCCTGGAGAA AGTTCGGAAG ATGTCGTCAT GATGAGCACG CCTGTGGTTA | 1440 |
| | AAGCAGCCTT GGAAATGGGC TTCAGTAGGA GCCTGGTGAG ACAGACGGTT CAGCGGCAGA | 1500 |
| | TCCTGGCCAC TGGTGAGAAC TACAGGACCG TCAGTGACCT CGTTATAGGC TTA CTGATG | 1560 |
| | CAGAAGACGA GATGAGAGAG GAGCAGATGG AGCAGGCGGC CGAGGAGGAG GAGTCAGATG | 1620 |
| 20 | ATCTAGCACT AATCCGGAAG AACAAAATGG TGCTTTTCCA ACATTTGACG TGTGTGACAC | 1680 |
| | CAATGCTGTA TTGCCTCCTA AGTGCAAGGG CCATCACTGA ACAGGAGTGC AATGCTGTGA | 1740 |
| | AACAGAAACC ACACACCTTA CAAGCAAGCA CACTGATTGA TACTGTGTTA GCAAAAGGAA | 1800 |
| | ACACTGCAGC AACCTCATT CAGAACTCCC TTCGGGAAAT TGACCCTGCG TTATACAGAG | 1860 |
| | ATATATTTGT GCAACAGGAC ATTAGGAGTC TTCCACAGA TGACATTGCA GCTCTACCAA | 1920 |
| 25 | TGGAAGAACA GTTGCGGAAA CTCCAGGAGG AAAGAATGTG TAAAGTGTGT ATGGACCGAG | 1980 |
| | AGGTATCCAT CGTGTTCAAT CCCTGTGGCC ATCTGGTCGT GTGCAAAGAC TGCGCTCCCT | 2040 |
| | CTCTGAGGAA GTGTCCATC TGTAGAGGGA CCATCAAGGG CACAGTGC GCACATTTCTCT | 2100 |
| | CCTGAACAAG ACTAATGGTC CATGGCTGCA ACTTCAGCCA GGAGGAAGTT CACTGTCACT | 2160 |
| | CCCAGCTCCA TTCGGAACCT GAGGCCAGCC TGGATAGCAC GAGACACCGC CAAACACACA | 2220 |
| 30 | AATATAAACA TGAAAACTT TTGTCTGAAG TCAAGAATGA ATGAATTACT TATATAATAA | 2280 |
| | TTTAAATTGG TTTCTTAAAG AGTGCTATTT GTTCCCAACT CAGAAAATTG TTTTCTGTAA | 2340 |
| | ACATATTTAC ATACTACCTG CATCTAAAGT ATTCATATAT TCATATATTC AGATGTCATG | 2400 |
| | AGAGAGGGTT TTGTTCTTGT TCCTGAAAAG CAGGGATTGC CTGCACTCCT GAAATTCTCA | 2460 |
| | GAAAGATTTA CAATGTTGGC ATTTATGGTT CAGAACTAG AATCTTCTCC CGTTGCTTTA | 2520 |
| 35 | AGAACCGGGA GCACAGATGT CCATGTGTTT TATGTATAGA AATTCCTGTT ATTTATTGGA | 2580 |
| | TGACATTTTA GGGATATGAA ATTTTATAA AGAATTTGTG AGAAAAAGTT AATAAAGCAA | 2640 |
| | CATAATTACC TCTTTTTTTT TAAAGAAAAA AAAAAA | 2676 |

- 85 -

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 600 amino acids

(B) TYPE: amino acid

5 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met Val Gln Asp Ser Ala Phe Leu Ala Lys Leu Met Lys Ser Ala Asp
 10 1 5 10 15
 Thr Phe Glu Leu Lys Tyr Asp Phe Ser Cys Glu Leu Tyr Arg Leu Ser
 20 25 30
 Thr Tyr Ser Ala Phe Pro Arg Gly Val Pro Val Ser Glu Arg Ser Leu
 35 40 45
 15 Ala Arg Ala Gly Phe Tyr Tyr Thr Gly Ala Asn Asp Lys Val Lys Cys
 50 55 60
 Phe Cys Cys Gly Leu Met Leu Asp Asn Trp Lys Gln Gly Asp Ser Pro
 65 70 75 80
 Met Glu Lys His Arg Lys Leu Tyr Pro Ser Cys Asn Phe Val Gln Thr
 20 85 90 95
 Leu Asn Pro Ala Asn Ser Leu Glu Ala Ser Pro Arg Pro Ser Leu Pro
 100 105 110
 Ser Thr Ala Met Ser Thr Met Pro Leu Ser Phe Ala Ser Ser Glu Asn
 115 120 125
 25 Thr Gly Tyr Phe Ser Gly Ser Tyr Ser Ser Phe Pro Ser Asp Pro Val
 130 135 140
 Asn Phe Arg Ala Asn Gln Asp Cys Pro Ala Leu Ser Thr Ser Pro Tyr
 145 150 155 160
 His Phe Ala Met Asn Thr Glu Lys Ala Arg Leu Leu Thr Tyr Glu Thr
 30 165 170 175
 Trp Pro Leu Ser Phe Leu Ser Pro Ala Lys Leu Ala Lys Ala Gly Phe
 180 185 190
 Tyr Tyr Ile Gly Pro Gly Asp Arg Val Ala Cys Phe Ala Cys Asp Gly

- 86 -

| | | | |
|----|---|-------------------------|-------------|
| | 195 | 200 | 205 |
| | Lys Leu Ser Asn Trp Glu Arg | Lys Asp Asp Ala Met Ser | Glu His Gln |
| | 210 | 215 | 220 |
| | Arg His Phe Pro Ser Cys Pro Phe Leu Lys Asp Leu Gly Gln Ser Ala | | |
| 5 | 225 | 230 | 235 240 |
| | Ser Arg Tyr Thr Val Ser Asn Leu Ser Met Gln Thr His Ala Ala Arg | | |
| | 245 | 250 | 255 |
| | Ile Arg Thr Phe Ser Asn Trp Pro Ser Ser Ala Leu Val His Ser Gln | | |
| | 260 | 265 | 270 |
| 10 | Glu Leu Ala Ser Ala Gly Phe Tyr Tyr Thr Gly His Ser Asp Asp Val | | |
| | 275 | 280 | 285 |
| | Lys Cys Phe Cys Cys Asp Gly Gly Leu Arg Cys Trp Glu Ser Gly Asp | | |
| | 290 | 295 | 300 |
| | Asp Pro Trp Val Glu His Ala Lys Trp Phe Pro Arg Cys Glu Tyr Leu | | |
| 15 | 305 | 310 | 315 320 |
| | Leu Arg Ile Lys Gly Gln Glu Phe Val Ser Gln Val Gln Ala Gly Tyr | | |
| | 325 | 330 | 335 |
| | Pro His Leu Leu Glu Gln Leu Leu Ser Thr Ser Asp Ser Pro Glu Asp | | |
| | 340 | 345 | 350 |
| 20 | Glu Asn Ala Asp Ala Ala Ile Val His Phe Gly Pro Gly Glu Ser Ser | | |
| | 355 | 360 | 365 |
| | Glu Asp Val Val Met Met Ser Thr Pro Val Val Lys Ala Ala Leu Glu | | |
| | 370 | 375 | 380 |
| | Met Gly Phe Ser Arg Ser Leu Val Arg Gln Thr Val Gln Arg Gln Ile | | |
| 25 | 385 | 390 | 395 400 |
| | Leu Ala Thr Gly Glu Asn Tyr Arg Thr Val Ser Asp Leu Val Ile Gly | | |
| | 405 | 410 | 415 |
| | Leu Leu Asp Ala Glu Asp Glu Met Arg Glu Glu Gln Met Glu Gln Ala | | |
| | 420 | 425 | 430 |
| 30 | Ala Glu Glu Glu Glu Ser Asp Asp Leu Ala Leu Ile Arg Lys Asn Lys | | |
| | 435 | 440 | 445 |
| | Met Val Leu Phe Gln His Leu Thr Cys Val Thr Pro Met Leu Tyr Cys | | |
| | 450 | 455 | 460 |
| | Leu Leu Ser Ala Arg Ala Ile Thr Glu Gln Glu Cys Asn Ala Val Lys | | |
| 35 | 465 | 470 | 475 480 |
| | Gln Lys Pro His Thr Leu Gln Ala Ser Thr Leu Ile Asp Thr Val Leu | | |
| | 485 | 490 | 495 |

- 87 -

Ala Lys Gly Asn Thr Ala Ala Thr Ser Phe Arg Asn Ser Leu Arg Glu
 500 505 510
 Ile Asp Pro Ala Leu Tyr Arg Asp Ile Phe Val Gln Gln Asp Ile Arg
 515 520 525
 5 Ser Leu Pro Thr Asp Asp Ile Ala Ala Leu Pro Met Glu Glu Gln Leu
 530 535 540
 Arg Lys Leu Gln Glu Glu Arg Met Cys Lys Val Cys Met Asp Arg Glu
 545 550 555 560
 Val Ser Ile Val Phe Ile Pro Cys Gly His Leu Val Val Cys Lys Asp
 10 565 570 575
 Cys Ala Pro Ser Leu Arg Lys Cys Pro Ile Cys Arg Gly Thr Ile Lys
 580 585 590
 Gly Thr Val Arg Thr Phe Leu Ser
 595 600

15 (2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3151 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 20 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

AGTTATATAA AATACGAAGT TTTCAAAAAG AAGGCTAGTG CAACAGAAAA GCTTTGCTAA 60
 AACAGATTCT TAGTTATTTG AGGTAACAAA AGAAAGCCAT GTCTTGAATT GATTCGTTCT 120
 25 TAATTATAAC AGACTTATAG TGGAAAGGGC CTTAAACACA GGCGGACTTT ATAAAATGCA 180
 GTCTTAGGTT TATGTGCAAA ATACTGTCTG TTGACCAGAT GTATTCACAT GATATATACA 240
 GAGTCAAGGT GGTGATATAG AAGATTAAAC AGTGAGGGAG TTAACAGTCT GTGCTTTAAG 300
 CGCAGTTCCT TTACAGTGAA TACTGTAGTC TTAATAGACC TGAGCTGACT GCTGCAGTTG 360
 ATGTAACCCA CTTTAGAGAA TACTGTATGA CATCTTCTCT AAGGAAAACC AGCTGCAGAC 420
 30 TTCACTCAGT TCCTTTCATT TCATAGGAAA AGGAGTAGTT CAGATGTCAT GTTTAAGTCC 480
 TTATAAGGGA AAAGAGCCTG AATATATGCC CTAGTACCTA GGCTTCATAA CTAGTAATAA 540
 GAAGTTAGTT ATGGGTAAAT AGATCTCAGG TTACCCAGAA GAGTTCATGT GACCCCCAAA 600

- 88 -

| | | |
|----|--|------|
| | GAGTCCTAAC TAGTGTCTTG GCAAGTGAGA CAGATTTGTC CTGTGAGGGT GTCAATTCAC | 660 |
| | CAGTCCAAGC AGAAGACAAT GAATCTATCC AGTCAGGTGT CTGTGGTGGA GATCTAGTGT | 720 |
| | CCAAGTGGTG AGAAACTTCA TCTGGAAGTT TAAGCGGTCA GAAATACTAT TACTACTCAT | 780 |
| | GGACAAAAC GTCTCCCAGA GACTCGGCCA AGGTACCTTA CACCAAAAAC TTAAACGTAT | 840 |
| 5 | AATGGAGAAG AGCACAATCT TGTCAAATTG GACAAAGGAG AGCGAAGAAA AAATGAAGTT | 900 |
| | TGACTTTTCG TGTGAATCT ACCGAATGTC TACATATTCA GCTTTTCCCA GGGGAGTTCC | 960 |
| | TGTCTCAGAG AGGAGTCTGG CTCGTGCTGG CTTTTATTAT ACAGGTGTGA ATGACAAAGT | 1020 |
| | CAAGTGCTTC TGCTGTGGCC TGATGTTGGA TAAGTGAAA CAAGGGGACA GTCCTGTTGA | 1080 |
| | AAAGCACAGA CAGTTCTATC CCAGCTGCAG CTTTGTACAG ACTCTGCTTT CAGCCAGTCT | 1140 |
| 10 | GCAGTCTCCA TCTAAGAATA TGTCTCCTGT GAAAAGTAGA TTTGCACATT CGTCACCTCT | 1200 |
| | GGAACGAGGT GGCATTCACCT CCAACCTGTG CTCTAGCCCT CTTAATTCTA GAGCAGTGGA | 1260 |
| | AGACTTCTCA TCAAGGATGG ATCCCTGCAG CTATGCCATG AGTACAGAAG AGGCCAGATT | 1320 |
| | TCTTACTTAC AGTATGTGGC CTTTAAGTTT TCTGTCACCA GCAGAGCTGG CCAGAGCTGG | 1380 |
| | CTTCTATTAC ATAGGGCCTG GAGACAGGGT GGCCTGTTTT GCCTGTGGTG GGAAACTGAG | 1440 |
| 15 | CAACTGGGAA CCAAAGGATG ATGCTATGTC AGAGCACCGC AGACATTTTC CCCACTGTCC | 1500 |
| | ATTCTCTGAA AATACTTCAG AAACACAGAG GTTTAGTATA TCAAATCTAA GTATGCAGAC | 1560 |
| | ACACTCTGCT CGATTGAGGA CATTTCTGTA CTGGCCACCT AGTGTTCCCTG TTCAGCCCGA | 1620 |
| | GCAGCTTGCA AGTGCTGGAT TCTATTACGT GGATCGCAAT GATGATGTCA AGTGCTTTTG | 1680 |
| | TTGTGATGGT GGCTTGAGAT GTTGGGAACC TGGAGATGAC CCCTGGATAG AACACGCCAA | 1740 |
| 20 | ATGGTTTCCA AGGTGTGAGT TCTTGATACG GATGAAGGGT CAGGAGTTTG TTGATGAGAT | 1800 |
| | TCAAGCTAGA TATCTCATC TTCTTGAGCA GCTGTTGTCC ACTTCAGACA CCCAGGAGA | 1860 |
| | AGAAAATGCT GACCCTACAG AGACAGTGGT GCATTTTGGC CCTGGAGAAA GTTCGAAAGA | 1920 |
| | TGTCGTCATG ATGAGCACGC CTGTGGTTAA AGCAGCCTTG GAAATGGGCT TCAGTAGGAG | 1980 |
| | CCTGGTGAGA CAGACGGTTC AGCGGCAGAT CCTGGCCACT GGTGAGAACT ACAGGACCGT | 2040 |
| 25 | CAATGATATT GTCTCAGTAC TTTTGAATGC TGAAGATGAG AGAAGAGAAG AGGAGAAGGA | 2100 |
| | AAGACAGACT GAAGAGATGG CATCAGGTGA CTTATCACTG ATTCGGAAGA ATAGAATGGC | 2160 |
| | CCTCTTTCAA CAGTTGACAC ATGTCCTTCC TATCCTGGAT AATCTTCTTG AGGCCAGTGT | 2220 |
| | AATTACAAAA CAGGAACATG ATATTATTAG ACAGAAAACA CAGATACCCT TACAAGCAAG | 2280 |
| | AGAGCTTATT GACACCGTTT TAGTCAAGGG AAATGCTGCA GCCAACATCT TCAAAAATC | 2340 |
| 30 | TCTGAAGGAA ATTGACTCCA CGTTATATGA AAACCTATTT GTGGAAAAGA ATATGAAGTA | 2400 |
| | TATTCACAAC GAAGACGTTT CAGGCTTGTC ATTGGAAGAG CAGTTGCGGA GATTACAAGA | 2460 |
| | AGAACGAACT TGCAAAGTGT GTATGGACAG AGAGGTTTCT ATTGTGTTCA TTCCGTGTGG | 2520 |
| | TCATCTAGTA GTCTGCCAGG AATGTGCCCC TTCTCTAAGG AAGTGCCCCA TCTGCAGGGG | 2580 |
| | GACAAATCAAG GGGACTGTGC GCACATTTCT CTCATGAGTG AAGAATGGTC TGAAAGTATT | 2640 |
| 35 | GTTGGACATC AGAAGCTGTC AGAACAAAGA ATGAACTACT GATTTCAGCT CTTCAGCAGG | 2700 |
| | ACATTCTACT CTCTTTCAAG ATTAGTAATC TTGCTTTATG AAGGGTAGCA TTGTATATTT | 2760 |
| | AAGCTTAGTC TGTTGCAAGG GAAGGTCTAT GCTGTTGAGC TACAGGACTG TGTCTGTTCC | 2820 |

- 89 -

AGAGCAGGAG TTGGGATGCT TGCTGTATGT CCTTCAGGAC TTCTTGGATT TGGAATTTGT 2880
 GAAAGCTTTG GATTCAGGTG ATGTGGAGCT CAGAAATCCT GAAACCAGTG GCTCTGGTAC 2940
 TCAGTAGTTA GGGTACCCTG TGCTTCTTGG TGCTTTTCCT TTCTGGAAAA TAAGGATTTT 3000
 TCTGCTACTG GTAAATATTT TCTGTTTGTG AGAAATATAT TAAAGTGTTC CTTTAAAGG 3060
 5 CGTGCATCAT TGTAGTGTGT GCAGGGATGT ATGCAGGCAA AACACTGTGT ATATAATAAA 3120
 TAAATCTTTT TAAAAAGTGT AAAAAAAAAA A 3151

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 612 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

15 Met Asp Lys Thr Val Ser Gln Arg Leu Gly Gln Gly Thr Leu His Gln
 1 5 10 15
 Lys Leu Lys Arg Ile Met Glu Lys Ser Thr Ile Leu Ser Asn Trp Thr
 20 25 30
 Lys Glu Ser Glu Glu Lys Met Lys Phe Asp Phe Ser Cys Glu Leu Tyr
 35 40 45
 Arg Met Ser Thr Tyr Ser Ala Phe Pro Arg Gly Val Pro Val Ser Glu
 50 55 60
 Arg Ser Leu Ala Arg Ala Gly Phe Tyr Tyr Thr Gly Val Asn Asp Lys
 65 70 75 80
 25 Val Lys Cys Phe Cys Cys Gly Leu Met Leu Asp Asn Trp Lys Gln Gly
 85 90 95
 Asp Ser Pro Val Glu Lys His Arg Gln Phe Tyr Pro Ser Cys Ser Phe
 100 105 110
 Val Gln Thr Leu Leu Ser Ala Ser Leu Gln Ser Pro Ser Lys Asn Met
 30 115 120 125
 Ser Pro Val Lys Ser Arg Phe Ala His Ser Ser Pro Leu Glu Arg Gly
 130 135 140

- 90 -

Gly Ile His Ser Asn Leu Cys Ser Ser Pro Leu Asn Ser Arg Ala Val
 145 150 155 160
 Glu Asp Phe Ser Ser Arg Met Asp Pro Cys Ser Tyr Ala Met Ser Thr
 165 170 175
 5 Glu Glu Ala Arg Phe Leu Thr Tyr Ser Met Trp Pro Leu Ser Phe Leu
 180 185 190
 Ser Pro Ala Glu Leu Ala Arg Ala Gly Phe Tyr Tyr Ile Gly Pro Gly
 195 200 205
 Asp Arg Val Ala Cys Phe Ala Cys Gly Gly Lys Leu Ser Asn Trp Glu
 10 210 215 220
 Pro Lys Asp Asp Ala Met Ser Glu His Arg Arg His Phe Pro His Cys
 225 230 235 240
 Pro Phe Leu Glu Asn Thr Ser Glu Thr Gln Arg Phe Ser Ile Ser Asn
 245 250 255
 15 Leu Ser Met Gln Thr His Ser Ala Arg Leu Arg Thr Phe Leu Tyr Trp
 260 265 270
 Pro Pro Ser Val Pro Val Gln Pro Glu Gln Leu Ala Ser Ala Gly Phe
 275 280 285
 Tyr Tyr Val Asp Arg Asn Asp Asp Val Lys Cys Phe Cys Cys Asp Gly
 20 290 295 300
 Gly Leu Arg Cys Trp Glu Pro Gly Asp Asp Pro Trp Ile Glu His Ala
 305 310 315 320
 Lys Trp Phe Pro Arg Cys Glu Phe Leu Ile Arg Met Lys Gly Gln Glu
 325 330 335
 25 Phe Val Asp Glu Ile Gln Ala Arg Tyr Pro His Leu Leu Glu Gln Leu
 340 345 350
 Leu Ser Thr Ser Asp Thr Pro Gly Glu Glu Asn Ala Asp Pro Thr Glu
 355 360 365
 Thr Val Val His Phe Gly Pro Gly Glu Ser Ser Lys Asp Val Val Met
 30 370 375 380
 Met Ser Thr Pro Val Val Lys Ala Ala Leu Glu Met Gly Phe Ser Arg
 385 390 395 400
 Ser Leu Val Arg Gln Thr Val Gln Arg Gln Ile Leu Ala Thr Gly Glu
 405 410 415
 35 Asn Tyr Arg Thr Val Asn Asp Ile Val Ser Val Leu Leu Asn Ala Glu
 420 425 430
 Asp Glu Arg Arg Glu Glu Glu Lys Glu Arg Gln Thr Glu Glu Met Ala

- 91 -

```

          435          440          445
Ser Gly Asp Leu Ser Leu Ile Arg Lys Asn Arg Met Ala Leu Phe Gln
          450          455          460
Gln Leu Thr His Val Leu Pro Ile Leu Asp Asn Leu Leu Glu Ala Ser
5 465          470          475          480
Val Ile Thr Lys Gln Glu His Asp Ile Ile Arg Gln Lys Thr Gln Ile
          485          490          495
Pro Leu Gln Ala Arg Glu Leu Ile Asp Thr Val Leu Val Lys Gly Asn
          500          505          510
10 Ala Ala Ala Asn Ile Phe Lys Asn Ser Leu Lys Glu Ile Asp Ser Thr
          515          520          525
Leu Tyr Glu Asn Leu Phe Val Glu Lys Asn Met Lys Tyr Ile Pro Thr
          530          535          540
Glu Asp Val Ser Gly Leu Ser Leu Glu Glu Gln Leu Arg Arg Leu Gln
15 545          550          555          560
Glu Glu Arg Thr Cys Lys Val Cys Met Asp Arg Glu Val Ser Ile Val
          565          570          575
Phe Ile Pro Cys Gly His Leu Val Val Cys Gln Glu Cys Ala Pro Ser
          580          585          590
20 Leu Arg Lys Cys Pro Ile Cys Arg Gly Thr Ile Lys Gly Thr Val Arg
          595          600          605
Thr Phe Leu Ser
          610

```

(2) INFORMATION FOR SEQ ID NO:15:

25 (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

30 (ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

AGTGCGGGTT TTTATTATGT G

- 92 -

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Other

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

AGATGACCAC AAGGAATAAA CACTA

25

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 11 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Met Glu Gln Lys Leu Ile Ser Glu Glu Asp Leu

1

5

10

- 93 -

What is claimed is:

Claims

1. A method for enhancing apoptosis in a cell from a mammal with a proliferative disease, said method comprising administering to said cell a compound that inhibits the
5 biological activity of an IAP polypeptide or an NAIP polypeptide, said compound being administered to said cell in an amount sufficient to enhance apoptosis in said cell.
2. The method of claim 1, wherein said cell is proliferating in said proliferative disease.
3. The method of claim 1, wherein said biological activity is the level of expression
10 of said polypeptide; wherein said biological activity is the level of expression of an mRNA molecule encoding said polypeptide; or wherein said biological activity is an apoptosis-inhibiting activity.
4. The method of claim 3, wherein said level of expression is measured by assaying the amount of said polypeptide present in said cell.
- 15 5. The method of claim 1, wherein said polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP.
6. The method of claim 1, wherein said polypeptide is NAIP.
7. The method of claim 1, wherein said polypeptide is XIAP.
8. The method of claim 1, wherein said polypeptide is HIAP-1.
- 20 9. The method of claim 1, wherein said polypeptide is HIAP-2.
10. The method of claim 1, wherein said compound is a negative regulator of an IAP or an NAIP-dependent anti-apoptotic pathway; wherein said compound is a fragment of said

- 94 -

IAP polypeptide, said fragment comprising a ring zinc finger and having no more than two BIR domains; wherein said compound is a nucleic acid molecule encoding a ring zinc finger domain of said IAP polypeptide; wherein said compound is a compound that prevents cleavage of said IAP polypeptide or said NAIP polypeptide; wherein said compound is a
5 purified antibody or a fragment thereof that specifically binds to said IAP polypeptide or said NAIP polypeptide; wherein said compound is a ribozyme; or wherein said compound is an antisense nucleic acid molecule have a nucleic acid sequence that is complementary to the coding strand of a nucleic acid sequence encoding said IAP polypeptide or said NAIP polypeptide.

10 11. The method of claim 10, wherein said cleavage is decreased by at least 20% in said cell.

12. The method of claim 10, wherein said antibody binds to a BIR domain of said IAP polypeptide or said NAIP polypeptide.

13. The method of claim 10, wherein said nucleic acid sequence encoding said IAP
15 polypeptide or said NAIP polypeptide has about 50% or greater identity with the nucleotide sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, or the nucleic acid sequence of NAIP.

14. The method of claim 10, wherein said antisense nucleic acid molecule decreases the level of said nucleic acid sequence encoding said IAP polypeptide or said NAIP
20 polypeptide by at least 20%, said level being measured in the cytoplasm of said cell.

15. The method of claim 10, wherein said antisense nucleic acid molecule is encoded by a virus vector.

16. The method of claim 10, wherein said antisense nucleic acid molecule is encoded by a transgene.

- 95 -

17. The method of claim 1, wherein said mammal is a human or a mouse.

18. The method of claim 1, wherein said proliferative disease is cancer.

19. The method of claim 18, wherein said cancer is in a tissue selected from the group consisting of ovary, breast, pancreas, lymph node, skin, blood, lung, brain, kidney, liver, nasopharyngeal cavity, thyroid, central nervous system, prostate, colon, rectum, cervix, endometrium, and lung.

20. A method for detecting a proliferative disease or an increased likelihood of said proliferative disease in a mammal, said method comprising:

(a) contacting an IAP or a NAIP nucleic acid molecule that is greater than about 18 nucleotides in length with a preparation of nucleic acid from a cell of said mammal, said cell proliferating in said disease, said cell from a tissue; and

(b) measuring the amount of nucleic acid from said cell of said mammal that hybridizes to said molecule, an increase in the amount from said cell of said mammal relative to a control indicating a an increased likelihood of said mammal having or developing a proliferative disease.

21. The method of claim 20, wherein said method further comprises the steps of:

(a) contacting said molecule with a preparation of nucleic acid from said control, wherein said control is a cell from said tissue of a second mammal, said second mammal lacking a proliferative disease; and

(b) measuring the amount of nucleic acid from said control, an increase in the amount of said nucleic acid from said cell of said mammal that hybridizes to said molecule relative to said amount of said nucleic acid from said control indicating an increased likelihood of said mammal having or developing a proliferative disease.

22. The method of claim 20 or 21, said method further comprising the steps of:

(a) providing a pair of oligonucleotides having sequence identity to or being complementary to a region of said IAP or said NAIP nucleic acid molecule;

- 96 -

- (b) combining said pair of oligonucleotides with said nucleic acid under conditions suitable for polymerase chain reaction-mediated nucleic acid amplification; and
- (c) isolating said amplified nucleic acid or fragment thereof.

23. The method of claim 22, wherein said amplification is carried out using a
5 reverse-transcription polymerase chain reaction.

24. The method of claim 23, wherein said reverse-transcription polymerase chain reaction is RACE.

25. The method of claim 20, 21, or 22, wherein said method provides measuring said nucleic acid having a nucleotide sequence that has about 50% or greater identity with the
10 nucleotide sequence of SEQ ID NO: 3, SEQ ID NO: 5, SEQ ID NO: 7, SEQ ID NO: 9, SEQ ID NO: 11, SEQ ID NO: 13, or the nucleic acid sequence of NAIP.

26. The method of claim 20, 21, or 22, wherein said method provides measuring said nucleic acid having a nucleotide sequence that has about 50% or greater identity with the nucleotide sequence of SEQ ID NO: 3.

15 27. The method of claim 20, 21, or 22, wherein said method provides measuring said nucleic acid having a nucleotide sequence that has about 50% or greater identity with the nucleotide sequence of SEQ ID NO: 5.

28. The method of claim 20, 21, or 22, wherein said method provides measuring said nucleic acid having a nucleotide sequence that has about 50% or greater identity with the
20 nucleotide sequence of SEQ ID NO: 7.

29. The method of claim 20, 21, or 22, wherein said method provides measuring said nucleic acid having a nucleotide sequence that has about 50% or greater identity with the nucleotide sequence of NAIP.

- 97 -

30. A method for detecting a proliferative disease or an increased likelihood of developing said disease in a mammal, said method comprising measuring the level of biological activity of an IAP polypeptide or a NAIP polypeptide in a sample of said mammal, an increase in said level of said IAP polypeptide or said NAIP polypeptide relative to a
5 sample from a control mammal being an indication that said mammal has said disease or increased likelihood of developing said disease.

31. The method of claim 30, wherein said sample comprises a cell that is proliferating in said disease from said mammal, said cell from a tissue.

32. The method of claim 31, wherein said sample from a control mammal is from
10 said tissue, said sample consisting of healthy cells.

33. The method of claim 32, wherein said mammal and said control mammal are the same.

34. The method of claim 30, wherein said biological activity is the level of
15 expression of said polypeptide; wherein said biological activity is the level of expression of an mRNA molecule encoding said polypeptide; or wherein said biological activity is an apoptosis-inhibiting activity.

35. The method of claim 34, wherein said level of expression is measured by assaying the amount of said polypeptide present in said cell.

20 36. The method of claim 30, wherein said polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP.

37. The method of claim 30, wherein said polypeptide is NAIP.

38. The method of claim 30, wherein said polypeptide is XIAP.

- 98 -

39. The method of claim 30, wherein said polypeptide is HIAP-1.

40. The method of claim 30, wherein said polypeptide is HIAP-2.

41. A method for identifying a compound enhances apoptosis in an affected cell that is proliferating in a proliferative disease, said method comprising exposing a cell that
5 overexpresses an IAP polypeptide or a NAIP polypeptide to a candidate compound, a decrease the level of biological activity of said polypeptide indicating the presence of a compound that enhances apoptosis in said affected cell that is proliferating in said proliferative disease.

42. A method for identifying a compound that enhances apoptosis in an affected cell
10 that is proliferating in a proliferative disease, said method comprising the steps of:

(a) providing a cell comprising a nucleic acid molecule encoding a IAP polypeptide or a nucleic acid molecule encoding a NAIP polypeptide, said nucleic acid molecule being expressed in said cell; and

(b) contacting said cell with a candidate compound and monitoring level of biological
15 activity of said IAP polypeptide or said NAIP polypeptide in said cell, a decrease in the level of biological activity of said IAP polypeptide or said NAIP polypeptide in said cell in response to said candidate compound relative to a cell not contacted with said candidate compound indicating the presence of a compound that enhances apoptosis in said affected cell that is proliferating in said proliferative disease.

20 43. The method of claim 42, wherein said cell further expresses a p53 polypeptide associated with said proliferative disease.

44. The method of claim 41 or 42, wherein said biological activity is the level of expression of said polypeptide; wherein said biological activity is the level of expression of an mRNA molecule encoding said polypeptide; or wherein said biological activity is an
25 apoptosis-inhibiting activity.

- 99 -

45. The method of claim 44, wherein said level of expression is measured by assaying the amount of said polypeptide present in said cell.

46. The method of claim 41 or 42, wherein said polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP.

5 47. The method of claim 41 or 42, wherein said polypeptide is NAIP.

48. The method of claim 41 or 42, wherein said polypeptide is XIAP.

49. The method of claim 41 or 42, wherein said polypeptide is HIAP-1.

50. The method of claim 41 or 42, wherein said polypeptide is HIAP-2.

51. A method for determining the prognosis of a mammal diagnosed with a
10 proliferative disease, said method comprising the steps of:

(a) isolating a sample from a tissue from said mammal; and

(b) determining whether said sample has an increased an level of biological activity of an IAP polypeptide or an NAIP polypeptide relative to a control sample, an increase in said level in said sample being an indication that said mammal has a poor prognosis.

15 52. The method of claim 51, wherein said sample comprises a cells that is proliferating in said proliferative disease and said control sample is from said tissue, said control sample consisting of healthy cells.

53. The method of claim 52, wherein said sample and said control sample are from said mammal.

20 54. The method of claim 51, wherein said sample further comprises a cell expressing a p53 polypeptide associated with said proliferative disease.

- 100 -

55. The method of claim 51, wherein said biological activity is the level of expression of said polypeptide; wherein said biological activity is the level of expression of an mRNA molecule encoding said polypeptide; or wherein said biological activity is an apoptosis-inhibiting activity.

5 56. The method of claim 55, wherein said level of expression is measured by assaying the amount of said polypeptide present in said cell.

57. The method of claim 51, wherein said polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP.

58. The method of claim 51, wherein said polypeptide is NAIP.

10 59. The method of claim 51, wherein said polypeptide is XIAP.

60. The method of claim 51, wherein said polypeptide is HIAP-1.

61. The method of claim 51, wherein said polypeptide is HIAP-2.

62. The method of claim 51, wherein said level is assayed by measuring the amount of IAP peptide of less than 64 kDa present in said sample.

15 63. A method for determining the prognosis of a mammal diagnosed with a proliferative disease, said method comprising the steps of:

(a) isolating a sample from said mammal, said sample having a nuclear fraction; and

(b) measuring the amount of a polypeptide that is recognized by an antibody that specifically binds an IAP polypeptide or an antibody that specifically binds an NAIP polypeptide in said nuclear fraction of said sample relative an amount from a control sample, an increase in said amount from said sample being an indication that said mammal has a poor prognosis.

20

- 101 -

64. The method of claim 63, wherein said sample is from a tissue of said mammal, said sample comprising a cell that is proliferating in said proliferative disease, and said control sample is from said tissue, said control sample consisting of healthy cells.

65. The method of claim 64, wherein said sample and said control sample are from
5 said mammal.

66. The method of claim 63, wherein said biological activity is the level of expression of said polypeptide; wherein said biological activity is the level of expression of an mRNA molecule encoding said polypeptide; or wherein said biological activity is an apoptosis-inhibiting activity.

10 67. The method of claim 66, wherein said level of expression is measured by assaying the amount of said polypeptide present in said cell.

68. The method of claim 63, wherein said polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP.

69. The method of claim 63, wherein said polypeptide is NAIP.

15 70. The method of claim 63, wherein said polypeptide is XIAP.

71. The method of claim 63, wherein said polypeptide is HIAP-1.

72. The method of claim 63, wherein said polypeptide is HIAP-2.

73. The method of claim 63, wherein said amount is measured by immunological methods.

20 74. A method for treating a mammal diagnosed as having a proliferative disease, said method comprising the steps of:

- 102 -

(a) measuring the amount of an IAP or NAIP polypeptide in a first sample from a tissue from said mammal, said first sample comprising a cell that is proliferating in said proliferative disease;

(b) measuring the amount of said polypeptide in a second sample from said tissue,
5 said second sample consisting of healthy cells;

(c) detecting an increase in the amount of said polypeptide in said first sample to the amount of said polypeptide in said second sample; and

(d) treating said mammal with a compound that decreases the biological activity of said polypeptide.

10 75. The method of claim 74, wherein said first sample and said second sample are from said mammal.

76. The method of claim 74, wherein said biological activity is the level of expression of said polypeptide; wherein said biological activity is the level of expression of an mRNA molecule encoding said polypeptide; or wherein said biological activity is an
15 apoptosis-inhibiting activity.

77. The method of claim 76, wherein said level of expression is measured by assaying the amount of said polypeptide present in said cell.

78. The method of claim 74, wherein said polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP.

20 79. The method of claim 74, wherein said polypeptide is NAIP.

80. The method of claim 74, wherein said polypeptide is XIAP.

81. The method of claim 74, wherein said polypeptide is HIAP-1.

82. The method of claim 74, wherein said polypeptide is HIAP-2.

- 103 -

83. Use of a compound that decreases the biological activity an IAP polypeptide or a NAIP polypeptide for the manufacture of a medicament for the enhancement of apoptosis.

84. The use of claim 83, wherein said biological activity is the level of expression of said polypeptide; wherein said biological activity is the level of expression of an mRNA molecule encoding said polypeptide; or wherein said biological activity is an apoptosis-inhibiting activity.

85. The use of claim 84, wherein said level of expression is measured by assaying the amount of said polypeptide present in said cell.

86. The use of claim 83, wherein said polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP.

87. The use of claim 83, wherein said polypeptide is NAIP.

88. The use of claim 83, wherein said polypeptide is XIAP.

89. The use of claim 83, wherein said polypeptide is HIAP-1.

90. The use of claim 83, wherein said polypeptide is HIAP-2.

91. A kit for diagnosing a mammal for the presence of a proliferative disease or an increased likelihood of developing a proliferative disease, said kit comprising an oligonucleotide that hybridizes to a nucleic acid sequence that encodes an IAP polypeptide or a NAIP polypeptide.

92. The kit of claim 91, wherein said polypeptide is selected from the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP.

93. The kit of claim 91, wherein said polypeptide is NAIP.

- 104 -

94. The kit of claim 91, wherein said polypeptide is XIAP.

95. The kit of claim 91, wherein said polypeptide is HIAP-1.

96. The kit of claim 91, wherein said polypeptide is HIAP-2.

97. A transgenic mammal, said mammal having an elevated level of biological
5 activity of an IAP polypeptide or a NAIP polypeptide.

98. The transgenic mammal of claim 97, wherein said biological activity is the level
of expression of said polypeptide; wherein said biological activity is the level of expression
of an mRNA molecule encoding said polypeptide; or wherein said biological activity is an
apoptosis-inhibiting activity.

10 99. The transgenic mammal of claim 98, wherein said level of expression is
measured by assaying the amount of said polypeptide present in said cell.

100. The transgenic mammal of claim 97, wherein said polypeptide is selected from
the group consisting of HIAP-1, m-HIAP-1, HIAP-2, m-HIAP-2, XIAP, and m-XIAP.

101. The transgenic mammal of claim 97, wherein said polypeptide is NAIP.

15 102. The transgenic mammal of claim 97, wherein said polypeptide is XIAP.

103. The transgenic mammal of claim 97, wherein said polypeptide is HIAP-1.

104. The transgenic mammal of claim 97, wherein said polypeptide is HIAP-2.

1/42

SEQ ID 3—1 GAAAAGGTGGACAAGTCCTATTTTCAAGAGAAGATGACTTTTAAACAGTTTGAAGGATCT 60
SEQ ID 4—1 M T F N S F E G S 9

61 AAAACTTGTGTACCTGCAGACATCAATAAGGAAGAAGATTTGTAGAAGAGTTTAATAGA 120
10 K T C V P A D I N K E E E F V E E F N R 29

121 TTAAAAACTTTTGCTAATTTTCCAAGTGGTAGTCCTGTTTCAGCATCAACACTGGCACGA 180
30 L K T F A N F P S G S P V S A S T L A R 49

181 GCAGGGTTTCTTTATACTGGTGAAGGAGATACCGTGCGGTGCTTTAGTTGTCATGCAGCT 240
50 A G F L Y T G E G D T V R C F S C H A A 69

241 GTAGATAGATGGCAATATGGAGACTCAGCAGTTGGAAGACACAGGAAAGTATCCCCAAAT 300
70 V D R W Q Y G D S A V G R H R K V S P N 89

301 TGCAGATTTATCAACGGCTTTTATCTTGAAAATAGTGCCACGCAGTCTACAAATTCTGGT 360
90 C R F I N G F Y L E N S A T Q S T N S G 109

361 ATCCAGAATGGTCAGTACAAAGTTGAAAACATCTGGAAGCAGAGATCATTTTGCCTTA 420
110 I Q N G Q Y K V E N Y L G S R D H F A L 129

421 GACAGGCCATCTGAGACACATGCAGACTATCTTTGAGAACTGGGCAGGTTGTAGATATA 480
130 D R P S E T H A D Y L L R T G Q V V D I 149

481 TCAGACACCATATACCCGAGGAACCTGCCATGTATaGTGAAGAAGCTAGATTAAAGTCC 540
150 S D T I Y P R N P A M Y S E E A R L K S 169

541 TTTCAGAACTGGCCAGACTATGCTCACCTAACCCCAAGAGAGTTAGCAAGTGCTGGACTC 600
170 F Q N W P D Y A H L T P R E L A S A G L 189

601 TACTACACAGGTATTGGTGACCAAGTGCAGTGCTTTTGTGTGGTGGAAAAC TGAAAAT 660
190 Y Y T G I G D Q V Q C F C C G G K L K N 209

661 TGGGAACCTTGTGATCGTGCCTGGTCAGAACACAGGCGACACTTTCCTAATTGCTTCTTT 720
210 W E P C D R A W S E H R R H F P N C F F 229

721 GTTTTGGGCCGAATCTTAATATTCCAAGTGAATCTGATGCTGTGAGTTCTGATAGGAAT 780
230 V L G R N L N I R S E S D A V S S D R N 249

781 TTCCCAAATCAACAAATCTTCCAAGAAATCCATCCATGGCAGATTATGAAGCACGGATC 840
250 F P N S T N L P R N P S M A D Y E A R I 269

841 TTTACTTTTGGGACATGGATATACTCAGTTAACAAGGAGCAGCTTGCAAGAGCTGGATTT 900
270 F T F G T W I Y S V N K E Q L A R A G F 289

901 TATGCTTTAG^{1,2}GTGAAGGTGATAAAGTAAAGTGCTTTCACTGTGGAGGAGGGCTAACTGAT 960
290 Y A L G E G D K V K C F H C G G G L T D 309

961 TGGAAGCCCACTGAAGACCTTGGGAACAACATGCTAAATGGTATCCAG^{2,3}GTGCAATAT 1020
310 W K P S E D P W E Q H A K W Y P G C K Y 329

1021 CTGTTAGAACAGAAGGGACAAGAATATATAAACAATATTCAATTTAACTCATTCACTTGAG 1080
330 L L E Q K G Q E Y I N N I H L T H S L E 349

Fig. 1

SUBSTITUTE SHEET (RULE 26)

Fig 1
Sheet 2 of 3

3/42

3061 AATATTGGCAAGAAAAGAAGAAATAGTTGTTTAAATATTTTTTAAAAAACACTTGAATAAG 3120
3121 AATCAGTAGGGTATAAACTAGAAGTTTAAAAATGCCTCATAGAACGTCCAGGGTTTACAT 3180
3181 TACAAGATTCTCACAACAAACCCATTGTAGAGGTGAGTAAGGCATGTTACTACAGAGGAA 3240
3241 AGTTTGACAGTAAACTGTAAAAAATTATATTTTTGTTGTACTTTCTAAGAGAAAGAGTA 3300
3301 TTGTTATGTTCTCCTAACTTCTGTTGATTACTACTTTAAGTGATATTCATTTAAAAACATT 3360
3361 GCAAATTTATTTTATTTATTTAATTTTCTTTTGGAGATGGAGTCTTGCTTGTCACCCAGG 3420
3421 CTGGAGTGCAGTGGAGTGATCTCTGCTCACTGCAACCTCCGCCTTCTGGGTTCAGCGAT 3480
3481 TCTCGTGCTCAGCTTCCTGAGTAGCTGGAATTACAGGCAGGTGCCACCATGCCCGACTA 3540
3541 ATTTTTTTTTTATTTTAGTAGAGACGGGGTTTCACCATGTTGGCCAGGCTGGTATCAAAC 3600
3601 TCCTGACCTCAAGAGATCCACTCGCCTTGCCCTCCCAAAGTGCTGGGATTACAGGCTTGA 3660
3661 GCCACCACGCCCCGGCTAAACATTGCAAATTTAAATGAGAGTTTTAAAAATTAAATAATG 3720
3721 ACTGCCCTGTTTCTGTTTATGATGTAAATCCTCAGTTCTTCACCTTTCAGTGTCTGCC 3780
3781 ACTTAGTTTGGTTATATAGTCATTAACCTGAATTTGGTCTGTATAGTCTAGACTTTAAAT 3840
3841 TTAAAGTTTCTACAAGGGGAGAAAAGTGTTAAATTTTTTAAATATGTTTCCAGGACA 3900
3901 CTTCACTTCCAAGTCAGGTAGGTAGTTCAATCTAGTTGTTAGCCAAGGACTCAAGGACTG 3960
3961 AATTGTTTTAACATAAGGCTTTTCTGTTCTGGGAGCCGCACTTCATTAATAATCTTCTTA 4020
4021 AAACCTGTATGTTTAGAGTTAAGCAAGACTTTTTTCTTCTCTCCATGAGTTGTGAAAT 4080
4081 TTAATGCACAACGCTGATGTGGCTAACAAGTTTATTTTAAAGAAATGTTTAGAAATGCTGT 4140
4141 TGCTTCAGGTTCTTAAATCACTCAGCACTCCAACCTTCAATCAAATTTTGGAGACTTA 4200
4201 ACAGCATTTGTCTGTGTTTGAACATAAAAAGCACCGGATCTTTCCATCTAATTCGCCA 4260
4261 AAAATTGATCATTTGCAAAGTCAAAACTATAGCCATATCCAAATCTTTCCCCCTCCCAA 4320
4321 GAGTTCTCAGTGTCTACATGTAGACTATTCCTTTTCTGTATAAAGTTCACTCTAGGATTT 4380
4381 CAAGTCACCACTTATTTTACATTTTAGTCATGCAAAGATTCAAGTAGTTTTCGAATAAGT 4440
4441 ACTTATCTTTATTTGAATAATTTAGTCTGCTGATCAAAAGCATTTGTCTTAATTTTGTAG 4500
4501 AACTGGTTTTAGCATTTACAACTAAATTCAGTTAATTAATTAATAGCTTTATATTGCC 4560
4561 TTTCTCTGTACATTTGGTTTTTTCCCTGTCCCTTTGATTACGGGCTAAGGTAGGGTAAG 4620
4621 AXXGGGTGTAGTGAGTGATATATAATGTGATTTGGCCCTGTGTATTATGATATTTGTTAT 4680
4681 TTTTGTGTATATTTATTTACATTTTCAGTAGTTGTTTTTGTGTTTCCATTTTAGGGGAT 4740
4741 AAAATTTGTATTTTGAACATGAATGGAGACTACGCCCCAGCATTAGTTTTCACATGATA 4800
4801 TACCCTTTAAACCCGAATCATTGTTTTATTTCCCTGATTACACAGGTGTTGAATGGGGAAA 4860
4861 GGGGCTAGTATATCAGTAGGATATACTATGGGATGTATATATATCATTGCTGTAGAGAA 4920
4921 ATGAAATAAAATGGGGCTGGGCTCAGTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGG 4980
4981 CTGAGGCAGGTGGATCACGAGGTCAGGAGATCGAGACCATCCTGGCTAACACGCTGAAAC 5040
5041 CCCGTCTCTACTAAAAACAGAAAATTAGCCGGGCGTGGTGGCGGGCGCCTGTAGTCCCA 5100
5101 GCTACTCGGGAGGCTGAGGCAGGAGAATGGTGTGAACCCGGGAGGCAGAGCTTGCACTGA 5160
5161 GCCGAGATCTCGCCACTGCACTCCAGCCTGGGCAACAGAGCAAGACTCTGTCTCAAAAAA 5220
5221 AAAAAAAAAAAG 5232

Fig. 1 (cont.)

SUBSTITUTE SHEET (RULE 26)

4/42

SEQ. ID 5—1 TTGCTCTGTCACCCAGTTTGGAGTGCAGTTATGCAGTCTCACACTGCAAGCTCTGCCTCA 60
61 TGGGCTCAAGTGAACCTCCTGCCTCAGCCTCTCAAGTAGCTGGGACCACAGGCAGGTGCC 120
121 ACCATGTCTGGCTAATTTTGTAGTTCTTTGTAGAGATGGTGTCTTTGCCAAGTCACCCAG 180
181 TTTGAGGCTGGTCTCAAACACCTGGGCTCAAGCAATCCATCTACCTCAGCCTCCCAAAGT 240
241 GCTGGGATTACAGGAGTGAGCCATGGCATGAGGCCTTGTGGGGTGTCTCTTTTAAATGAA 300
301 AGCATACTCTGTTTACGTATTTGATATGAAGGAATATCCTTCCTTTCCACAAAGACAAAA 360
361 ATTATCCTATTTTCTCAAAACATATGTCCTTTTCTCTACTTTTCATTTTGTACTTT 420
421 TGATGGACACATGTGTTACATTGATTTCACTTTCTCATAATTCTGCTGTAAGAAAAACAA 480
481 TAGTGCCAGTTCAATGACAAATAGCAACAGTCTGTTATTGCTAGACTGTTACTGTTAGTG 540
541 GAGACTACCAGAACAGTCACTCCCAGTGTCAAGGAATCAAAGAGAACATGTTCCCTCTCT 600
601 AAAGGGCACAGCTGCTGCTCAGCTTTAGCTGATTGCTGCCCTGCAGGACTATAGGCCAG 660
661 TGTGCTAGATCTTTTGATGTTTCAAGAGAAGCTTGAATCTAGAATGTGATGGGAAGTC 720
721 TCTTACATTTAAACATGTTGGCAATTAATGGTAAGATTTAAAAATACTGTGGTCCAAGAA 780
781 AAAAATGGATTGGAAACTGGATTAAATTCAAATGAGGCATGCAGATTAATCTACAGCAT 840
841 GGTACAATGTGAATTTTCTGGTTTCTTTAATTGCACTGTAAATAGGTAAGATGTTAGCTT 900
901 TGGGGAAGCTAAGTGCAGAGTATGCAGAACTATATTTTTGTAAAGTTTTCTCTAAGTAT 960
961 AAATAAATTTCAAATAAAAAATAAAAACTTAGTAAAGAACTATAATGCAATTCATGTAA 1020
1021 GCCAAACATAATATGCTTCCAGTTTGAAACCTCTGGGTTTTATTTTATTTTATTTTATT 1080
1081 TTTGAGACAGAGTCTTGCTGTGTCAACCAGGCTGGAGTGTAGTGGCACTATTTCCGCCCA 1140
1141 CTGCAACCTCCACCTCCAGGCTCAAATGATTCTCCTGCCTCAGCCTCCGGAGTAGCTGG 1200
1201 GATTACAGGCGGTACCACCACACCCAGCTAATTTTGTATTTTGTAGTAGAGATGGGGTT 1260
1261 TCACCATTTTGGCCAGGCTGGTTTTGAACTCCTGACCTCAAGTGATCCACTTGTCTTGGC 1320
1321 CTCCAAAATGCTGGGATTACAGGCGTGAGCCACTGCACCAGGCAGAGGCCTCTGTTTTT 1380
1381 TATCTCTTTTTTGGCCTCTACAGTGCCTAGTAAAGCACCTGATACATGGTAAACGATCAGT 1440
1441 AATTACTAGTACTCTATTTTGGAGAAAATGATTTTTTAAAAAGTCATTGTGTTCCATCCA 1500
1501 TGAGTCGTTTGTAGTTTTTAAACTGTCTTTTTGTGTTGTTTTGAACAGGTTTACAAAGGAG 1560
1561 GAAAACGACTTCTCTAGATTTTTTTTTTTCAGTTTCTCTATAAATCAAACATCTCAAAA 1620
1621 TGGAGACCTAAATCCTTAAAGGGACTTAGTCTAATCTCGGGAGGTAGTTTGTGCATGG 1680
1681 GTAAACAAATTAAGTATTAAGTATGCTGTTTACTATCCAAAGAAATGCTAATTTTATAAACA 1740
1741 TGATCAGGTTATATAAGGTATACCATATGAGTTGAGTTTGATTTTGAATTTGATTGTTGAAAT 1800
1801 AAAGGAAAAGTGATTCTAGCTGGGGCATATTGTTTAAAGCATTTTTTTCAGAGTTGGCCAG 1860
1861 GCAGTCTCCTACTGGCACATTCTCCCATATGTAGAATAGAAATAGTACCTGTGTTTGGG 1920
1921 AAAGATTTTAAATGAGTGACAGTTATTTGGAACAAAGAGCTAATAATCAATCCACTGCA 1980
1981 AATTAAAGAAACATGCAGATGAAAGTTTGCACATTAAATACTTCTACAGTGACAAAG 2040
2041 AAAAATCAAGAACAAAGCTTTTTGATATGTGCAACAAATTTAGAGGAAGTAAAAAGATAA 2100
2101 ATGTGATGATTGGTCAAGAAATATCCAGTTATTTACAAGGCCACTGATATTTTAAACCT 2160
2161 CCAAAGTTTGTTTAAATGGGCTGTTACCGCTGAGAATGATGAGGATGAGAATGATGGTT 2220
2221 GAAGGTTACATTTTAGGAAATGAAGAACTTAGAAAATTAATAAAGACAGTGATGAAT 2280
2281 ACAAGAAGATTTTATAACAATGTGTAAAATTTTTGGCCAGGGAAAGGAATATTGAAGT 2340
2341 TAGATACAATTACTTACCTTTGAGGGAATAATTGTTGGTAATGAGATGTGATGTTTCTC 2400
2401 CTGCCACCTGGAAACAAAGCATTGAAGTCTGCAGTTGAAAAGCCCAACGTCTGTGAGATC 2460
2461 CAGGAAACCATGCTTGCAAAACCACTGGTAAAAAAAAAAAAAAAAAAAAAAAAAAGCCACAG 2520
2521 TGACTTGCTTATTTGGTCAATGCTAGTATTATCGACTCAGAACCCTTTTACTAATGGCTAG 2580
2581 TAAATCATAATTGAGAAATCTGAATTTTGACAAGGTCTCTGCTGTTGAAATGGTAAATT 2640
2641 TATTATTTTTTTTGTGATGATAAATCTGGTTCAAGGTATGCTATCCATGAAATAATTTT 2700
2701 TGACCAAAACTAAATTGATGCAATTTGATTATCCATCTTAGCCTACAGATGGCATCTGGT 2760
2761 AACTTTTGACTGTTTTAAAAATAAATCCACTATCAGAGTAGATTGATGTTGGCTTCAG 2820
2821 AAACATTTAGAAAAACAAAAGTTCAAAAATGTTTTTCAGGAGGTGATAAGTTGAATAACTC 2880
2881 TACAATGTTAGTTCTTTGAGGGGGACARAAAATTTAAATCTTTGAAAGGTCTTATTTTA 2940
2941 CAGCCATATCTAAATTATCTTAAGAAAATTTTAAACAAAGGGAATGAAATATATATCATG 3000
3001 ATTCTGTTTTTCCAAAAGTAACCTGAATATAGCAATGAAGTTCAGTTTGTATTGGTAG 3060
3061 TTTGGGCAGAGTCTCTTTTGCAGCAGCTGTTGTCTACCATAATTACAGAGGACATTTCC 3120
3121 ATGTTCTAGCCAAGTATACTATTAGAATAAARAACTTAACATTGAGTTGCTTCAACAGC 3180

Fig. 2

SUBSTITUTE SHEET (RULE 26)

5/42

3181 ATGAAACTGAGTCCAAAAGACCAAATGAACAAACACATTAATCTCTGATTATTTATTTTA 3240
 3241 AATAGAAATATTTAATTGTGTGAAGATCTAATAGTATCATTATACCTTAAGCAATCATATTCC 3300
 3301 TGATGATCTATGGGAAATAACTATTATTTAATTAATATTGAAACCAGGTTTTAAGATGTG 3360
 3361 TTAGCCAGTCTGTACTAGTAAATCTCTTTATTTGGAGAGAAATTTAGATTGTTTTGT 3420
 3421 TCTCCTTATTAGAAGGATTGTAGAAAGAAAAAATGACTAATTGGAGAAAAATGGGGAT 3480
 3481 ATATCATATTTCACTGAATTCAAAATGTCTTCAGTTGTAAATCTTACCATTATTTTACGT 3540
 3541 ACCTCTAAGAAATAAAAGTGCTTCTAATTAAAAATATGATGTCATTAATTATGAAATACTT 3600
 3601 CTTGATAACAGAAGTTTTAAAAATAGCCATCTTAGAATCAGTGAAATATGGTAATGTATTA 3660
 3661 TTTTCCTCCTTTGAGTNAGGTCTTGTGCTTTTNTTCTGGCCACTAAATNTCCACCATNT 3720
 3721 CCAANAAGCAAANTAAACCTATTCTGAATATTTTGTGTGAAACACTTGNCAGCAGAGC 3780
 3781 TTTCCCNCCATGNNAGAAGCTTCATGAGTCACACATTACATCTTTGGGTTGATTGAATGC 3840
 3841 CACTGAAACATTTCTAGTAGCCTGGAGNAGTTGACCTACCTGTGGAGATGCCTGCCATTA 3900
 3901 AATGGCATCCTGATGGCTTAATACACATCACTCTTCTGTGNAGGGTTTTAATTTTCAACA 3960
 3961 CAGCTTACTCTGTAGCATCATGTTTACATTGTATGTATAAAGATTATACNAAGGTGCAAT 4020
 4021 TGTGTATTTCTTCCTTAAATGTATCAGTATAGGATTTAGAATCTCCATGTTGAAACTCT 4080
 4081 AAATGCATAGAAATAAAAAATAAAAAAATTTTCATTTTGGCTTTTCAGCCTAGTATTA 4140
 4141 AAATGATAAAAGCAAAGCCATGCACAAACTACCTCCCTAGAGAAAGGCTAGTCCCTTT 4200
 4201 TCTTCCCATTCATTTCATTATGAACATAGTAGAAACAGCATATCTTATCAAATTTGA 4260
 SEQ. ID 6-1 M N I V E N S I F L S N L M 14

4261 TGAAAAGCGCCAACACGTTTGAAGTGAATACGACTTGTCATGTGAAGTGTACCGAATGT 4320
 15 K S A N T F E L K Y D L S C E L Y R M S 34

4321 CTACGTATTCCACTTTTCTGCTGGGGTTCTGTCTCAGAAAGGAGTCTTGCTCGTGCTG 4380
 35 T Y S T F P A G V P V S E R S L A R A G 54

4381 GTTCTATTACACTGGTGTGAATGACAAGGTCAAATGCTTCTGTTGTGGCCTGATGCTGG 4440
 55 F Y Y T G V N D K V K C F C C G L M L D 74

4441 ATAAGTGGAAAAGAGGAGACAGTCTACTGAAAAGCATAAAAAGTTGTATCCTAGCTGCA 4500
 75 N W K R G D S P T E K H K K L Y P S C R 94

4501 GATTCGTTTCAAGTCTAAATTCGGTTAACAACCTTGAAGCTACCTCTCAGCCTACTTTTC 4560
 95 F V Q S L N S V N N L E A T S Q P T F P 114

4561 CTTCTTCAGTAACACATTCCACACACTCATTACTCCGGGTACAGAAAACAGTGGATATT 4620
 115 S S V T H S T H S L L P G T E N S G Y F 134

4621 TCCGTGGCTCTTATTCAAACCTCTCCATCAAATCCTGTAAACTCCAGAGCAAATCAAGAAT 4680
 135 R G S Y S N S P S N P V N S R A N Q E F 154

4681 TTTCTGCCTTGATGAGAAGTTCCTACCCCTGTCCAATGAATAACGAAAATGCCAGATTAC 4740
 155 S A L M R S S Y P C P M N N E N A R L L 174

4741 TTACTTTTCAGACATGGCCATTGACTTTTCTGTGCGCAACAGATCTGGCACGAGCAGGCT 4800
 175 T F Q T W P L T F L S P T D L A R A G F 194

4801 TTTACTACATAGGACCTGGAGACAGAGTGGCTTGCTTTGCCTGTGGTGGAAAATTGAGCA 4860
 195 Y Y I G P G D R V A C F A C G G K L S N 214

4861 ATTGGGAACCGAAGGATAATGCTATGTGAGAACACCTGAGACATTTTCCCAAATGCCCAT 4920
 215 W E P K D N A M S E H L R H F P K C P F 234

Fig. 2 (cont.)

SUBSTITUTE SHEET (RULE 26)

6/42

4921 TTATAGAAAATCAGCTTCAAGACACTTCAAGATACACAGTTTCTAATCTGAGCATGCAGA 4980
235 I E N Q L Q D T S R Y T V S N L S M Q T 254

4981 CACATGCAGCCCGCTTTAAACATTCTTTAACTGGCCCTCTAGTGTCTAGTTAATCCTG 5040
255 H A A R F K T F F N W P S S V L V N P E 274

5041 AGCAGCTTGCAAGTGC GG G T T T T T A T T A T G T G G T A A C A G T G A T G T C A A A T G C T T T T 5100
275 Q L A S A G F Y Y V G N S D D V K C F C 294

5101 GCTGTGATGGTGGACTCAGGTGTTGGGAATCTGGAGATGATCCATGGGTTCAACATGCCA 5160
295 C D G G L R C W E S G D D P W V Q H A K 314

5161 AGTGGTTTCCAAGGTGTGAGTACTTGATAAGAATTAAGGACAGGAGTTCATCCGTCAAG 5220
315 W F P R C E Y L I R I K G Q E F I R Q V 334

5221 TTCAAGCCAGTTACCCTCATCTACTTGAACAGCTGCTATCCACATCAGACAGCCCAGGAG 5280
335 Q A S Y P H L L E Q L L S T S D S P G D 354

5281 ATGAAAATGCAGAGTCATCAATTATCCATTTTGAACCTGGAGAAGACCATTTCAGAAGATG 5340
355 E N A E S S I I H F E P G E D H S E D A 374

5341 CAATCATGATGAATACTCCTGTGATTAATGCTGCCGTGGAAATGGGCTTTAGTAGAAGCC 5400
375 I M M N T P V I N A A V E M G F S R S L 394

5401 TGGTAAACAGACAGTTCAGAGAAAAATCCTAGCAACTGGAGAGAATTATAGACTAGTCA 5460
395 V K Q T V Q R K I L A T G E N Y R L V N 414

5461 ATGATCTTGTGTTAGACTTACTCAATGCAGAAGATGAAATAAGGGAAGAGGAGAGAGAAA 5520
415 D L V L D L L N A E D E I R E E E R E R 434

5521 GAGCAACTGAGGAAAAAGAAATCAATGATTTATTATTAATCCGGAAGAATAGAATGGCAC 5580
435 A T E E K E S N D L L L I R K N R M A L 454

5581 TTTTTCACATTTGACTTGTGTAATCCAATCCTGGATAGTCTACTAACTGCCGGAATTA 5640
455 F Q H L T C V I P I L D S L L T A G I I 474

5641 TTAATGAACAAGAACATGATGTTATTAAACAGAAGACACAGACGTCTTTACAAGCAAGAG 5700
475 N E Q E H D V I K Q K T Q T S L Q A R E 494

5701 AACTGATTGATACGATTTTAGTAAAAGGAAATATTGCAGCCACTGTATTCAGAACTCTC 5760
495 L I D T I L V K G N I A A T V F R N S L 514

5761 TGCAAGAAGCTGAAGCTGTGTTATATGAGCATTATTTGTGCAACAGGACATAAAATATA 5820
515 Q E A E A V L Y E H L F V Q Q D I K Y I 534

5821 TTCCACAGAAGATGTTTCAGATCTACCAGTGAAGAACAATTGCGGAGACTACAAGAAG 5880
535 P T E D V S D L P V E E Q L R R L Q E E 554

5881 AAAGAACATGTAAAGTGTGTATGGACAAAGAAGTGTCATAGTGTGTTTATCCTTGTGGTC 5940
555 R T C K V C M D K E V S I V F I P C G H 574

5941 ATCTAGTAGTATGCAAGATTGTGCTCCTTCTTTAAGAAAGTGTCTTATTTGTAGGACTA 6000
575 L V V C K D C A P S L R K C P I C R S T 594

Fig. 2 (cont.)

SUBSTITUTE SHEET (RULE 26)

7/42

6001 CAATCAAGGGTACAGTTCGTACATTTCTTTTCATGAAGAAGAACCAAAACATCGTCTAAAC 6060
595 I K G T V R T F L S * 604

6061 TTTAGAATTAATTTATTAAATGTATTATAACTTTAACTTTTATCCTAATTTGGTTTCCTT 6120
6121 AAAATTTTTATTTATTTTACAACCTCAAAAAACATTGTTTTGTGTAACATATTTATATATGT 6180
6181 ATCTAAACCATATGAACATATATTTTTTTAGAACTAAGAGAATGATAGGCTTTTGTCTT 6240
6241 ATGAACGAAAAAGAGGTAGCACTACAAACACAATATTCAATCAAAATTTTCAGCATTATTG 6300
6301 AAATTGTAAGTGAAGTAAACTTAAGATATTTGAGTTAACCTTTAAGAATTTTAAATATT 6360
6361 TTGGCATTGTACTAATACCGGAACATGAAGCCAGGTGTGGTGGTATGTGCCTGTAGTCC 6420
6421 CAGGCTGAGGCAAGAGAATTACTTGAGCCCAGGAGTTTGAATCCATCCTGGGCAGCATAC 6480
6481 TGAGACCCTGCCTTTAAAAACAAACAGAACAAAAACAAACACCAGGGACACATTTCTCT 6540
6541 GTCTTTTTTGATCAGTGTCTTATACATCGAAGGTGTGCATATATGTTGAATCACATTTTA 6600
6601 GGGACATGGTGTTTTTTATAAAGAATTCTGTGAGAAAAAATTTAATAAAGCAACCAAAAA 6660
6661 AAAAAAAA 6669

Fig. 2 (cont.)

8/42

SEQ. ID 7— 1 GAGCGCCCGGG^{1,2}CTGATCCGAGCCGAGCGGGCCGTATCTCCTTGTCGGCGCCGCTGATTCC 60
 61 CGGCTCTGCGGAGGCCTCTAGGCAGCCGCGCAGCTTCCGTGTTTGCTGCGCCCGCACTGC 120
 121 ^{2,3}GATTTACAACCCTGAAGAATCTCCCTATCCCTATTTTGTCCCCCTGCAGTAATAAATCCC 180
 181 ATTATGGAGATCTCGAACTTTATAAAGGGATATAGTTGAATTCTATGGAGTGTAATTT 240
 241 TGTGTATGAATTATATTTTAAACATTGAAGAGTTTTCAGAAAGAAGGCTAGTAGAGTT 300
 301 GATTACTGATACTTTATGCTAAGCAGTACTTTTTTGGTAGTACAATATTTTGTAGGCCGT 360
 361 TTCTGATAACACTAGAAAGGACAAGTTTATCTTGTGATAAATTGATTAATGTTTACAAC 420
 421 ATGACTGATAATTATAGCTGAATAGTCCTTAAATGATGAACAGGTTATTTAGTTTTTAAA 480
 481 TGCAGTGTA AAAAGTGTGCTGTGGAATTTTATGGCTAACTAAGTTTATGGAGAAAATAC 540
 541 CTTCAAGTTGATCAAGAATAATAGTGGTATACAAAGTTAGGAAGAAAGTCAACATGATGCT 600
 601 GCAGGAAATGGAACAAATACAAATGATATTTAACAAGATAGAGTTTACAGTTTTTGAA 660
 661 CTTTAAGCCAAATTCATTTGACATCAAGCACTATAGCAGGCACAGGTTCAACAAAGCTTG 720
 721 TGGGTATTGACTTCCCCCAAAGTTGTCAGCTGAAGTAATTTAGCCCACTTAAGTAAATA 780
 781 CTATGATGATAAGCTGTGTGAACCTAGCTTTTAAATAGTGTGACCATATGAAGGTTTTAA 840
 841 TTACTTTTGTATTGGAATAAAATGAGATTTTTTGGGTGTCATGTTAAAGTGCTTATA 900
 901 GGGAAAGAAGCCTGCATATAATTTTTTACCTTGTGGCATAATCAGTAATGGTCTGTTAT 960
 961 TCAGGCTTCATAGCTTGTAAACARATATAAATAAAGGCATAATTTAGGTATTCTATAGT 1020
 1021 TGCTTAGAATTTTGTTAATATAAATCTCTGTGAAAAATCAAGGAGTTTAAATATTTTTCAG 1080
 1081 AAGTGCATCCACCTTTTCAAGGGCTTTAAGTTAGTATTAAGTCAAGATTATGAACAAATAGC 1140
 1141 ACTTAGGTTACCTGAAAGAGTTACTACAACCCCAAAGAGTTGTGTTCTAAGTAGTATCTT 1200
 1201 GGTAATTCAGAGAGATACTCATCTACCTGAATATAAACTGAGATAAATCCAGTAAAGAA 1260
 1261 AGTGTAGTAAATTCTACATAAGAGTCTATCATGATTTCTTTTTGTGGTAAAAATCTTAG 1320
 1321 TTCATGTGAAGAAATTTTATGTGAATGTTTTAGCTATCAAACAGTACTGTACCTACTCA 1380

M 1

SEQ. ID 8— 2 TGCACAAAACCTGCCTCCCAAAGACTTTTCCCAGGTCCCTCGTATCAAAACATTAAGAGTA 1440
 H K T A S Q R L F P G P S Y Q N I K S I 21
 1441 TAATGGAAGATAGCACGATCTTGTGAGATTGGACAAACAGCAACAAACAAAAATGAAGT 1500
 22 M E D S T I L S D W T N S N K Q K M K Y 41
 1501 ATGACTTTTCCTGTGAACCTACAGAATGTCTACATATTCAACTTTCCCCCGCGGGGTGC 1560
 42 D F S C E L Y R M S T Y S T F P A G V P 61
 1561 CTGTCTCAGAAAGGAGTCTTGTCTCGTGTGGTTTTTATTATACTGGTGTGAATGACAAGG 1620
 62 V S E R S L A R A G F Y Y T G V N D K V 81
 1621 TCAATGCTTCTGTTGTGGCCTGATGCTGGATAACTGGAACTAGGAGACAGTCCTATTC 1680
 82 K C F C C G L M L D N W K L G D S P I Q 101
 1681 AAAAGCATAAACAGCTATATCCTAGCTGTAGCTTTTATTCAGAATCTGGTTTCAGCTAGTC 1740
 102 K H K Q L Y P S C S F I Q N L V S A S L 121
 1741 TGGGATCCACCTCTAAGAATACGTCTCCAATGAGAAACAGTTTTCACATTCAATTATCTC 1800
 122 G S T S K N T S P M R N S F A H S L S P 141
 1801 CCACCTTGAACATAGTAGCTTGTTCAGTGGTCTTACTCCAGCCTTTCTCCAAACCCTC 1860
 142 T L E H S S L F S G S Y S S L S P N P L 161
 1861 TTAATTCTAGAGCAGTTGAAGACATCTCTTCATCGAGGACTAACCCCTACAGTTATGCAA 1920
 162 N S R A V E D I S S S R T N P Y S Y A M 181
 1921 TGAGTACTGAAGAAGCCAGATTTCTTACCTACCATATGTGGCCATTAAGTTTGTGTCAC 1980
 182 S T E E A R F L T Y H M W P L T F L S P 201

Fig. 3

SUBSTITUTE SHEET (RULE 26)

9/42

1981 CATCAGAATTGGCAAGAGCTGGTTTTTATTATATAGGACCTGGAGATAGGGTAGCCTGCT 2040
202 S E L A R A G F Y Y I G P G D R V A C F 221

2041 TTGCCTGTGGTGGGAAGCTCAGTAAGCTGGGAACCAAAGGATGATGCTATGTCAGAACACC 2100
222 A C G G K L S N W E P K D D A M S E H R 241

2101 GGAGGCATTTTCCCAACTGTCCATTTTTGAAAAATTCTCTAGAACTCTGAGGTTTAGCA 2160
242 R H F P N C P F L E N S L E T L R F S I 261

2161 TTTCAAATCTGAGCATGCAGACACATGCAGCTCGAATGAGAACATTTATGTACTGGCCAT 2220
262 S N L S M Q T H A A R M R T F M Y W P S 281

2221 CTAGTGTTCAGTTTCAGCCTGAGCAGCTTGCAAGTGCTGGTTTTTATTATGTGGGTCGCA 2280
282 S V P V Q P E Q L A S A G F Y Y V G R N 301

2281 ATGATGATGTCAAATGCTTTTGTGTGATGGTGGCTTGAGGTGTTGGGAATCTGGAGATG 2340
302 D D V K C F C C D G G L R C W E S G D D 321

2341 ATCCATGGGTAGAACATGCCAAGTGGTTTCCAAGTGTGAGTTCTTGATACGAATGAAAG 2400
322 P W V E H A K W F P R C E F L I R M K G 341

2401 GCCAAGAGTTTGTGATGAGATTCAAGGTAGATATCCTCATCTTCTTGAACAGCTGTTGT 2460
342 Q E F V D E I Q G R Y P H L L E Q L L S 361

2461 CAACTTCAGATACCACTGGAGAAGAAAATGCTGACCCACCAATTATTCATTTTGGACCTG 2520
362 T S D T T G E E N A D P P I I H F G P G 381

2521 GAGAAAGTTCTTCAGAAGATGCTGTGATGAATACACCTGTGGTTAAATCTGCCTTGG 2580
382 E S S S E D A V M M N T P V V K S A L E 401

2581 AAATGGGCTTTAATAGAGACCTGGTGAAACAAACAGTTCAAAGTAAATCCTGACAACTG 2640
402 M G F N R D L V K Q T V Q S K I L T T G 421

2641 GAGAGAACTATAAAACAGTTAATGATATTGTGTGACACTTCTTAATGCTGAAGATGAAA 2700
422 E N Y K T V N D I V S A L L N A E D E K 441

2701 AAAGAGAAGAGGAGAAGGAAAAACAAGCTGAAGAAATGGCATCAGATGATTTGTCTATTAA 2760
442 R E E E K E K Q A E E M A S D D L S L I 461

2761 TTCGGAAGAACAGAATGGCTCTCTTTCAACAATTGACATGTGTGCTTCCTATCCTGGATA 2820
462 R K N R M A L F Q Q L T C V L P I L D N 481

2821 ATCTTTTAAAGCCAATGTAATTAATAACAGGAACATGATATTATTAAACAAAAAACAC 2880
482 L L K A N V I N K Q E H D I I K Q K T Q 501

2881 AGATACCTTTACAAGCGAGAGAACTGATTGATACCATTTTGGTTAAAGGAAATGCTGCGG 2940
502 I P L Q A R E L I D T I L V K G N A A A 521

2941 CCAACATCTTCAAAAAGTGTCTAAAGAAATTGACTCTACATTGTATAAGAACTTATTG 3000
522 N I F K N C L K E I D S T L Y K N L F V 541

3001 TGGATAAGAATATGAAGTATATCCCAACAGAAGATGTTTCAGGTCTGTCACTGGAAGAAC 3060
542 D K N M K Y I P T E D V S G L S L E E Q 561

Fig. 3 (cont.)

SUBSTITUTE SHEET (RULE 26)

10/42

3061 AATTGAGGAGGTTGCAAGAAGAACGAACTTGTAAAGTGTGTATGGACAAAGAAGTTTCTG 3120
562 L R R L Q E E R T C K V C M D K E V S V 581

3121 TTGTATTTATTCCTTGTGGTCATCTGGTAGTATGCCAGGAATGTGCCCTTCTCTAAGAA 3180
582 V F I P C G H L V V C Q E C A P S L R K 601

3181 AATGCCCTATTTGCAGGGGTATAATCAAGGGTACTGTTCTGACATTTCTCTCTTAAAGAA 3240
602 C P I C R G I I K G T V R T F L S * 618

3241 AAATAGTCTATATTTTAACCTGCATAAAAAGGTCTTTAAAAATATTGTTGAACACTTGAAG 3300
3301 CCATCTAAAGTAAAAAGGAATTATGAGTTTTTCAATTAGTAACATTCATGTTCTAGTCT 3360
3361 GCTTTGGTACTAATAATCTTGTCTTCTGAAAAGATGGTATCATATATTTAATCTTAATCTG 3420
3421 TTTATTTACAAGGGAAGATTTATGTTTGGTGAACATATTAGTATGTATGTGTACCTAAG 3480
3481 GGAGTAGTGTCACTGCTTGTATGCATCATTTTCAGGAGTTACTGGATTGTTGTTCTTTC 3540
3541 AGAAAGCTTTGAATACTAAATTATAGTGTAGAAAAGAACTGGAAACCAGGAACCTCTGGAG 3600
3601 TTCATCAGAGTTATGGTGCCGAATTGTCTTTGGTGCTTTTCACTTGTGTTTTAAATAAG 3660
3661 GATTTTTCTCTTATTTCTCCCCCTAGTTTGTGAGAAACATCTCAATAAAGTGCTTTAAAA 3720
3721 AGAAAAAAAAA 3732

Fig. 3 (cont.)

11/42

SEQ. ID 9—1 ATTTTAAATTGATGCATTAACATTCTAAACATTATCTGTTTTAAATAGTAAAAATT 60
61 GAACTTTGCCTTGAATATGTAATGATTCAATFATAACAATTATGCATAGTCTTTAATAATC 120
121 TGCATATTTTATGCTGCTTTCATGTTTTTCCTAATTAATGACTTCACATGTTTAAATATT 180
181 ATAATTTTCTGTCATAGTTTCCATATTTATATAAAATGAATACTTAAGATCAGTAATTC 240
241 TGCTCTGTTTGTATATACTATTTCCATCAAAAGACAAAATGGGACTGAGGTTGAGGC 300
301 TCGTTGCTAAAGCACTTTCCTAAAATGCAAAAGGCCCTATGATGGATCCCTAGTACTTAT 360
361 TTAAGTGAGAGAGAAACAGGCTGGGGGTGTAGGTCTGTTAGAGCATGTGTTTGGCATTAT 420
421 GTGAAGCCCAACACTAAAAAAGGAGAACAAACAAAAGCGCAGACTTTAAACTCAAGTG 480
481 GTTTGGTAATGTACGACTCTACTGTTTGAATTTAAATGTGCTTAGTTATTGTGCCATT 540
541 ATTTTATGTCATCACTGGATAATATATTAGTGCTTAGTATCAGAAATAGTCCTTATGCT 600
601 TTGTGTTTTGAAGTTCCTAATGCAATGTTCTCTTTCTAGAAAAGGTGGACAAGTCTTATT 660
661 TTCCAGAGAAGATGACTTTTAACAGTTTTGAAGGAAGTAGAACTTTTGTACTTGCAGACA 720
SEQ. ID 10—1 M T F N S F E G T R T F V L A D T 17
721 CCAATAAGGATGAAGAATTTGTAGAAGAGTTTAATAGATTAAAAACATTTGCTAACTTCC 780
18 N K D E E F V E E F N R L K T F A N F P 37
781 CAAGTAGTAGTCTGTTTTCAGCATCAACATTGGCGCGAGCTGGGTTTCTTTATACCGGTG 840
38 S S S P V S A S T L A R A G F L Y T G E 57
841 AAGGAGACACCGTGCAATGTTTCAGTTGTCATGCGGCAATAGATAGATGGCAGTATGGAG 900
58 G D T V Q C F S C H A A I D R W Q Y G D 77
901 ACTCAGCTGTTGGAAGACACAGGAGAATATCCCCAAATTGCAGATTTATCAATGGTTTTT 960
78 S A V G R H R R I S P N C R F I N G F Y 97
961 ATTTTGAAAATGGTGCTGCACAGTCTACAAATCCTGGTATCCAAATGGCCAGTACAAAT 1020
98 F E N G A A Q S T N P G I Q N G Q Y K S 117
1021 CTGAAAACGTGTGTTGGGAAATAGAAATCCTTTTGCCCTGACAGGCCACCTGAGACTCATG 1080
118 E N C V G N R N P F A P D R P P E T H A 137
1081 CTGATTATCTCTTGAGAACTGGACAGGTTGTAGATATTTAGACACCATATACCCGAGGA 1140
138 D Y L L R T G Q V V D I S D T I Y P R N 157
1141 ACCCTGCCATGTGTAGTGAAGAAGCCAGATTGAAGTCATTTTCAGAACTGGCCGGACTATG 1200
158 P A M C S E E A R L K S F Q N W P D Y A 177
1201 CTCATTTAACCCCGAGAGAGTTAGCTAGTGCTGGCCTCTACTACACAGGGGCTGATGATC 1260
178 H L T P R E L A S A G L Y Y T G A D D Q 197
1261 AAGTGCAATGCTTTTGTGTGGGGGAAAACGAAAAATTGGGAACCCTGTGATCGTGCCT 1320
198 V Q C F C C G G K L K N W E P C D R A W 217
1321 GGTGAGAACACAGGAGACACTTTCCCAATTGCTTTTTTGTGTTTGGGCCGGAACGTTAATG 1380
218 S E H R R H F P N C F F V L G R N V N V 237
1381 TTCGAAGTGAATCTGGTGTGAGTTCTGATAGGAATTTCCCAAATTCAACAACTCTCCAA 1440
238 R S E S G V S S D R N F P N S T N S P R 257
1441 GAAATCCAGCCATGGCAGAAATATGAAGCACGGATCGTTACTTTTGAACATGGACATCCT 1500
258 N P A M A E Y E A R I V T F G T W T S S 277

Fig. 4

SUBSTITUTE SHEET (RULE 26)

12/42

1501 CAGTTAACAAGGAGCAGCTTGCAAGAGCTGGATTTTATG^{1,2}CTTTAGGTGAAGGCGATAAAG 1560
278 V N K E Q L A R A G F Y A L G E G D K V 297

1561 TGAAGTGCTTCCACTGTGGAGGAGGGCTCACGGATTGGAAGCCAAGTGAAGACCCCTGGG 1620
298 K C F H C G G G L T D W K P S E D P W D 317

1621 ACCAGCATGCTAAGTGCTACCCAG^{2,3}GGTGCAAATACCTATTGGATGAGAAGGGGCAAGAAT 1680
318 Q H A K C Y P G C K Y L L D E K G Q E Y 337

1681 ATATAAATAATATTCATTTAACCCATCCACTTGAGGAATCTTT^{3,4}GGAAGAACTGCTGAAA 1740
338 I N N I H L T H P L E E S L G R T A E K 357

1741 AAACACCACCGCTAACTAAAAAATCG^{4,5}ATGATACCATCTTCCAGAATCCTATGGTGAAG 1800
358 T P P L T K K I D D T I F Q N P M V Q E 377

1801 AAGCTATACGAATGGGATTTAGCTTCAAGGACCTTAAGAAAACAATGGAAGAAAAATCC 1860
378 A I R M G F S F K D L K K T M E E K I Q 397

1861 AAACATCCGGGAGCAGCTATCTATCACTTGAGGTCCTGATTGCAGATCTTGTGAGTGCTC 1920
398 T S G S S Y L S L E V L I A D L V S A Q 417

1921 AGAAAGATAATACGAGGATGAGTCAAGTCAAACCTCATTGCAGAAAG^{5,6}ACATTAGTACTG 1980
418 K D N T E D E S S Q T S L Q K D I S T E 437

1981 AAGAGCAGCTAAGGCGCTACAAGAGGAGAAGCTTTCCAAAATCTGTATGGATAGAAATA 2040
438 E Q L R R L Q E E K L S K I C M D R N I 457

2041 TTGCTATCGTTTTTTTTTCTTGTGGACATCTGGCCACTTGTAACAGTGTGCAGAAGCAG 2100
458 A I V F F P C G H L A T C K Q C A E A V 477

2101 TTGACAAATGTCCCATGTGCTACACCGTCATTACGTTCAACCAAAAAATTTTTATGTCTT 2160
478 D K C P M C Y T V I T F N Q K I F M S * 496

2161 AGTGGGGCACCACATGTTATGTTCTTCTTGTCTCTAATTGAATGTGTAATGGGAGCGAACT 2220
2221 TTAAGTAATCCTGCATTTGCATTCCATTAGCATCCTGCTGTTTCCAAATGGAGACCAATG 2280
2281 CTAACAGCACTGTTTCCGTCTAAACATTCAATTTCTGGATCTTTTCGAGTTATCAGCTGTA 2340
2341 TCATTTAGCCAGTGTTTTACTCGATTGAAACCTTAGACAGAGAAGCATTTTATAGCTTTT 2400
2401 CACATGTATATTGGTAGTACACTGACTTGATTTCTATATGTAAGTGAATTCATCACCTGC 2460
2461 ATGTTTCATGCCTTTTGCATAAGCTTAACAAATGGAGTGTCTGTATAAGCATGGAGATG 2520
2521 TGATGGAATCTGCCAATGACTTTAATTGGCTTATTGTAAACACGGAAAGAACTGCCCCA 2580
2581 CGCTGCTGGGAGGATAAAGATTGTTTATAGATGCTCACTTCTGTGTTTATGAGATTCTGCC 2640
2641 ATTTACTTGAATTTATTGGAGTTATAATGTACTTATATGATATTTCCGAA 2691

Fig. 4 (cont.)

13/42

SEQ. ID 11—1 TGGGAGTTCCCCGAGCCCTGGAGGAAAGCACCGCAGGTCTGAGCAGCCCTGAGCCGGGC 60
 61 AGGGTGGGGGCAGTGGCTAAGGCCTAGCTGGGGACGATTTAAAGGTATCGCGCCACCCAG 120
 121 CCACACCCACAGGCCAGGCGAGGGTGCCACCCCGGAGATCAGAGGTCATTGCTGGCGT 180
 181 TCAGAGCCTAGGAAGTGGGCTGCGGTATCAGCCTAGCAGTAAAACCGACCAGAAGCCATG 240
 241 CACAAACTACATCCCCAGAGAAAGACTTGTCCCTTCCCCTCCCTGTCATCTCACCATGA 300
 301 ACATGGTTCAAGACAGCGCCTTTCTAGCCAAGCTGATGAAGAGTGTGACACCTTTGAGT 360
 SEQ. ID 12—1 M V Q D S A F L A K L M K S A D T F E L 20
 361 TGAAGTATGACTTTTCCTGTGAGCTGTACCGATTGTCCACGTATTCAGCTTTTCCCAGGG 420
 21 K Y D F S C E L Y R L S T Y S A F P R G 40
 421 GAGTTCCTGTGTCAGAAAGGAGTCTGGCTCGTGTGGCTTTTACTACACTGGTGCCAATG 480
 41 V P V S E R S L A R A G F Y Y T G A N D 60
 481 ACAAGGTCAAGTGCTTCTGCTGTGGCCTGATGCTAGACAACCTGGAAACAAGGGGACAGTC 540
 61 K V K C F C C G L M L D N W K Q G D S P 80
 541 CCATGGAGAAGCACAGAAAGTTGTACCCAGCTGCAACTTTGTACAGACTTTGAATCCAG 600
 81 M E K H R K L Y P S C N F V Q T L N P A 100
 601 CCAACAGTCTGGAAGCTAGTCCTCGGCCTTCTCTTCCCTCCACGGCGATGAGCACCATGC 660
 101 N S L E A S P R P S L P S T A M S T M P 120
 661 CTTTGAGCTTTGCAAGTTCTGAGAATACTGGCTATTTCACTGGCTCTTACTCGAGCTTTC 720
 121 L S F A S S E N T G Y F S G S Y S S F P 140
 721 CCTCAGACCCTGTGAACCTCCGAGCAAATCAAGATTGTCTGCTTTGAGCACAAGTCCCT 780
 141 S D P V N F R A N Q D C P A L S T S P Y 160
 781 ACCACTTTGCAATGAACACAGAGAAGGCCAGATTACTCACCTATGAAACATGGCCATTGT 840
 161 H F A M N T E K A R L L T Y E T W P L S 180
 841 CTTTTCTGTACCAGCAAAGCTGGCCAAAGCAGGCTTCTACTACATAGGACCTGGAGATA 900
 181 F L S P A K L A K A G F Y Y I G P G D R 200
 901 GAGTGGCCTGCTTTGCGTGCGATGGGAAACTGAGCAACTGGGAACGTAAGGATGATGCTA 960
 201 V A C F A C D G K L S N W E R K D D A M 220
 961 TGTCAGAGCACCAGAGGCATTTCCCCAGCTGTCCGTTCTTAAAAGACTTGGGTGAGTCTG 1020
 221 S E H Q R H F P S C P F L K D L G Q S A 240
 1021 CTTCGAGATACACTGTCTCTAACCTGAGCATGCAGACACACGAGCCCGTATTAGAACAT 1080
 241 S R Y T V S N L S M Q T H A A R I R T F 260
 1081 TCTCTAACTGGCCTTCTAGTGCCTAGTTTCATCCCAGGAACTTGCAAGTGCGGGCTTTT 1140
 261 S N W P S S A L V H S Q E L A S A G F Y 280
 1141 ATTATACAGGACACAGTGATGATGTCAAGTGTTCCTGCTGTGATGGTGGGCTGAGGTGCT 1200
 281 Y T G H S D D V K C F C C D G G L R C W 300
 1201 GGGAACTCTGGAGATGACCCCTGGGTGGAACATGCCAAGTGGTTTCCAAAGTGTGAGTACT 1260
 301 E S G D D P W V E H A K W F P R C E Y L 320

Fig. 5
SUBSTITUTE SHEET (RULE 26)

14/42

1261 TGCTCAGAATCAAAGGCCAAGAATTTGTCAGCCAAGTTCAAGCTGGCTATCCTCATCTAC 1320
321 L R I K G Q E F V S Q V Q A G Y P H L L 340
1321 TTGAGCAGCTATTATCTACGTCTCAGACTCCCCAGAAGATGAGAATGCAGACGCAGCAATCG 1380
341 E Q L L S T S D S P E D E N A D A A I V 360
1381 TGCATTTTGGCCCTGGAGAAAGTTTCGGAAGATGTCGTCTATGATGAGCACGCCTGTGGTTA 1440
361 H F G P G E S S E D V V M M S T P V V K 380
1441 AAGCAGCCTTGGAATGGGCTTCAGTAGGAGCCTGGTGAGACAGACGGTTCAGCGGCAGA 1500
381 A A L E M G F S R S L V R Q T V Q R Q I 400
1501 TCCTGGCCACTGGTGAGAACTACAGGACCGTCTGACCTCGTTATAGGCTTACTCGATG 1560
401 L A T G E N Y R T V S D L V I G L L D A 420
1561 CAGAAGACGAGATGAGAGAGGAGCAGATGGAGCAGCGGCCGAGGAGGAGGAGTCAATG 1620
421 E D E M R E E Q M E Q A A E E E E S D D 440
1621 ATCTAGCACTAATCCGGAAGAACAATAATGGTGCTTTTCCAACATTTGACGTGTGTGACAC 1680
441 L A L I R K N K M V L F Q H L T C V T P 460
1681 CAATGCTGTATTGCCTCCTAAGTGCAAGGGCCATCACTGAACAGGAGTGAATGCTGTGA 1740
461 M L Y C L L S A R A I T E Q E C N A V K 480
1741 AACAGAAACCACACACCTTACAAGCAAGCACACTGATTGATACTGTGTTAGCAAAAGGAA 1800
481 Q K P H T L Q A S T L I D T V L A K G N 500
1801 AACTGCGCAACCTCATTGAGAACTCCCTTCGGGAAATTGACCCTGCGTTATACAGAG 1860
501 T A A T S F R N S L R E I D P A L Y R D 520
1861 ATATATTTGTGCAACAGGACATTAGGAGTCTTCCACAGATGACATTGCAGCTCTACCAA 1920
521 I F V Q Q D I R S L P T D D I A A L P M 540
1921 TGGAAGAACAGTTGCGGAACTCCAGGAGGAAAGAATGTGTAAAGTGTGTATGGACCGAG 1980
541 E E Q L R K L Q E E R M C K V C M D R E 560
1981 AGGTATCCATCGTGTTCATTCCCTGTGGCCATCTGGTGTGTGCAAGACTGCGCTCCCT 2040
561 V S I V F I P C G H L V V C K D C A P S 580
2041 CTCTGAGGAAGTGTCCATCTGTAGAGGGACCATCAAGGGCACAGTGCGCACATTTCTCT 2100
581 L R K C P I C R G T I K G T V R T F L S 600
2101 CCTGAACAAGACTAATGGTCCATGGCTGCAACTTCAGCCAGGAGGAAGTTCACTGTCACT 2160
*
2161 CCCAGCTCCATTCCGAACTTGAGGCCAGCCTGGATAGCACGAGACACCGCCAAACACACA 2220
2221 AATATAAACATGAAAACTTTGTCTGAAGTCAAGAATGAATGAATTACTTATATAATAA 2280
2281 TTTTAATTGGTTTCCTTAAAGTGCTATTTGTTCCCAACTCAGAAAATTGTTTCTGTAA 2340
2341 ACATATTTACATACTACCTGCATCTAAAGTATTCATATATTCATATATTCAGATGTCATG 2400
2401 AGAGAGGGTTTTGTTCTTGTTCCTGAAAAGCAGGGATTGCCTGCACTCCTGAAATTCCTCA 2460
2461 GAAAGATTTACAATGTTGGCATTTATGGTTTCAGAACTAGAATCTTCTCCCGTTGCTTTA 2520
2521 AGAACCAGGAGCAGATGTCCATGTGTTTATGTATAGAAATTCCTGTTATTTATTTGA 2580
2581 TGACATTTTAGGGATATGAAATTTTATAAAGAATTTGTGAGAAAAAGTTAATAAGCAA 2640
2641 CATAATTACCTCTTTTTTTTTTAAAGAAAAA 2676

Fig. 5 (cont.)

SUBSTITUTE SHEET (RULE 26)

15/42

SEQ. ID 13—1 AGTTATATAAAATACGAAGTTTCAAAAAGAAGGCTAGTGCAACAGAAAAGCTTTGCTAA 60
 61 AACAGATTCTTAGTTATTTGAGGTAACAAAAGAAAGCCATGTCTTGAATTGATTCGTTCT 120
 121 TAATTATAACAGACTTATAGTGGAAAGGGCCTTAAACACAGGCGGACTTTATAAAATGCA 180
 181 GTCTTAGGTTTATGTGCAAAATACTGTCTGTTGACCAGATGTATTCACATGATATATACA 240
 241 GAGTCAAGGTGGTGATATAGAAGATTTAACAGTGAGGGAGTTAACAGTCTGTGCTTTAAG 300
 301 CGCAGTTCCTTTACAGTGAATACTGTAGTCTTAATAGACCTGAGCTGACTGCTGCAGTTG 360
 361 ATGTAACCCACTTTAGAGAATACTGTATGACATCTTCTTAAGGAAAACCAGCTGCAGAC 420
 421 TTCACTCAGTTCCTTTTCATTTTCATAGGAAAAGGAGTAGTTTCAGATGTATGTTTAAGTCC 480
 481 TTATAAGGGAAGAGCCTGAATATATGCCCTAGTACCTAGGCTTCATAACTAGTAATAA 540
 541 GAAGTTAGTTATGGGTAAATAGATCTCAGGTTACCCAGAAGAGTTCATGTGACCCCCAAA 600
 601 GAGTCCTAACTAGTGTCTTGGCAAGTGACAGATTGTCTCTGTGAGGGTGTCAATTCAC 660
 661 CAGTCCAAGCAGAAGACAATGAATCTATCCAGTCAGGTGTCTGTGGTGGAGATCTAGTGT 720
 721 CCAAGTGGTGAGAACTTCATCTGGAAGTTTAAGCGGTGAGAAATACTATTACTACTCAT 780
 1 M 1

781 GGACAAAAGTGTCTCCAGAGACTCGGCCAAGGTACCTTACACCAAAAAGCTTAAACGTAT 840
 SEQ. ID 14—2 D K T V S Q R L G Q G T L H Q K L K R I 21

841 AATGGAGAAGAGCACAATCTTGTCAAATTGGACAAAGGAGAGCGAAGAAAAATGAAGTT 900
 22 M E K S T I L S N W T K E S E E K M K F 41

901 TGAAGTTTCGTGTGAAGTCTACCGAATGTCTACATATTCAGCTTTTCCAGGGGAGTTCC 960
 42 D F S C E L Y R M S T Y S A F P R G V P 61

961 TGTCTCAGAGAGGAGTCTGGCTCGTGGCTTTTATTATACAGGTGTGAATGACAAAGT 1020
 62 V S E R S L A R A G F Y Y T G V N D K V 81

1021 CAAGTGCTTCTGCTGTGGCCTGATGTTGGATAACTGGAAACAAGGGGACAGTCCTGTGTA 1080
 82 K C F C C G L M L D N W K Q G D S P V E 101

1081 AAAGCACAGACAGTTCTATCCCAGCTGCAGCTTTGTACAGACTCTGCTTTTCAGCCAGTCT 1140
 102 K H R Q F Y P S C S F V Q T L L S A S L 121

1141 GCAGTCTCCATCTAAGAATATGTCTCCTGTGAAAAGTAGATTTCACATTCGTCACCTCT 1200
 122 Q S P S K N M S P V K S R F A H S S P L 141

1201 GGAACGAGGTGGCATTCACTCCAACCTGTGCTCTAGCCCTCTTAATTCTAGAGCAGTGA 1260
 142 E R G G I H S N L C S S P L N S R A V E 161

1261 AGACTTCTCATCAAGGATGGATCCCTGCAGCTATGCCATGAGTACAGAAGAGGCCAGATT 1320
 162 D F S S R M D P C S Y A M S T E E A R F 181

1321 TCTTACTTACAGTATGTGGCCTTTAAGTTTCTGTACCAGCAGAGCTGGCCAGAGCTGG 1380
 182 L T Y S M W P L S F L S P A E L A R A G 201

1381 CTTCTATTACATAGGGCCTGGAGACAGGGTGGCCTGTTTTGCCTGTGGTGGGAAAGTGA 1440
 202 F Y Y I G P G D R V A C F A C G G K L S 221

1441 CAACTGGGAACCAAAGGATGATGCTATGTCAGAGCACCGCAGACATTTTCCCCACTGTCC 1500
 222 N W E P K D D A M S E H R R H F P H C P 241

1501 ATTTCTGAAAATACTTCAGAAACACAGAGGTTTAGTATATCAAATCTAAGTATGCAGAC 1560
 242 F L E N T S E T Q R F S I S N L S M Q T 261

Fig. 6

SUBSTITUTE SHEET (RULE 26)

16/42

1561 ACACTCTGCTCGATTGAGGACATTTCTGTACTGGCCACCTAGTGTTCCTGTTTCAGCCCGA 1620
 262 H S A R L R T F L Y W P P S V P V Q P E 281
 1621 GCAGCTTGCAAGTGCTGGATTCTATTACGTGGATCGCAATGATGATGTCAAGTGCTTTTG 1680
 282 Q L A S A G F Y Y V D R N D D V K C F C 301
 1681 TTGTGATGGTGGCTTGAGATGTTGGGAACCTGGAGATGACCCCTGGATAGAACACGCCAA 1740
 302 C D G G L R C W E P G D D P W I E H A K 321
 1741 ATGGTTTCCAAGGTGTGAGTTCTTGATACGGATGAAGGGTCAGGAGTTGTTGATGAGAT 1800
 322 W F P R C E F L I R M K G Q E F V D E I 341
 1801 TCAAGCTAGATATCCTCATCTTCTTGAGCAGCTGTTGTCCACTTCAGACACCCCAGGAGA 1860
 342 Q A R Y P H L L E Q L L S T S D T P G E 361
 1861 AGAAAATGCTGACCCCTACAGAGACAGTGGTGCATTTTGGCCCTGGAGAAAAGTTCGAAAGA 1920
 362 E N A D P T E T V V H F G P G E S S K D 381
 1921 TGTCGTCATGATGAGCAGCCTGTGGTTAAAGCAGCCTTGGAAATGGGCTTCAGTAGGAG 1980
 382 V V M M S T P V V K A A L E M G F S R S 401
 1981 CCTGGTGAGACAGACGGTTCAGCGGCAGATCCTGGCCACTGGTGAGAACTACAGGACCGT 2040
 402 L V R Q T V Q R Q I L A T G E N Y R T V 421
 2041 CAATGATATTGTCTCAGTACTTTTGAATGCTGAAGATGAGAGAAGAGAAGAGGAGAAGGA 2100
 422 N D I V S V L L N A E D E R R E E E K E 441
 2101 AAGACAGACTGAAGAGATGGCATCAGGTGACTTATCACTGATTCCGAAGAATAGAATGGC 2160
 442 R Q T E E M A S G D L S L I R K N R M A 461
 2161 CCTCTTTCAACAGTTGACACATGTCTTCCTATCCTGGATAATCTTCTTGAGGCCAGTGT 2220
 462 L F Q Q L T H V L P I L D N L L E A S V 481
 2221 AATTACAAAACAGGAACATGATATTATTAGACAGAAAACACAGATACCCTTACAAGCAAG 2280
 482 I T K Q E H D I I R Q K T Q I P L Q A R 501
 2281 AGAGCTTATTGACACCGTTTTAGTCAAGGGAAATGCTGCAGCCAACATCTTCAAAAACCTC 2340
 502 E L I D T V L V K G N A A A N I F K N S 521
 2341 TCTGAAGGAAATTGACTCCACGTTATATGAAAACCTATTGTTGGAAAAGAATATGAAGTA 2400
 522 L K E I D S T L Y E N L F V E K N M K Y 541
 2401 TATTCCAACAGAAGACGTTTCAGGCTTGTCAATTGGAAGAGCAGTTGCGGAGATTACAAGA 2460
 542 I P T E D V S G L S L E E Q L R R L Q E 561
 2461 AGAACGAACCTGCAAAGTGTGTATGGACAGAGAGGTTTCTATTGTGTTTCATTCCGTGTGG 2520
 562 E R T C K V C M D R E V S I V F I P C G 581
 2521 TCATCTAGTAGTCTGCCAGGAATGTGCCCTTCTCTAAGGAAGTGCCCCATCTGCAGGGG 2580
 582 H L V V C Q E C A P S L R K C P I C R G 601
 2581 GACAATCAAGGGGACTGTGCGCACATTTCTCTCATGAGTGAAGAATGGTCTGAAAGTATT 2640
 602 T I K G T V R T F L S * 612

Fig. 6 (cont.)

SUBSTITUTE SHEET (RULE 26)

17/42

2641 GTTGGACATCAGAAGCTGTCAGAACAAAGAATGAACTACTGATTTTCAGCTCTTCAGCAGG 2700
2701 ACATTCTACTCTCTTTC AAGATTAGTAATCTTGCTTTATGAAGGGTAGCATTGTATATTT 2760
2761 AAGCTTAGTCTGTTGCAAGGGAAGGTCTATGCTGTTGAGCTACAGGACTGTGTCTGTTCC 2820
2821 AGAGCAGGAGTTGGGATGCTTGCTGTATGTCCTTCAGGACTTCTTGGATTTGGAATTTGT 2880
2881 GAAAGCTTTGGATTCAAGGTGATGTGGAGCTCAGAAATCCTGAAACCAGTGGCTCTGGTAC 2940
2941 TCAGTAGTTAGGGTACCCTGTGCTTCTTGGTGCTTTTCCTTTCTGGAAAATAAGGATTTT 3000
3001 TCTGCTACTGGTAAATATTTTCTGTTTGTGAGAAATATATTAAAGTGTTCCTTTTAAAGG 3060
3061 CGTGCATCATTGTAGTGTGTGCAGGGATGTATGCAGGCAAAACACTGTGTATATAATAAA 3120
3121 TAAATCTTTTAAAAAGTGTA AAAAAAAAAA 3151

Fig. 6 (cont.)

18/42




Fig. 7




Fig. 8

19/42




Fig. 9

20/42




Fig. 10A


MENADIONE (20 μ M), CHO CELLS: 24hr SURVIVAL

Fig. 10B

21/42


STAUROSPOURINE (1 μ M), RAT-1 CELLS, 24 HOUR SURVIVAL

Fig. 10C


MENADIONE (10 μ M), RAT-1 CELLS, 18 HOUR SURVIVAL

Fig. 10D

22/42




Fig. 11




Fig. 12

23/42




Fig. 13




Fig. 14

24/42




Fig. 15A

Fig. 15B

25/42




Fig. 16A




Fig. 16B

26/42




Fig. 17

27/42




Fig. 18

28/42




Fig. 19

29/42

INFLUENCE OF TAXOL ON DNA FRAGMENTATION IN
CISPLATIN-SENSITIVE (OV2008) AND -RESISTANT (C13)
HUMAN OVARIAN EPITHELIAL CANCER




Fig. 20

30/42

SELECTIVE INFLUENCE OF CISPLATIN ON DNA FRAGMENTATION
IN SENSITIVE (OV2008) AND -RESISTANT (C13)
HUMAN OVARIAN EPITHELIAL CANCER




Fig. 21

31/42

EFFECTS OF TAXOL ON XIAP AND HIAP-2 PROTEIN CONTENT IN
CISPLATIN-RESISTANT (C13) AND -SENSITIVE (OV2008)
HUMAN OVARIAN EPITHELIAL CANCER CELLS IN VITRO




Fig. 22

32/42

INFLUENCE OF TAXOL and TGF β ON HIAP-2 mRNA ABUNDANCE
IN CISPLATIN-SENSITIVE (OV2008) AND -RESISTANT (C13)
HUMAN EPITHELIAL CANCER CELLS IN VITRO




Fig. 23A




Fig. 23B

33/42

INFLUENCE OF TGF β ON XIAP PROTEIN EXPRESSION AND DNA
FRAGMENTATION IN CISPLATIN-SENSITIVE (OV2008) AND -RESISTANT (C13)
HUMAN OVARIAN EPITHELIAL CANCER CELLS IN VITRO




Fig. 24A




Fig. 24B

34/42








Fig. 25

35/42



36/42




Fig. 27A



Fig. 27B




Fig. 27C

37/42




Fig. 28A




Fig. 28B

38/42




Fig. 29A

39/42




Fig. 29B

40/42




Fig. 30A








Fig. 30B

41/42






Fig. 32A




Fig. 32B



Fig. 32C




Fig. 32D

INTERNATIONAL SEARCH REPORT

Int'l. Application No.

PCT/IB 98/00781

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K38/17 A61K31/70 A61K39/395 C12N15/11 C12Q1/68
G01N33/50 G01N33/574 A01K67/027 C12N15/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K C12N C12Q G01N A01K C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| A | P. LISTON ET AL.: "SUPPRESSION OF APOPTOSIS IN MAMMALIAN CELLS BY NAIP AND A RELATED FAMILY OF IAP GENES." NATURE, vol. 379, 25 January 1996, pages 349-353, XP002032296 LONDON GB see the whole document --- -/-- | 1-104 |

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

20 August 1998

Date of mailing of the international search report

27/08/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ryckebosch, A

INTERNATIONAL SEARCH REPORT

Int. Patent Application No
PCT/IB 98/00781

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|---|-----------------------|
| P,X | <p>WO 97 06255 A (UNIVERSITY OF OTTAWA) 20 February 1997 cited in the application see page 4, line 11 - line 28; claims 1-16,49,74,78,85-94; figures 1-4 see page 40, line 27 - page 43, line 21 see page 50, line 10 - page 52, line 8 see page 6, line 24 - page 7, line 9 see page 11, line 21 - line 29 see page 12, line 7 - page 13, line 9 see page 14, line 31 - page 15, line 9 see page 35, line 20 - line 23 see page 39, line 33 - page 40, line 2 ----</p> | 1-104 |
| P,X | <p>WO 97 26331 A (UNIVERSITY OF OTTAWA) 24 July 1997 cited in the application see page 3, line 27 - page 4, line 8 see page 37, line 9 - line 11; claims 18-20,62-78 see page 23, paragraph 4 - page 24, line 2 see page 25, line 9 - page 27, line 4 see page 28, paragraph 4 see page 7, paragraph 3 see page 8, line 12 - line 20 see page 8, line 28 - page 9, line 6 see page 10, line 4 - line 11 ----</p> | 1-104 |
| E | <p>WO 98 22131 A (UNIVERSITY OF OTTAWA) 28 May 1998 cited in the application see page 3, line 20 - page 4, line 3; claims 16,17,20-22,24-27,29-35 see page 10, line 12 - line 14 see page 15, line 22 - line 24 see page 18, line 27 - page 19, line 2 see page 21, line 13 - line 14 see page 7, line 4 - line 6 -----</p> | 1-104 |

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB 98/00781

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-19 (as far as being related to an in vivo method) and 74-82 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 98/00781

| Patent document cited in search report | Publication date | Patent family member(s) | Publication date |
|---|---------------------|----------------------------|---------------------|
| WO 9706255 A | 20-02-1997 | EP 0837939 A | 29-04-1998 |
| WO 9726331 A | 24-07-1997 | AU 1614997 A | 11-08-1997 |
| | | CA 2215793 A | 24-07-1997 |
| | | EP 0815231 A | 07-01-1998 |
| WO 9822131 A | 28-05-1998 | NONE | |

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☒ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.